DURNAL DE PHYSIQUE

us le patronage de la Société Française de Physique oncours du ational de la Recherche Scientifique

Europhysics Journal

Luiophysics douine

, 1981

ISSN 0302-0738

x Général

lications 1981





JOURNAL DE PHYSIQUE

Tome 42 — 1981

PHYSIQUE MATHÉMATIQUE

dit (Ph.). — Utilisation des matrices cycliques pour l'étude des cristaux parfaits : discussion générale	903	Fröman (N.) and Fröman (P. O.). — Generalization of the Fermi-Segrè formula.	1491
	903	Fröman (P. O.). — Voir Fröman (N.)	
Ell (R.) and Mehta (M. L.). — Sequence of invariants for	1102	Mehta (M. L.). — Voir Ball (R.)	
knots and links	1193	Muller (J.), Esling (C.) and Bunge (H. J.). — An inversion	-
Inge (H. J.). — Voir Muller (J.)	161	formula expressing the texture function in terms of	
Fing (C.). — Voir Muller (J.)	161	angular distribution functions	161
MÉCANIQUE	E, HY	DRODYNAMIQUE	
Akerson (B. J.) and Clark (N. A.). — Dynamic light scat-		Pomeau (Y.) and Manneville (P.).— Wavelength selection	
tering at low rates of shear	929	in axisymmetric cellular structures	1067
B'gé (P.). — Voir Dubois (M.)	167	Pomeau (Y.) and Zaleski (S.). — Wavelength selection in one-	in la
Bitgen (W.). — Voir Sadik (J.)	915	dimensional cellular structures	515
Crk (N. A.). — Voir Ackerson (B. J.)	929	Rothen (F.). — Voir Sadik (J.)	915
Dois (M.) et Bergé (P.). — Instabilités de couche limite dans un fluide en convection. Evolution vers la turbu-		Sadik (J.), Rothen (F.), Bestgen (W.) and Dubois-Violette (E.). — Theoretical study of the instability of a homeo-	
lence	167	tropic nematic submitted to an elliptical shear	915
Dois-Violette (E.). — Voir Sadik (J.)	915	Wu (F.) et Perio (P.). — La phase en acoustique musicale. II.	
Vnneville (P.). — Voir Pomeau (Y.)	1067	Le rayonnement des instruments à vent	627
Pio (P.). — Voir Wu (F.)	627	Zaleski (S.). — Voir Pomeau (Y.).	515
MÉCANI	OUE	STATISTIQUE	
	-	team (B.) Seriou ellis more tales de deseguera	
A stuey (A.) and Jancovici (B.). — On the classical two-		correlation functions of number fluctuations in simple	
dimensional one-component Coulomb plasma	1	fluids with radial many-body interactions	1373
Bc (P.). — Voir Villain (J.)	657	Knoll (A.), Schäfer (L.) and Witten (T. A.). — The thermo-	
Baux (R.) and De Seze (L.). — Exact solution of the 2D		dynamic scaling function of a polymer solution	767
« brickwork » Ising model with second neighbour	201	Lajzerowicz (J.). — Voir Vallade (M.)	1505
interactions in the horizontal direction	371	Lubensky (T. C.), Isaacson (J.) and Obukhov (S. P.). — Field	1501
Cnielowski (W.). — Voir Knast (K.).	1373	theory for ARB ₂ branched polymers	1591
Doud (M.) and Joanny (J. F.). — Conformation of branched	1359	Lubensky (T. C.) and Isaacson (J.). — Field theory and polymer size distribution for branched polymers	175
polymers	1339	Lubensky (T. C.). — Voir Dunn (S. G.)	1201
frontiers of Potts ferromagnets	1515	Luck (J. M.) Voir Drouffe (J. M.).	653
Di Cloizeaux (J.). — Polymers in solutions: principles and	1000	Moussa (P.). — Voir Jullien (R.)	1075
applications of a direct renormalization method	635	Obukhov (S. P.) Voir Lubensky (T. C.)	1591
DSeze (L.). — Voir Bidaux (R.)	371	Pfeuty (P.). — Voir Jullien (R.)	1075
Druffe (J. M.), Luck (J. M.) and Zuber (J. B.). — The step		Pokrovsky (V. L.). — Splitting of commensurate-incom-	
free-energy and the behaviour at the roughening tran-		mensurate phase transition	
sition of lattice gauge theories	653	Ruskin (H. J.). — Voir Duarte (J. A. M. S.)	1585
Dirte (J. A. M. S.) and Ruskin (H. J.). — The branching of	1505	Schäfer (L.). — Voir Knoll (A.)	767
real lattice trees as dilute polymers	1585	Tsallis (C.). — Voir De Magalhães (A. C. N.)	1515
Dm (S. G.) and Lubensky (T. C.). — Gauge-dependent		q-state Potts model: weak-coupling expansion	793
critical properties of the nematic to smectic-A phase transition in the 1/N-expansion	1201	Uzelac (K.). — Voir Jullien (R.).	1075
Faily (F.). — Polymer excluded volume exponent v in three	1201	Vallade (M.) et Lajzerowicz (J.). — Transition rugueuse et	2010
dimensions by direct renormalization	189	localisation pour une singularité linéaire dans un espace	
Isicson (J.). — Voir Lubensky (T. C.)	1591	à deux ou trois dimensions	1505
Isicson (J.). — Voir Lubensky (T. C.)	175	Villain (J.) and Bak (P.). — Two-dimensional Ising model	
Jacovici (B.). — Voir Alastuey (A.)	1	with competing interactions: floating phase, walls and	
Jenny (J. F.). — Voir Daoud (M.)	1359	dislocations	657
Jtien (R.), Uzelac (K.), Pfeuty (P.) and Moussa (P.).		Witten (T. A.). — Voir Knoll (A.).	767
The Yang-Lee edge singularity studied by a four-level	1075	Zinn-Justin (J.). — Analysis of high temperature series of	793
quantum renormalization-group blocking method	1075	the spin S Ising model on the body-centred cubic lattice.	783 653
Kest (K.) and Chmielowski (W.) Study of the higher-order		Zuber (J. B.). — Voir Drouffe (J. M.)	023

PHYSIQUE NUCLÉAIRE

Chaudhury (M. L.) and Sen (D. K.). — Effect of non-local	Plo (M.). — Voir Dupuis (P.)	16
barrier and relative intensities of alpha-spectra in rare-	Pomorski (K.) and Hofmann (H.). — On the dynamics of	
earth nuclei	fission as a dissipative process	3
Dasgupta (S. S.). — Voir Slobodrian (R. J.)	Rioux (C.). — Voir Slobodrian (R. J.)	
Dupuis (P.), Leccia (F.), Mennrath (P.), Morales (J.) et	Roy (R.). — Voir Slobodrian (R. J.)	
Plo (M.). — Etude de la décroissance β^+ de ³³ Cl 1603	Sen (D. K.). — Voir Chaudhury (M. L.)	
Gary (S.). — Voir Volant (C.)	Silvestre-Brac (B.) and Liotta (R. J.). — Full nuclear field	
Hofmann (H.). — Voir Pomorski (K.)	theory treatment of two-particle-one-hole-excitations.	7
Issa (A.). — Pairing quadrupolaire et propriétés des états	Slobodrian (R. J.), Dasgupta (S. S.), Rioux (C.), Lahlou (F.)	
(O ⁺) des noyaux des terres rares	et Roy (R.). — Déviations par rapport à l'espace des	
Lahlou (F.). — Voir Slobodrian (R. J.).	phases et forces dans les réactions $p + d \rightarrow p + p + n$	
Leccia (F.). — Voir Dupuis (P.)	et $d + p \rightarrow p + p + n$ très près du seuil	
Liotta (R. J.). — Voir Silvestre-Brac (B.)	Volant (C.) and Gary (S.). — Complete fusion of the	
Mennrath (P.). — Voir Dupuis (P.)	$^{14}N + ^{16}O$ and $^{15}N + ^{16}O$ systems	
Morales (J.). — Voir Dupuis (P.).		

PHYSIQUE ATOMIQUE ET MOLÉCULAIRE

Alamichel (C.). — Voir Bensari-Zizi (N.)	209
Alamichel (C.). — Voir Bensari-Zizi (N.)	827
André (J.). — Voir Vedel (F.)	-1611
André (J.). — Voir Vedel (M.)	541
Barrat (JP.). — Voir Lémery (H.)	1081
Bensari-Zizi (N.) and Alamichel (C.). — Study of the v_1 - v_3	
infrared hot band of CH ₃ Cl ³⁵	209
Bensari-Zizi (N.), Alamichel (C.) and Guelachvili (G.).	
Study of the anharmonic resonance between $v_2 + v_4^{\pm 1}$	
and $v_2 + 3 v_6^{\pm 1}$ and between $v_3 + v_4^{\pm 1}$ and $v_3 + 3 v_6^{\pm 1}$	
infrared bands of CH ₃ Cl ³⁵	827
Blain (M.). — Voir Brigot (N.).	979
Bliman (S.), Dousson (S.), Geller (R.), Jacquot (B.) et Van	
Houtte (D.). — Sections efficaces d'échange de charges	
dans la collision des ions multichargés O ^{q +} et N ^{q +} sur	
le deutérium moléculaire	399
Bliman (S.), Dousson (S.), Geller (R.), Jacquot (B.) et Van	
Houtte (D.). — Capture multiple d'électrons dans la	
collision des ions multiplement chargés d'argon sur	
l'argon	
Bliman (S.), Dousson (S.), Jacquot (B.) and Van Houtte (D.).	
Charge exchange cross sections of argon ions colliding	
with rare gas targets at keV energies	1387
Bloch (D.). — Voir Ducloy (M.)	711
Bordé (Ch. J.), Camy (G.), Decomps (B.), Descoubes (JP.)	
and Vigué (J.). — High precision saturation spectroscopy	
of ¹²⁷ I ₂ with argon lasers at 5 145 Å and 5 017 Å : I	
Mean resonances	1393
Bordé (J.). — Voir Palma (M. L.)	1239
Bouchiat (M. A.), Coblentz (A.), Guena (J.) and Pottier (L.).	
Can imperfect light polarization mimic parity violation	
in Stark experiments on forbidden M1 transitions?	985
Boulet (C.) and Rosenberg (A.). — Pressure induced shifts	HPAR.
in HCl/HF and HF/HCl systems	203
Bousquet (C.). — Voir Bras (N.).	215
Bras (N.) et Bousquet (C.). — Etude du satellite « bleu » de la	
raie 1 849 Å du mercure autoélargie. Mise en évidence	
d'une variation du moment de transition avec la distance	
internucléaire	215
Brigot (N.), Cohen de Lara (E.) et Blain (M.). — Calcul	
ab initio de l'interaction d'une molécule d'azote avec	
un ion Na ⁺ . Comparaison avec les données expérimen-	
tales du spectre infrarouge de N ₂ adsorbés dans la	
zéolithe NaA	979
Broyer (M.). — Voir Vigué (J.)	0.000
Camy (G.). — Voir Bordé (Ch. J.)	-1393

Carré (M.), Zgainski (A.), Gaillard (M.), Nouh (M.) et
Lombardi (M.). — Détermination des populations rela-
tives des sous-niveaux magnétiques du niveau 4 1D
de HeI excité par impact d'ions lourds
Chaillat-Negrel (S.). — Voir Vedel (M.)
Coblentz (A.) Voir Bouchiat (M. A.)
Cohen de Lara (E.). — Voir Brigot (N.).
Decomps (B.). — Voir Bordé (Ch. J.) 1.
Delsart (C.), Keller (JC.) and Kaftandjian (V. P.). — Inves-
tigation of the dynamic Stark effect of a $J = 0 \leftrightarrow J = 1$ optical transition
Descoubes (JP.). — Voir Bordé (Ch. J.)
Di Lauro (C.) et Lattanzi (F.). — Angles de phases dans les
éléments matriciels en spectroscopie ro-vibronique
Dousson (S.). — Voir Bliman (S.)
Ducloy (M.) and Bloch (D.). — Theory of degenerate four-
wave mixing in resonant Doppler-broadened systems.
I. Angular dependence of intensity and lineshape of
phase-conjugate emission
Gaillard (M.). — Voir Carré (M.)
Gaillard (M.). — Voir Carré (M.). Geller (R.). — Voir Bliman (S.). 399-
Goulet (P.). — Voir Verry (C.)
Grandin (J. P.) et Husson (X.). — Etude de la dépendance
en température des sections efficaces de dépolarisation
par collision des niveaux 2p de l'atome d'argon
Guelachvili (G.). — Voir Bensari-Zizi (N.).
Guéna (J.). — Voir Bouchiat (M. A.).
Hamel (J.). — Voir Lémery (H.)
Husson (X.). — Voir Grandin (J. P.).
Husson (X.). — Voir Grandin (J. P.). Jacquot (B.). — Voir Bliman (S.). 399-705-1 Jeannet (J. C.). — Voir Perrin (D.). 1
Jeannet (J. C.). — Voir Perrin (D.)
Jurek (R.). — Voir Verry (C.)
Attachandian (v. 1.). Von Delsatt (C.)
Kazantsev (A. P.), Surdutovich (G. I.) and Yakovlev (V. P.). —
The motion of atoms and molecules in a resonance
light field
Keller (JC.). — Voir Delsart (C.)
Lamalle (B.). — Voir Verry (C.). Lattanzi (F.). — Voir Di Lauro (C.)
Lattanzi (F.). — Voir Di Lauro (C.)
Leboucher (E.). — Aspect dynamique des collisions dans le
calcul des profils de raies atomiques. Application à la
recherche des potentiels interatomiques Hg(6 ³ P ₁)-gaz
rares lourds
Lehmann (J. C.). — Voir Vigué (J.)
Lémery (H.), Hamel (J.) et Barrat (JP.). — Mesure de la
structure hyperfine des niveaux 3 ¹ D ₂ , 4 ¹ D ₂ et 5 ¹ D ₂
de l'hélium 3

ns
de 1611
391-1611
n-
de
. 541
es
223
al
te.
937
al
te. 0-
949
al
te. 961
. 1393
1231
. 235
gy
1421 1525 1421
1421
1421 1525 1421 1525
1421 1525 1421 1525
1421 1525 1421 1525 ee- 723
1421 1525 1421 1525 ee- 723
1421 1525 1421 1525 ee- 723
1421 1525 1421 1525 ee- 723
1421 1525 1421 1525 723
1421 1525 1421 1525 723
1421 1525 1421 1525 ee- 723
1421 1525 1421 1525 ee- 723
1421 1525 1421 1525 ee- 723
1421 1525 1421 1525 ee- 723
1421 1525 1421 1525 723 723 723 1279 1651 353 605 1129 129
1421 1525. 1421 1525 1525 723 723 1279 1651 353 605 1129 593 113
1421 1525. 1421 1525 ee- 723 1279 dd on 445 991 1651 353 605 1129 ee- dd 593 1157 293
1421 1525. 1421 1525 ee- 723 1279 dd 1279 dd 1651 353 605 1129 e- dd 593 1157 293
1421 1525 1421 1525 ee- 723 dd 1279 dd 1279 dd 1651 353 1055 1129 ee- dd 593 1157 293 345
1421 1525. 1421 1525 ee- 723 1279 dd 1279 dd 1651 353 605 1129 e- dd 593 1157 293
the state of the s

Blanchard (C.). — Voir Deville (A.)	1641	Destrade (C.). — Voir Levelut (A. M.)	
Bloemen (E.) and Garland (C. W.). — Calorimetric investi-		Deveaud (B.). — Voir Picoli (G.)	133
gation of phase transitions in butyloxybenzylidene		Deville (A.), Arzoumanian (C.), Gaillard (B.), Blanchard (C.),	
heptylaniline (40.7)		Jamet (J. P.) and Maletta (H.). — Electron resonance	1641
Boulon (G.). — Voir Moine (B.)	499	in the insulating spin glass Eu _{0.4} Sr _{0.6} S	1641
Bourcet (J. C.). — Voir Moine (B.)	499	Dondos (A.). — Voir Izumi (Y.)	353
Bourdon (L.), Kléman (M.), Lejcek (L.) and Taupin (D.).		Dorner (B.). — Voir Levelut (A. M.)	
On static helical instabilities of screw dislocations in a	261		165
SmA phase and on their consequence on plasticity	261	Draman (C.). — Voir Haydar (A.)	092
Briggs (A.), Monceau (P.), Nunez-Regueiro (M.), Ribault	10	Dreyfus (JM.) and Guyon (E.). — Convective instabilities	283
(M.) and Richard (J.). — Superconductivity of NbSe ₃	1453	in nematics caused by an elliptical shear Dreyfus (JM.) and Pieranski (P.). — Distorsion waves and	201
Brochard (F.). — Polymer networks swollen by a homo-		phase slippage in nematics	459
polymer solution		Dreysse (H.) et Riedinger (R.). — Structure électronique auto-	433
Brochard (F.). — Voir Joanny (J. F.).	1145	cohérente d'une surface traitée en liaisons fortes	437
Broniatowski (A.). — Charged grain boundaries in germa-	741	Dubois-Violette (E.). — Voir Pieranski (P.)	53
nium		Dujardin (F.) and Decruppe (J. P.). — Attempt to determine	
Brot (C.). — Voir Tresser (C.).	427	the band parameters of graphite by a theoretical calcu-	
Brundle (C. R.). — Voir Schlapbach (L.)	1025	lation	1167
Brunet (M.). — Voir Perez (A.)	1559	Dupas (C.), Le Dang (K.), Renard (J. P.), Veillet (P.), Miran-	
Buckmaster (H. A.). — Voir Malhotra (V. M.)	1053	day (J. P.) and Jacoboni (C.) Susceptibi-	
Buevoz (J. L.). — Voir Meyer (C.)	003	lity and N.M.R. study of the vitrous fluorides	
Bunning (J. D.), Faber (T. E.) and Sherrell (P. L.). — The		$Pb_2MnFe_{1-x}Ga_xF_9$ and $Pb_2Fe-Mn_{1-x}Zn_xF_9$	1345
Franck constants of nematic 5CB at atmospheric pressure	1175	Durand (G.) Voir Martinot-Lagarde (Ph.)	
	1113	Durand (G.). — Voir Bartolino (R.)	1445
Cabane (B.). — Structure of the water/surfactant interface		Duval (Ph.), Peyre (H.) et Henry (L.). — Etude expérimen-	
in micelles : an NMR study of SDS micelles labelled	847	tale des franges d'interférences au bord des images	
with paramagnetic ions Caroli (B.). — Voir Abraham-Ibrahim (S.).		électroniques inélastiques et quasi élastiques	751
	1137	Faber (T. E.). — Voir Bunning (J. D.)	
Caroli (C.). — Voir Abraham-Ibrahim (S.). Cavagnat (R.). — Voir Besnard (M.).	1137 593	Fabre (C.). — Voir Barbarin (F.)	1183
Chakraverty (B. K.). — Bipolarons and superconductivity	1351	Faure (P.). — Dynamique cristalline avec charges tenso-	- 43
Chan (W. K.) and Webb (W. W.). — Observation of elemen-	1331	rielles. Application à la glace hexagonale	579
tary edge dislocations in phospholipid multilayers and		Feigelman (M.) and Pokrovsky (V.). — Long-range order	
of their annealing as a determination of the permeation		and diffraction of X-ray waves on multilayered crys-	124
coefficient	1007	talline films	125
Charvolin (J.). — Voir Hendrikx (Y.)	1427	Ferrari (G.). — Voir Porteseil (J. L.)	125
Chausse (J. P.). — Voir Barbarin (F.)	1183	Findeisen (M.). — Voir Limmer (St.)	1665
Chervin (J. C.). — Voir Piccioli (N.)	1129	Flapper (S. D. P.), Vertogen (G.) and Leenhouts (F.). — On the elastic constants of nematics; a comparison	
Chevalier (J.), Baixeras (J.) et Andro (P.) Etude expéri-		between experiment and molecular theory	1647
· mentale de différents modes de conduction par sauts		and still on extractively during the two promition my turns and	
dans un alliage désordonné Nb-N	405	Flechon (J.). — Voir Machizaud (F.)	97
Chizhikov (V. I.), Panyutin (V. L.), Ponedelnikov (B. E.) et		Flores (F.). — Voir Louis (E.)	
Rozenson (A. E.). — Energie de liaison des excitons		Gaillard (B.). — Voir Deville (A.). Galland (D.). — Voir Picoli (G.).	164:
directs dans $CdGa_2(S_xSe_{1-x})_4$	1003	Garland (C. W.). — Voir Bloemen (E.)	129
Chizhikov (V. I.), Panyutin (V. L.), Ponedelnikov (B. E.) et		Gasparoux (H.). — Voir Levelut (A. M.).	14
Rozenson (A. E.). — Dependance en température de la		Germain (J. P.). — Voir Barbarin (F.)	1183
largeur de bande interdite dans CdIn ₂ Se ₄	1039	Gervais (F.). — Voir Quittet (A. M.).	493
Chouteau (G.) and Veyret-Jeandey (Cl.). — Metamagnetism	MILLIA	Giraud (G.). — Voir Guyon (E.).	1553
in tanol suberate	1441	Glogarová (M.). — Anchoring effect of triglycine sulphate	
Clerc (J. P.). — Voir Guyon (E.).	1553	cleavage plane on MBBA liquid crystal	1569
Cohen de Lara (E.) and Kahn (R.). — Neutron and infrared		Goldfarb (D.), Luz (Z.) and Zimmermann (H.). — A deute-	
study of the dynamical behaviour of methane in NaA	1000	rium NMR study of the discotic mesophase of hexa-	
zeolite	1029	hexyloxy-triphenylene	1303
Comes (R.). — Voir Yamaji (K.)		Grant (P.). — Voir Joanny (J. F.)	1045
Coret (A.). — Voir Haydar (A.)	893	Gröger (B.). — Voir Kronmüller (H.)	
Coste (J.). — Voir Tresser (C.).	427	Gros (Y.). — Voir Meyer (C.)	
Cotillard (J. C.). — Voir Porteseil (J. L.)	1253	Guillot (G.) et Nouailhat (A.). — Piégeage des interstitiels	
Coulomb (J. P.), Bienfait (M.) and Thorel (P.). — Mobility		par des impuretés divalentes dans KBr : Sr	1577
measurements of two kinds of two-dimensional fluids.		Guittard (M.). — Voir Vautier (R.)	885
Methane adsorbed on graphite	293	Guyon (E.), Clerc (J. P.), Giraud (G.) and Roussenq (J.).	
Coutine (L.). — Voir Le Paillier-Malecot (A.).			1553
Curie (D.). — Voir Hitter (G.)	479	Guyon (E.). — Voir Dreyfus (JM.)	283
Daniel (E.). — Voir Koenig (C.)	1015	Haldane (F. D. M.) and Villain (J.). — Commensurate-	
Danot (M.). — Voir Vautier (R.). Decruppe (J. P.). — Voir Dujardin (F.).		incommensurate transitions of physisorbed films on	1.000
Defresne (A.). — Voir Machizaud (F.).		anisotropic substrates.	1673
De Gennes (P. G.). — Motions of one stiff molecule in an		Hardouin (F.), Levelut (A. M.) and Sigaud (G.). — Anomalies	
entangled polymer melt		of periodicity in some liquid crystalline cyano deri-	71
De Gennes (P. G.). — Coherent scattering by one reptating	4/3	vatives Hardouin (F.). — Voir Sigaud (G.).	107
chain	735	Hardouin (F.). — Voir Sigaud (G.). Hardouin (F.). — Voir Levelut (A. M.).	
Demangeat (C.). — Voir Khan (M. A.).	573	Hartmann-Boutron (F.). — Voir Meyer (C.).	
		(-), (-), (-), (-), (-), (-), (-), (-),	200

Haydar (A.), Coret (A.), Draman (C.) et Oudjehane (N.).		Le Paillier-Malecot (A.) and Couture (L.) Properties of	
Photoconductivité excitonique de Cu ₂ O déformé plas-		the single and double D4d groups and their isomor-	
tiquement	893	phisms with D ₈ and C _{8v} groups	
Hendrikx (Y.) and Charvolin (J.). — Structural relations between lyotropic phases in the vicinity of the nematic		Le Toullec (R.). — Voir Piccioli (N.).	1129
phases	1427	Levelut (A. M.), Moussa (F.), Doucet (J.), Benattar (J. J.),	
Henry (L.). — Voir Duval (Ph.).	1427	Lambert (M.) and Dorner (B.). — Local order and lattice dynamics in the ordered phases of T.B.B.A.	
Hillner (B.). — Voir Limmer (St.)	751	(Terephthal-Bis-Butyl-Aniline)	1651
Hinov (H. P.). — Formation of a smectic A phase from a	-1003	Levelut (A. M.). — Voir Hardouin (F.)	71
strongly deformed planar nematic liquid crystal CBOOA	307	Levelut (A. M.). — Voir Sigaud (G.)	107
Hirlimann (C.). — Voir Oueslatti (M.)	1151	Levelut (A. M.), Hardouin (F.), Gasparoux (H.), Destrade (C.)	
Hitier (G.), Curie (D.) et Visocekas (R.). — Emission de	1000	and Nguyen Huu Tinh X-ray investigations and	
bandes larges dans ZnSe : cathodoluminescence, ther-		magnetic field effect on a nematic phase of disc-like	1 47
moluminescence, transitions dues aux centres profonds.	479	molecules Leyris (J. P.). — Voir Aguero (G.)	317
Horn (R. G.), Israelachvili (J. N.) and Perez (E.). — Forces		Limmer (St.), Findeisen (M.), Schmiedel (H.) and Hillner (B.).	317
due to structure in a thin liquid crystal film	39	- Investigations on the ordering of five p, p'-di-n-	
Israelachvili (J. N.). — Voir Horn (R. G.)	39	alkyloxy-azoxybenzenes in their nematic phases by	
viscosity measurements in the binary mixture butyl		proton NMR	1665
cellosolve-water near its upper and lower critical		Limmer (St.), Schmiedel (H.), Hillner (B.) and Lösche (A.). —	
consolute points	353	Transfer of the macroscopic alignment of meso-	
Jacoboni (C.). — Voir Dupas (C.)	1345	phases into the solid state	11111
Jamet (J. P.). — Voir Deville (A.)	1641	Lösche (A.). — Voir Limmer (St.). Louis (E.) and Flores (F.). — Schottky-barriers for clean,	1111
Jannin (M.). — Corrélation entre le bruit anormal et l'évo-		etched and reactive metal-semiconductor junctions	1313
lution de la structure en domaine des ferroélectriques	1269	Luz(Z.). — Voir Goldfarb (D.)	1303
Jeandey-Veyret (Cl.). — Exchange couplings in the free		Machizaud (F.), Kuhnast (FA.), Flechon (J.), Auguin (B.)	
radical nitroxide tanol suberate determined by static		and Defresne (A.). — Electron transport and kine-	
magnetic methods and room temperature proton relaxa-	875	tics of transformations in the amorphous alloy Ni ₆₆ B ₃₄	97
		Madej (C.). — Voir Pedrini (C.)	323
Jérôme (D.). — Voir Schulz (H. J.)	991	Mai (C.) Voir Tatibouet (J.)	1473
properties of the ordered VPt ₃ alloy. I. — Ferromagnetic		Malletta (H.). — Voir Deville (A.). Malhotra (V. M.) and Buckmaster (H. A.). — Estimation of	1641
behaviour	1157	τ ₁ for the host Kramers' Ln ³⁺ ions (Ln ≡ Nd, Sm and	
Joanny (J. F.), Grant (P.), Turkevich (L. A.) and Pincus (P.).		Yb) from the EPR linewidths of Gd ³⁺ impurity ions in	
Conformations of polydisperse polymer solutions:		Ln ₂ (SO ₄) ₃ .8 H ₂ O	1053
bimodal distribution	1045	Marais (A.). — Voir Vautier (R.).	885
Joanny (J. F.) and Brochard (F.). — Collapse of a polymer	1	Marcus (M.). — Quasicrystalline behaviour and phase tran-	
chain in a melt of incompatible polymer chains	1145	sition in cholesteric « blue » phase	61
Kahn (R.). — Voir Cohen de Lara (E.). Kalisky (Y.). — Voir Moine (B.).	1029	Marion (G.). — Etude aux rayons X de la phase incommen-	460
Khan (M. A.), Khwaja (Y.) and Demangeat (C.). — Origin	433	surable dans $(N(CH_3)_4)_2MCl_4$, $M = Co$ et Zn Martinot-Lagarde (Ph.) and Durand (G.). — Dielectric	469
of giant moments in non-stoichiometric FeRh alloys	573	relaxation in a ferroelectric liquid crystal	269
Khwaja (Y.). — Voir Khan (M. A.)	573	Martinoty (P.) and Bader (M.). — Measurement of the bire-	207
Kim (Y. H.). — Helix-coil transitions of dilute polymers		fringence induced in liquids by ultrasonic waves; appli-	
under a velocity gradient	247	cation to the study of the isotropic phase of PAA near	
Kléman (M.). — Voir Bourdon (L.)	261	the transition point	1097
Kléman (M.). — Barkhausen jumps in soft magnetic materials	1262	Mazaud (A.). — Voir Schulz (H. J.)	991
at low fields	1263 1461	McClure (D. S.). — Voir Pedrini (C.)	323 1327
Kléman (M.). — Voir Oswald (P.)	1401	Meister (H.). — Voir Yamaji (K.) Meister (H.). — Voir Bastie (P.)	445
ASA de la structure électronique d'impuretés « s-p »		Metzger (J.). — Voir Blaive (B.)	1533
dans le fer ferromagnétique	1015	Meyer (C.), Hartmann-Boutron (F.), Gros (Y.), Berthier (Y.)	KERT
Kostopoulos (D.), Mourikis (S.) and Varotsos (P.) Ionic		and Buevoz (J. L.). — Detailed study of NdFe2 and	
thermocurrent in KCl doped with SO ₄ ⁻²	1481	additional results relative to PrFe2 and YbFe2. Compa-	a policy
Kronmüller (H.) and Gröger (B.). — Domains, domain walls		rison with other R.E. Fe ₂ compounds	605
and the coercive field of amorphous ferromagnets	1285	Miranday (J. P.). — Voir Dupas (C.)	1345
Kuentzler (R.). — Voir Jesser (R.)	1157	Moine (B.), Bourcet (J. C.), Boulon (G.), Reisfeld (R.) and Kalisky (Y.). — Interaction mechanisms in the Bi ³⁺ -	
Kuhnast (FA.). — Voir Machizaud (F.)	97	Eu ³⁺ energy transfer in germanate glass at low tempe-	
Laheurte (J. P.), Noiray (J. C.), Obadia (M.) and Romagnan (J. P.). — Dielectric measurements of adsorbed kryp-		rature	499
ton on lamellar halides	559	Monceau (P.). — Voir Briggs (A.)	1453
Lambert (M.). — Voir Levelut (A. M.)	1651	Mott (N. F.). — Metal-insulator transition in Ti ₂ O ₃	277
Laridjani (M.) and Sadoc (J. F.) Structure study of		Mourikis (S.). — Voir Kostopoulos (D.)	1481
amorphous Gd-Y alloys	1293	Moussa (F.). — Voir Levelut (A. M.)	1651
Lascombe (J.). — Voir Besnard (M.)	593	Müller (K. A.). — Paramagnetic point and pair defects in oxide perovskites	551
Le Dang (K.). — Voir Dupas (C.)	1345	Nguyen Huu Tinh. — Voir Levelut (A. M.)	147
Leenhouts (F.). — Voir Flapper (S. D. P.)	1647	Noiray (J. C.). — Voir Laheurte (J. P.)	559
Lejcek (L.). — Voir Bourdon (L.) Leleyter (M.). — Emission ionique secondaire et structure	261	Nouailhat (A.). — Voir Guillot (G.)	1577
électronique des petits agrégats VC _n		Nunez-Regueiro (M.). — Voir Briggs (A.)	1453
Léonard (P) Voir Koenig (C)	1015	Obadia (M.). — Voir Laheurte (J. P.)	559

Oswald (P.) et Kléman (M.). — Défauts dans une mésophase	Schlapbach (L.) and Brundle (C. R.). — XPS study of the
hexagonale discotique: disinclinaisons et parois 1461	chemisorption induced surface segregation in LaNi ₅
Oudjehane (N.). — Voir Haydar (A.)	and ThNi ₅ 1025
Oueslatti (M.), Hirlimann (C.) et Balkanski (M.). — Diffu-	Schlenker (M.). — Voir Baruchel (J.) 1279
sion inélastique de la lumière à la résonance dans	Schmiedel (H.). — Voir Limmer (St.)
GaAs _{1-x} P _x 1151	Schulz (H. J.), Jérome (D.), Mazaud (A.), Ribault (M.) and
Pagost (P. O.). — Voir Pedrini (C.)	Bechgaard (K.). — Possibility of superconducting pre-
Palmer (S. B.). — Voir Baruchel (J.)	cursor effects in quasi-one-dimensional organic con-
Panyutin (V. L.). — Voir Chizhikov (V. I.)	ductors: theory and experiments
Parisi (G.). — Voir Vannimenus (J.)	Servoin (J. L.). — Voir Quittet (A. M.)
Parodi (O.). — Voir Perez (A.)	Sherrell (P. L.). — Voir Bunning (J. D.).
Pedrini (C.), Pagost (P. O.), Madej (C.) et McClure (D. S.).	Sigaud (G.). — Voir Hardouin (F.)
Photoconductivité due à l'autoionisation de Ce ²⁺ dans	Sigaud (G.), Hardouin (F.), Achard (M. F.) and Levelut
CaF_2 , SrF_2 et BaF_2	(A. M.). — A new type of smectic A phase with long
Perez (A.), Brunet (M.) and Parodi (O.). — Défauts dans	range modulation in the layers
les smectiques C chiraux : III. — Défauts toriques de	Spiesser (M.). — Voir Vautier (R.)
surface 1559	Strzelecki (L.). — Voir Pieranski (P.)
Perez (E.). — Voir Horn (R. G.)	Tatibouet (J.), Mai (C.), Perez (J.) et Vassoille (R.). — Etude
Perez (J.). — Voir Tatibouet (J.)	du vieillissement de la glace I _h par mesures de frotte-
Peyre (H.). — Voir Duval (Ph.)	ment interne et par observation en topographie aux
Piccioli (N.), Le Toullec (R.), Bertrand (F.) et Chervin (J. C.).	rayons X 1473
Constantes optiques de InSe entre 10 500 cm ⁻¹	Taupin (D.). — Voir Bourdon (L.)
(1,30 eV) et 22 500 cm ⁻¹ (2,78 eV) 1129	Thorel (P.). — Voir Coulomb (J. P.)
Picoli (G.), Deveaud (B.) and Galland (D.). — Interpretation	Thurston (R. N.) and Almgren (F. J.) Liquid crystals
of luminescence in GaAs : Cr : 0.839 eV and 0.574 eV	and geodesics
lines	
	Thurston (R. N.). — Stability of nematic liquid crystal
Picot (C.). — Voir Izumi (Y.)	configurations
Pieranski (P.). — Voir Dreyfus (JM.)	Toulouse (G.). — Voir Vannimenus (J.) 56.
Pieranski (P.), Dubois-Violette (E.), Rothen (F.) and Strze-	Travers (J. P.). — Mouvement aléatoire unidimensionnel
lecki (L.). — Geometry of Kossel lines in colloidal	de sphères dures : influence d'une interaction
crystals 53	d'échange 110.
Pincus (P.). — Voir Joanny (J. F.)	Trebin (HR.). — Elastic energies of a directional medium 1573
Pokrovsky (V.). — Voir Feigelman (M.)	Tresser (C.), Brot (C.) and Coste (J.). — Random walks
Ponedelnikov (B. E.). — Voir Chizhikov (V. I.) 1003-1039	with sign reversals and the intrinsic angular momentum
Porteseil (J. L.), Cotillard (J. C.), Vergne (R.) et Ferrari (G.).	behaviour on lattices
— Etude expérimentale des mécanismes d'activation	
	Troadec (J. P.) et Bideau (D.).—Propriétés de transport
thermique dans le déplacement d'une paroi de Bloch	d'un mélange de sphères conductrices et isolantes : étude
à 180°	des effets de percolation
Prost (J.). — Voir Ricard (L.)	Turkevich (L. A.). — Voir Joanny (J. F.)
Ptak (M.). — Voir Sanson (A.)	Vallade (M.). — Voir Bastie (P.)
Quittet (A. M.), Servoin (J. L.) and Gervais (F.). — Correla-	Vannimenus (J.), Toulouse (G.) and Parisi (G.). — Study of
tion of the soft modes in the orthorhombic and cubic	a simple hypothesis for the mean-field theory of spin-
phases of KNbO ₃	glasses
Reisfeld (R.). — Voir Moine (B.)	Varotsos (P.). — Voir Kostopoulos (D.)
Renard (J. P.). — Voir Dupas (C.)	Vassoille (R.). — Voir Tatibouet (J.)
Ribault (M.). — Voir Schulz (H. J.)	
Ribault (M.). — Voir Briggs (A.)	Vautier (R.), Marais (A.), Villers (G.), Guittard (M.),
Ricard (L.) and Prost (J.). — Critical behaviour of second	Danot (M.) et Spiesser (M.). — Propriétés magnétiques
sound near the smetic A nematic phase transition 861	de FeTi ₂ S ₄ et des composés Fe _{1+ε} Ti _{2+2ε} S ₄ ($\varepsilon > 0$) 885
TANAMAY NOV - 4 President	Veillet (P.). — Voir Dupas (C.)
Richard (J.). — Voir Briggs (A.)	Vergne (R.). — Voir Porteseil (J. L.)
Riedinger (R.). — Voir Dreysse (H.)	Vertogen (G.). — Voir Flapper (S. D. P.)
Romagnan (J. P.). — Voir Laheurte (J. P.) 559	Vettier (C.). — Voir Bastie (P.)
Rothen (F.). — Voir Pieranski (P.)	Veyret-Jeandey (Cl.). — Voir Chouteau (G.)
Roulet (B.). — Voir Abraham-Ibrahim (S.)	Villa (M.). — Voir Bjorkstam (J. L.)
Roussenq (J.). — Voir Guyon (E.)	Villain (J.). — Voir Haldane (F. D. M.)
Rozenson (A. E.). — Voir Chizhikov (V. I.) 1003-1039	Villers (G.). — Voir Vautier (R.)
Sadoc (J. F.). — Voir Laridjani (M.)	
Sanson (A.) et Ptak (M.). — Etude par RPE de l'organisation	
de la phase smectique C du 4-4' di-heptyloxyazoxyben-	Webb (W. W.). — Voir Chan (W. K.)
zène	Yamaji (K.), Megtert (S.) and Comes (R.). — 2D displace-
	ment pattern in TSeF-TCNQ model analysis of the
Saubade (Ch.). — Indice de réfraction de l'eau pure aux	2 k _F diffuse lines
basses températures, pour la longueur d'onde de	Zeyen (C. M. E.). — Voir Bastie (P.)
5 893 Å	Zimmermann (H.). — Voir Goldfarb (D.)
	CET ADDENDA
COMMENTAIRES	ET ADDENDA
Notice of Co. Co. Control of the Control of the Co.	
Barreto (E.) and Jurenka (H.). — Comments on «the	Callimberti (I) Discussion of the assessment on (the
mechanism of the long spark formation »	Gallimberti (I.). — Discussion of the comments on « the
133	mechanism of the long spark formation »
	Jurenka (H.). — Voir Barreto (E.)

JOURNAL DE PHYSIQUE — LETTRES

Tome 42 — 1981

MECANIQUE, HYDRODYNAMIQUE		Lubensky (T. C.) and McKane (A. J.). — Anderson localization, branched polymers and the Yang-Lee edge sin-	
Bachelart (S.). — Voir Pomeau (Y.)	L-271	gularity	L-331
Bailleux (R.). — Voir Pantaloni (J.)	L-147	Luck (J. M.). — Finite-size lattice method and the rou-	12-331
Barquins (M.). — Voir Maugis (D.)	L-95	ghening transition	L-275
Brand (H.) and Pleiner (H.) Hydrodynamics of choles-	told.	McKane (A. J.). — Voir Lubensky (T. C.)	L-331
teric liquid crystals with impurities	L-327	Martinez-Mekler (G. C.) and Moore (M. A.). — Diffusion	L-331
Cerisier (P.). — Voir Pantaloni (J.)	L-147	of a polymer in a random medium	L-413
Chauveteau (G.) and Moan (M.). — The onset of dilatant	brodo/	Milczarek (J.) and Wentowska (K.). — The nematic critical	L-413
behaviour in non-inertial flow of dilute polymer solu-		point in the molecular field approximation	L-335
tions through channels with varying cross-sections	L-201		L-333
Clavin (P.). — Voir Stauffer (D.)	L-353	Moore (M. A.). — Voir Martinez-Mekler (G. C.) Orland (H.), De Dominicis (C.) and Garel (T.). — Non-	L-413
Couder (Y.). — The observation of a shear flow instability	L-333		L-73
in a rotating system with a soap membrane	1-420	homogeneous mean field picture for spin-glasses	
Fauve (S.), Laroche (C.) and Libchaber (A.). — Effect of a	L-42)	Roussenq (J.). — Voir Clerc (J. P.)	L-249
horizontal magnetic field on convective instabilities in		Sourlas (N.). — Critical dimension and 1/D expansion for	1 222
mercury	I 455	the random anisotropy axis model	L-233
Gerbaud (C.). — Voir Pantaloni (J.)		Stauffer (D.). — Monte-Carlo simulation of Ising droplets	Y 00
		in correlated site-bond percolation	L-99
Laroche (C.). — Voir Fauve (S.)		Stauffer (D.). — Voir Clerc (J. P.)	L-249
Libchaber (A.). — Voir Fauve (S.)	L-455	Toulouse (G.) and Gabay (M.). — Mean field theory for	Y 102
Maugis (D.) and Barquins (M.). — Adhesive contact of a	1.05	Heisenberg spin glasses	L-103
conical punch on an elastic half-space	L-95	Tsallis (C.) and De Magalhães (A. C. N.). — Conjecture on	
Moan (M.). — Voir Chauveteau (G.)	L-201	the critical frontier of the Potts ferromagnet on planar	T 227
Pantaloni (J.), Cerisier (P.), Bailleux (R.) et Gerbaud (C.).		lattices	L-227
Convection de Bénard-Marangoni; un pendule de	Y 147	Wentowska (K.). — Voir Milczarek (J.)	L-335
Foucault ?			
Pleiner (H.). — Voir Brand (H.)	L-321	PHYSIQUE NUCLÉAIRE	
Pomeau (Y.). — Non linear pattern selection in a problem	Y 1	THISTQUE THE CERTIFICE	
of elasticity	L-1	Danier (T.E.) Chad (M) and Come (D.) Missosomic	
Pomeau (Y.), Roux (J. C.), Rossi (A.), Bachelart (S.) and		Berger (JF.), Girod (M.) and Gogny (D.). — Microscopic	1 500
Vidal (C.). — Intermittent behaviour in the Belousov-		description of nuclear fission	L-509
Zhabotinsky reaction		Bocquet (J. P.). — Voir Nifenecker (H.)	L-527
Rossi (A.). — Voir Pomeau (Y.)		Girod (M.). — Voir Berger (JF.).	L-509
Roux (J. C.). — Voir Pomeau (Y.)	L-271	Gogny (D.). — Voir Berger (JF.)	L-509
Strauffer (D.) and Clavin (P.). — Simple calculation of sedi-		Guet (C.). — Voir Signarbieux (C.)	L-437
mentation velocity in concentrated dispersions		Mariolopoulos (G.). — Voir Nifenecker (H.)	L-527
Vidal (C.). — Voir Pomeau (Y.)	L-271	Marquez (L.). — The nuclear radius from alpha decay	L-181
		Maurel (M.). — Voir Signarbieux (C.)	L-437
MÉCANIQUE STATISTIQUE		Mazur (C.). — Voir Signarbieux (C.)	L-437
		Montoya (M.). — Voir Signarbieux (C.)	L-437
Caillol (J. M.). — Exact results for a two-dimensional one-		Nifenecker (H.), Mariolopoulos (G.) and Bocquet (J. P.).	1 627
component plasma on a sphere	L-245	Pair breaking mechanism in cold fission	L-527
Carton (J. P.). — Voir Clerc (J. P.)	L-249	Perrin (P.). — Voir Signarbieux (C.)	L-437
Clerc (J. P.), Carton (J. P.), Roussenq (J.) and Stauffer (D.).	2217	Ribrag (M.). — Voir Signarbieux (C.)	L-437
— Surface effects for percolation	L-249	Signarbieux (C.), Montoya (M.), Ribrag (M.), Mazur (C.),	
De Dominicis (C.). — Voir Orland (H.)	L-73	Guet (C.), Perrin (P.) and Maurel (M.). — Evidence for	
De Gennes (P. G.). — Some effects of long range forces on	275	nucleon pair breaking even in the coldest scission con-	7 427
interfacial phenomena	L-377	figurations of ²³⁴ U and ²³⁶ U	L-437
De Magalhães (A. C. N.). — Voir Tsallis (C.)	L-227		
Des Cloizeaux (J.). — Ring polymers in solution: topolo-	1,-661	PHYSIQUE ATOMIQUE ET MOLÉCULA	IRE
	L-433	THISTQUE MICHIQUE ET MICHECULE	
gical effects	L-103	Aboumajd (A.). — Voir Berger (H.)	L-55
	L-73	Aspect (A.). — Voir Vigué (J.)	L-531
Garel (T.). — Voir Orland (H.) Jancovici (R.) — Charge distribution and kinetic pressure	13	Berger (H.) et Aboumajd (A.). — Expression des constantes	201
Jancovici (B.). — Charge distribution and kinetic pressure	L-223	de distorsion centrifuge des hexafluorures en fonction	
in a plasma: a soluble model	2000	des fréquences harmoniques. Détermination de quel-	
trolyte chain below six dimensions	L-409	ques constantes cubiques du potentiel	L-55
Vertical (I) Presentation of holes between everlapping	L-403	Bloch (D.), Raj (R. K.), Snyder (J. J.) and Ducloy (M.).	1-33
Kertész (J.). — Percolation of holes between overlapping		Heterodyne detection of phase-conjugate emission in	
spheres: Monte Carlo calculation of the critical volume	L-393	an Ar discharge with a low-power c.w. laser	L-31
fraction	1-373	an Ai discharge with a low-power c.w. laser	2001


Boudouris (G.). — Voir Kouroupetroglou (G. T.)	L-441	Gorse (C.). — Voir Capitelli (M.) L-185	5-L-469
Brault (J. W.). — Voir Camy-Peyret (C.)	L-279	Hansen (J. P.). — Plasmon dispersion of the strongly	
Camy-Peyret (C.), Flaud (J. M.), Delbouille (L.), Roland		coupled one component plasma in two and three	1 207
(G.), Brault (J. W.) and Testerman (L.). — Quadrupole		dimensions	L-397
transitions of the $1 \leftarrow 0$ band of N_2 observed in a high	Y 000	Hansen (J. P.). — Voir Leger (D.).	L-459
resolution atmospheric spectrum	L-279	Leger (D.), Deutsch (C.) and Hansen (J. P.). — Electronic	
Davy (P.). — Voir Pignolet (P.)	L-205	transport in strongly coupled and fully ionized H-He	L-459
Delbouille (L.). — Voir Camy-Peyret (C.)	L-279	mixtures	L-435
Denis (A.) and Desesquelles (J.). — Observation of optical	L-59	Oda (Y.). — Voir Furutani (Y.)	L-285
emission from beam-foil excited Li	L-59	Ricard (A.). — Voir Capitelli (M.) L-185	
Desesquelles (J.). — Voir Denis (A.)	L-205	means (As). For Capiton (I.E.).	2 103
Ducloy (M.). — Voir Bloch (D.)	L-31		
Flaud (J. M.). — Voir Camy-Peyret (C.)	L-279		
Fontaine (B.). — Voir Forestier (B.)	L-211	PHYSIQUE DE LA MATIÈRE CONDEN	SÉE
Forestier (B.), Fontaine (B.) and Solenne (T.). — XeCl laser		The second secon	
pumped by an electron beam and X-ray assisted dis-		Ablart (G.), Bourdel (D.), Pescia (J.), Clément (S.) and	
charge	L-211	Renard (J. P.). — Temperature and orientation depen-	
Grangier (P.). — Voir Vigué (J.)	L-531	dences of the E.S.R. linewidth in T.M.M.C.	L-219
Kobeissi (H.). — Sur une équation aux valeurs propres de		Achard (M. F.). — Voir Hardouin (F.)	
vibration-rotation pour tout potentiel électronique	00	Aldissi (M.). — Voir Bernier (P.)	
d'une molécule diatomique	L-151	Alizon (J.). — Voir Granier (W.)	L-301
Kobeissi (H.). — Sur une expression explicite de l'effet de		Allain (C.). — Voir Vidakovic (P.)	L-323
rotation dans la fonction d'onde de vibration-rotation		Almairac (R.), Regis (M.), Nouet (J.) and Filippini (C.).	
d'une molécule diatomique	L-215	Observation of forbidden Bragg reflections in BaMnF ₄	L-385
Kouroupetroglou (G. T.) and Boudouris (G.). — Collision	T 441	Andrieux (A.), Jérome (D.) and Bechgaard (K.). — Spin-	
induced absorption in CO ₂ at 0.091 cm ⁻¹	L-441	density wave ground state in the one-dimensional	
Minard (J.). — Voir Pignolet (P.)	L-25 L-205	conductor (TMTSF) ₂ PF ₆ : microscopic evidence from	111
Peresse (J.). — Voir Tuffin (F.)	L-25	⁷⁷ Se and ¹ H NMR experiments	L-87
Pignolet (P.), Dubreuil (B.), Davy (P.) and Minard (J.).	2-25	Arnold (W.), Doussineau (P.), Frénois (Ch.) and Levelut (A.).	
Observation of splitting and optical satellites on the		- On the ultrasonic absorption and dispersion in	7 200
H _a line of hydrogen due to non-resonant interaction		amorphous superconductors	L-289
with a CO ₂ laser radiation	L-205	Barois (P.), Coulon (C.) and Prost (J.). — Escape from	
Raj (R. K.). — Voir Bloch (D.)	L-31	incommensurability: the two-dimensionally ordered $S_{\tilde{A}}$ antiphase	I -107
Roger (G.). — Voir Vigué (J.)	L-531	Bechgaard (K.). — Voir Andrieux (A.).	
Roland (G.). — Voir Camy-Peyret (C.)	L-279	Bechgaard (K.). — Voir Fournel (A.)	
Snyder (J. J.). — Voir Bloch (D.)	L-31	Bechgaard (K.). — Voir More (C.)	
Solenne (T.). — Voir Forestier (B.)	L-211	Bechgaard (K.). — Voir Pouget (J. P.)	
Testerman (L.). — Voir Camy-Peyret (C.)	L-279	Bechgaard (K.). — Voir Schulz (H. J.)	L-51
Torchin (L). — Energy transfer in near-resonant molecular collisions from diatomic to symmetric top molecules	L-357	Bechler-Ferry (E.). — Voir Esling (C.)	L-141
Tuffin (F.), Le Coz (G.) et Peresse (J.). — Etude du spectre	L-331	Bellessa (G.). — Voir Duquesne (J. Y.)	L-491
en énergie des électrons éjectés de la molécule de gaz		Benattar (J. J.), Moussa (F.), Lambert (M.) and Ger-	
carbonique par impact d'atomes d'hélium dans les		main (C.). — Two kinds of two-dimensional order:	
états 2 ¹ S et 2 ³ S et de néon dans les états ³ P ₂ et ³ P ₀	L-25	the SmF and SmI phases	L-67
Vigué (J.), Grangier (P.), Roger (G.) and Aspect (A.).		Benguigui (L.) and Hardouin (F.). — Reentrant nematic and smectic A phases: the dielectric approach	T 111
Polarization of calcium atomic fluorescence due to a		Benguigui (L.) and Hardouin (F.). — Antiferroelectric liquid	L-111
coherence effect in the photodissociation of Ca ₂ mole-		crystals	L-381
cules	L-531	Bernier (P.), Linaya (C.), Rolland (M.) and Aldissi (M.).	2 301
		EPR study of the cis-trans isomerization process in	
		polyacetylene films	L-295
PHYSIQUE DES PLASMAS		Bernier (P.). — Voir Granier (W.)	L-301
TITISIQUE DES TEASIVIAS		Bertault (M.), Collet (A.) and Schott (M.). — A calori-	
		metric study of pTS (bis-p-toluenesulphonate of 2,4-	
Bacal (M.), Bruneteau (A. M.) and Nachman (M.). — Elec-		hexadiyne 1,6-diol) and its phase transitions as a func-	
tron collection radius of an electrostatic probe immers-		tion of polymerization	L-131
ed in a low-pressure plasma	L-5	Beuneu (B.) and Quéré (Y.). — Paraelastic interstitials in	W 10.
Bernu (B.). — One-component plasma in a strong uniform	1 262	molybdenum	L-465
magnetic field Bruneteau (A. M.). — Voir Bacal (M.)	L-253 L-5	Blumstein (A.). — Voir Volino (F.)	L-305
Capitelli (M.), Gorse (C.) and Ricard (A.). — Relaxation	12-3	Blumstein (R. B.). — Voir Volino (F.). Boeuf (A.), Fournier (J. M.), Heger (G.), Lehner (N.),	1303
of the vibrational distribution function in N ₂ time		Manes (L.), Rebizant (J.), Rustichelli (F.) and Spir-	
varying discharges	L-185	let (J. C.). — Recent results on the NpO ₂ 25 K tran-	
Capitelli (M.), Gorse (C.) and Ricard (A.). — Influence of	1000000	sition by neutron diffraction	L-401
superelastic vibrational collisions on the relaxation	50	Bonjour (E.). — Voir Calemczuk (R.)	L-501
of the electron energy distribution function in N2		Borckmans (P.). — Voir Dewel (G.)	L-361
post discharge regimes	L-469	Bouchet (B.) and Glättli (H.). — Spin-lattice relaxation mea-	
Deutsch (C.). — Voir Furutani (Y.)	L-285	surements and spin conversion in methane below 1 K	L-159
Deutsch (C.). — Voir Leger (D.)	L-459	Bourdel (D.). — Voir Ablart (G.)	L-219
Furutani (Y.), Deutsch (C.) and Oda (Y.). — Evaluation of the first Coulomb Bridge in three dimensions	1 205	Brownsey (G. J.) and Leadbetter (A. J.). — Novel liquid	
of the first Coulomb Bridge in three dimensions	L-285	crystal structures in cyano bi-cyclohexanes	L-135

Bruyère (J. C.). — Voir Currie (J. F.)	L-373	Esling (C.), Bechler-Ferry (E.) and Bunge (HJ.). —	
Buisson (R.) and Vial (J. C.). — Transfer inside pairs of		Numerical calculation of the odd part of the texture	
Pr3+ in LaF3 studied by up-conversion fluorescence	I_115	function calculation of the odd part of the texture	
Bunge (HJ.). — Voir Esling (C.)	L-113	function	L-141
Burger (J. P.). — Voir Sniadower (L.)	L-141	Faetti (S.). — Voir Domenici (C.)	L-347
Parigor (0. 1.), Volt Dimetorici (12)	L-13	Filippini (C.). — Voir Almairac (R.)	L-385
Cagnon (M.) and Durand (G.). — Newtonian shear creep		Fleury (G.). — Voir Viger (C.)	L-495
in a smectic B plastic crystal	L-451	Fournel (A.), More (C.), Roger (G.), Sorbier (J. P.), Del-	
Calemczuk (R.) and Bonjour (E.) Structural relaxation		rieu (J. M.), Jérome (D.), Ribault (M.) and Bechgaard	
of glassy selenium under γ-irradiation	L-501	(K.). — The three dimensional stabilization of super-	
Campbell (L A.). — Voir Hamzić (A.) L-1	7-L-309	conductivity at 12 K in the one dimensional organic	
Catlow (C. R. A.). — Voir Corish (J.)	L-369	superconductor (TMTSF) ₂ PF ₆ under 11 kbar	L-445
Chaillou (D.), Chambaudet (A.) et Zidler (B.) Sur la	2 307	Fournier (J. M.). — Voir Boeuf (A.)	
rétention des traces de fission de l'uranium dans les			L-401
		Frénois (Ch.). — Voir Arnold (W.)	L-289
minéraux	L-343	Fretier (P.). — Voir Gourdon (J. C.)	L-21
Chambaudet (A.). — Voir Chaillou (D.)	L-343	Fries (P. H.). — Quelques résultats théoriques sur l'étude	
Chaves (C. M.). — Voir Lederer (P.)	L-127	par R.M.N. des fluides polyatomiques plans	L-257
Clément (S.). — Voir Ablart (G.)	L-219	Fries (P. H.). — Etude par R.M.N. d'un fluide polyatomique	
Collet (A.). — Voir Bertault (M.)	L-131	physisorbé	L-513
Çomès (R.). — Voir Pouget (J. P.)	L-543	Frigerio (J. M.). — Voir Rivory (J.)	L-481
Čopič (M.), Zgonik (M.) and Lavrenčič (B. B.). — Brillouin		Fronzoni (L.). — Voir Domenici (C.)	L-347
scattering in S _A and S _B phases of butyloxy-benzilidene		Froyer (G.). — Voir Maurice (F.)	L-425
octylaniline	L-405	Gabay (M.). — Voir De Dominicis (C.)	L-523
Corbel (C.) and Gupta (R. P.) Positron lifetime in		Galarneau (S.). — Voir Currie (J. F.)	L-373
vacancy-impurity complexes	L-547	Caracha (D) and Vannaia (V. I.) Surgeon dustinity in	L-3/3
Corish (J.), Catlow (C. R. A.) and Jacobs (P. W. M.). — The	2011	Garoche (P.) and Veyssie (J. J.). — Superconductivity in	* 265
relationship between experimental and calculated		amorphous versus crystalline Cu _{0.33} Zr _{0.66} alloys	L-365
	T 260	Gauneau (M.). — Voir Maurice (F.)	L-425
point defect energies.	L-369	Germain (C.). — Voir Benattar (J. J.)	L-67
Couget (A.) — Voir Mamy (R.)	L-79	Glättli (H.). — Voir Bouchet (B.)	L-159
Coulon (C.) and Prost (J.). — Simple example of fluctuation		Globus (A.). — Voir Pascard (H.)	L-319
induced reentrant behaviour	L-241	Gourdon (J. C.), Fretier (P.) and Pescia (J.). — Spin-lattice	
Coulon (C.). — Voir Barois (P.)	L-107	relaxation in amorphous silicon. Evidence of ano-	
Croquette (V.) and Poitou (C.). — Cascade of period		malous temperature dependences	L-21
doubling bifurcations and large stochasticity in the		Granier (W.), Bernier (P.), Dohri (M.), Alizon (J.) and	
motions of a compass	L-537	Robert (H). — 19F nuclear relaxation in an ionic	
Currie (J. F.), Depelsenaire (P.), Galarneau (S.), L'Ecuyer		conductor: TlSn ₂ F ₅	L-301
(J.), Groleau (R.), Bruyère (J. C.) and Deneuville (A.).		Gressier (P.). — Voir Mihaly (G.)	L-263
- Thickness dependence of hydrogen in a-Si: H films		Groleau (R.). — Voir Currie (J. F.)	L-373
deposited on c-Si	L-373	Gross (J. G.). — Voir Lewonczuk (S.)	L-91
Dallaporta (H.). — Voir Humbert (A.)	L-167	Gupta (R. P.). — Voir Corbel (C.)	L-547
Daniel (E.). — Voir Koenig (C.)	L-193	Hamzić (A.) and Campbell (L. A.). — The ferromagnetic to	Y 10
Debever (J. M.). — Voir Humbert (A.)	L-167	spin glass transition	L-17
De Dominicis (C.), Gabay (M.) and Orland (H.). —		Hamzić (A.) and Campbell (I. A.). — The spin glass to ferro-	
Replica derivation of Sompolinsky free energy func-		magnetic order transition in AuFe alloys	L-309
tional for mean field spin glasses	L-523	Hanus (J.). — Voir Humbert (A.)	L-167
Delrieu (J. M.). — Voir Fournel (A.)	L-445	Hardouin (F.), Sigaud (G.), Nguyen Huu Tinh and Achard	
Deneuville (A.). — Voir Currie (J. F.)	L-373	(M. F.). — A fluid smectic A antiphase in a pure nitro	
Depelsenaire (P.). — Voir Currie (J. F.)	L-373	rod-like compound	L-63
Derzhanski (A.), Petrov (A. G.) and Pavloff (Y. V.).	D-373	Hardouin (F.). — Voir Benguigui (L.) L-111	-L-381
Curvature induced conductive and displacement cur-		Heger (G.). — Voir Boeuf (A.)	L-401
•	T 110	Housseau (N.). — Voir Mihaly (G.)	L-263
rents through lipid bilayers	L-417	Hu Hua-Chen. — Explanation of the anomalous enhance-	
Destrade (C.). — Voir Nguyen Huu Tinh		ment effect observed in single crystals neutron dif-	
Deutscher (G.). — Voir Kapitulnik (A.)	L-341	fraction of α-LiIO ₃ : a new method to determine the	
Dewel (G.), Walgraef (D.) and Borckmans (P.). — Layered	1 2/1		L-189
structures in twodimensional nonequilibrium systems	L-361	activation energy of α-LilO ₃ by this effect	107
Djafari Rouhani (B.) et Joyes (P.). — Sur la relaxation		Humbert (A.), Debever (J. M.), Dallaporta (H.) and Hanus	
des distances interatomiques par effet de taille	L-389	(J.). — Observation of surface plasmons by transition	T 167
Dohri (M.). — Voir Granier (W.)	L-301	radiation from smooth aluminium films	L-167
Domenici (C.), Fletti (S.) and Fronzoni (L.). — Inversion		Jacobs (P. W. M.). — Voir Corish (J.)	L-369
effect of electrohydrodynamic vortices in the isotropic		Jamet (J. P.). — Electric field phase diagram of thiourea	
and anisotropic phase of thin nematic layers with two		determined by optical birefringence	L-123
free surfaces	L-347	Jánossy (L). — Diffuso-mechanical coupling in cholesteric	
Doussineau (P.). — Ultrasonic propagation in amorphous		liquid crystals	L-41
PdSiCu at low temperatures	L-83	Jérome (D.). — Voir Andrieux (A.)	L-87
Doussineau (P.) and Matecki (M.). — Ultrasonic propa-		Jérome (D.). — Voir Fournel (A.)	L-445
notion in a fluoresimental class from 0.08 K to 250 K	L-267	Jérome (D.). — Voir More (C.)	L-313
gation in a fluorozirconate glass from 0.08 K to 250 K	L-289	Jérome (D.). — Voir Schulz (H. J.)	L-51
Doussineau (P.). — Voir Arnold (W.)		Jezequel (G.). — Voir Mamy (R.)	L-473
Dumoulin (L.). — Voir Sniadower (L.)	L-13		L-45
Duquesne (J. Y.) and Bellessa (G.). — Internal friction		Joffrin (J.). — Voir Lynn (J. W.)	
in amorphous polystyrene at low temperature and	Y 404	Joyes (P.). — Voir Djafari Rouhani (B.)	L-389
high frequency	L-491	Kapitulnik (A.), Rappaport (M. L.) and Deutscher (G.).	
Durand (G.). — Voir Cagnon (M.)	L-451	The effect of crystallinity on the morphology of eva-	
Durand (C) Vois Potroy (M)	L-519	porated Al-Ge thin films	L-541

Koenig (C.) and Daniel (E.). — LMTO-ASA method for the		Nouet (J.). — Voir Almairac (R.)	L-385
electronic structure of dilute alloys	L-193	Orland (H.). — Voir De Dominicis (C.)	L-523
Kutka (R.). — Voir Trebin (HR.)	L-421	Oswald (P.). — Observation in homeotropic anchoring of	
Laheurte (J. P.), Noiray (J. C.) and Romagnan (J. P.).		the $+\pi$ disclinations in a hexagonal discotic liquid	
Third sound velocity in thin films of ³ He- ⁴ He mixtures		crystal	L-171
at low temperature	L-197	Papanicolaou (N.). — Voir Lapujoulade (J.)	L-463
Lambert (M.). — Voir Benattar (J. J.)	L-67	Pascard (H.) and Globus (A.). — Volume anomaly in ferri-	
Laporte (P.) and Subtil (J. L.). — Reflectance measurement		magnetism	L-319
of the VUV spectrum of solid xenon and its tempe-		Pavloff (Y. V.). — Voir Derzhanski (A.)	L-119
rature dependence up to the triple point	L-339	Pelissier (J.). — Voir Mihaly (G.)	L-263
Lapujoulade (J.), Le Cruër (Y.), Lefort (M.), Lejay (Y.),			L-219
Maurel (E.) and Papanicolaou (N.). — Diffraction		Pescia (J.). — Voir Ablart (G.)	
of neon by a (117) copper surface	L-463	Pescia (J.). — Voir Gourdon (J. C.)	L-21
Laridjani (M.). — Voir Sadoc (J. F.)	L-485	Petrov (M.) and Durand (G.). — On the origin of elec-	
• • •	L-465	trohydrodynamic instabilities in smectic C liquid crys-	
Lauter (H. J.). — Voir Tiby (C.)	L-105	tals	
Lavrenčič (B. B.). — Voir Čopič (M.)		Petrov (A. G.). — Voir Derzhanski (A.)	L-119
Leadbetter (A. J.). — Voir Brownsey (G. J.)	L-135	Poitou (C.). — Voir Croquette (V.)	L-537
Le Cruër (Y.). — Voir Lapujoulade (J.)	L-463	Pouget (J. P.), Moret (R.), Comès (R.) and Bechgaard (K.).	
L'Ecuyer (J.). — Voir Currie (J. F.)	L-373	— X-ray diffuse scattering study of superstructure	
Lederer (P.) and Chaves (C. M.). — Phases diagram of		formation in tetramethyltetraselenafulvalenium per-	
thiourea at atmospheric pressure under electric field:	T 100	rhenate (TMTSF) ₂ ReO ₄ and nitrate (TMTSF) ₂ NO ₃	L-543
a theoretical analysis	L-127	Prost (J.). — Voir Barois (P.)	L-107
Lefort (M.). — Voir Lapujoulade (J.)	L-463	Prost (J.). — Voir Coulon (C.)	L-241
Lehner (N.). — Voir Boeuf (A.)	L-401	Pynn (R.). — Voir Lynn (J. W.)	L-45
Lejay (Y.). — Voir Lapujoulade (J.)	L-463	Quéré (Y.). — Voir Beuneu (B.)	L-465
Lemonnier (J. C.). — Voir Mamy (R.)	L-473	Raggazoni (A.) Voir Lynn (J. W.)	L-45
Levelut (A.). — Voir Arnold (W.)	L-289	Rappaport (M. L.). — Voir Kapitulnik (A.)	L-541
Lewonczuk (S.), Gross (J. G.) and Ringeissen (J.). — Tem-		Rebizant (J.). — Voir Boeuf (A.)	L-401
perature dependence of the spin-orbit coupling and		Rebouillat (J. P.) Voir Rivory (J.)	L-481
energy gap in cuprous bromide	L-91	Regis (M.). — Voir Almairac (R.)	L-385
Linaya (C.). — Voir Bernier (P.)	L-295	Renard (J. P.). — Voir Ablart (G.)	L-219
Lynn (J. W.), Raggazoni (A.), Pynn (R.) and Joffrin (J.). —		Renard (J. P.). — Voir Velu (E.)	L-237
Observation of long range magnetic order in the		Ribault (M.). — Voir Fournel (A.)	L-445
reentrant superconductor HoMo ₆ S ₈	L-45	Ribault (M.). — Voir More (C.)	L-313
Malthête (J.). — Voir Nguyen Huu Tinh	L-417		L-513
Mamy (R.), Thiry (P.), Vachier (G.) et Couget (A.). — Struc-	2 11	Ribault (M.). — Voir Schulz (H. J.)	L-91
			191
ture de bandes expérimentale de l'aS1 I déterminée			
ture de bandes expérimentale de TaS ₂ -1 T déterminée	179	Rivory (J.), Frigerio (J. M.) et Rebouillat (J. P.). — Contri-	
par photoémission angulaire	L-79	Rivory (J.), Frigerio (J. M.) et Rebouillat (J. P.). — Contribution à la détermination de la structure électronique	27.
par photoémission angulaire	L-79	Rivory (J.), Frigerio (J. M.) et Rebouillat (J. P.). — Contribution à la détermination de la structure électronique d'alliages métalliques amorphes Cu ₅₀ Zr ₅₀ par spec-	
par photoémission angulaire		Rivory (J.), Frigerio (J. M.) et Rebouillat (J. P.). — Contribution à la détermination de la structure électronique d'alliages métalliques amorphes Cu ₅₀ Zr ₅₀ par spectroscopie optique	L-481
par photoémission angulaire	L-473	Rivory (J.), Frigerio (J. M.) et Rebouillat (J. P.). — Contribution à la détermination de la structure électronique d'alliages métalliques amorphes Cu ₅₀ Zr ₅₀ par spectroscopie optique	L-481 L-301
par photoémission angulaire. Mamy (R.), Thomas (J.), Jezequel (G.) et Lemonnier (J. C.). — Anisotropie des transitions interbandes dans le nitrure de bore hexagonal. Manes (L.). — Voir Boeuf (A.).	L-473 L-401	Rivory (J.), Frigerio (J. M.) et Rebouillat (J. P.). — Contribution à la détermination de la structure électronique d'alliages métalliques amorphes Cu ₅₀ Zr ₅₀ par spectroscopie optique	L-481 L-301 L-445
par photoemission angulaire. Mamy (R.), Thomas (J.), Jezequel (G.) et Lemonnier (J. C.). — Anisotropie des transitions interbandes dans le nitrure de bore hexagonal. Manes (L.). — Voir Boeuf (A.). Martins (A. F.). — Voir Volino (F.).	L-473 L-401 L-305	Rivory (J.), Frigerio (J. M.) et Rebouillat (J. P.). — Contribution à la détermination de la structure électronique d'alliages métalliques amorphes Cu ₅₀ Zr ₅₀ par spectroscopie optique	L-481 L-301 L-445 L-313
par photoemission angulaire. Mamy (R.), Thomas (J.), Jezequel (G.) et Lemonnier (J. C.). — Anisotropie des transitions interbandes dans le nitrure de bore hexagonal. Manes (L.). — Voir Boeuf (A.) Martins (A. F.). — Voir Volino (F.). Matecki (M.). — Voir Doussineau (P.)	L-473 L-401 L-305 L-267	Rivory (J.), Frigerio (J. M.) et Rebouillat (J. P.). — Contribution à la détermination de la structure électronique d'alliages métalliques amorphes Cu ₅₀ Zr ₅₀ par spectroscopie optique	L-481 L-301 L-445 L-313 L-295
par photoémission angulaire. Mamy (R.), Thomas (J.), Jezequel (G.) et Lemonnier (J. C.). — Anisotropie des transitions interbandes dans le nitrure de bore hexagonal. Manes (L.). — Voir Boeuf (A.). Martins (A. F.). — Voir Volino (F.). Matecki (M.). — Voir Doussineau (P.) Maurel (E.). — Voir Lapujoulade (J.)	L-473 L-401 L-305	Rivory (J.), Frigerio (J. M.) et Rebouillat (J. P.). — Contribution à la détermination de la structure électronique d'alliages métalliques amorphes Cu ₅₀ Zr ₅₀ par spectroscopie optique	L-481 L-301 L-445 L-313 L-295 L-197
par photoémission angulaire. Mamy (R.), Thomas (J.), Jezequel (G.) et Lemonnier (J. C.). — Anisotropie des transitions interbandes dans le nitrure de bore hexagonal. Manes (L.). — Voir Boeuf (A.). Martins (A. F.). — Voir Volino (F.). Matecki (M.). — Voir Doussineau (P.). Maurel (E.). — Voir Lapujoulade (J.). Maurice (F.), Froyer (G.), Miner (M.) and Gauneau (M.). —	L-473 L-401 L-305 L-267	Rivory (J.), Frigerio (J. M.) et Rebouillat (J. P.). — Contribution à la détermination de la structure électronique d'alliages métalliques amorphes Cu ₅₀ Zr ₅₀ par spectroscopie optique	L-481 L-301 L-445 L-313 L-295
par photoémission angulaire. Mamy (R.), Thomas (J.), Jezequel (G.) et Lemonnier (J. C.). — Anisotropie des transitions interbandes dans le nitrure de bore hexagonal. Manes (L.). — Voir Boeuf (A.). Martins (A. F.). — Voir Volino (F.). Matecki (M.). — Voir Doussineau (P.). Maurel (E.). — Voir Lapujoulade (J.). Maurice (F.), Froyer (G.), Miner (M.) and Gauneau (M.). — Correlation between D.C. bulk conductivity and impu-	L-473 L-401 L-305 L-267 L-463	Rivory (J.), Frigerio (J. M.) et Rebouillat (J. P.). — Contribution à la détermination de la structure électronique d'alliages métalliques amorphes Cu ₅₀ Zr ₅₀ par spectroscopie optique	L-481 L-301 L-445 L-313 L-295 L-197 L-323
par photoémission angulaire. Mamy (R.), Thomas (J.), Jezequel (G.) et Lemonnier (J. C.). — Anisotropie des transitions interbandes dans le nitrure de bore hexagonal. Manes (L.). — Voir Boeuf (A.) Martins (A. F.). — Voir Volino (F.). Matecki (M.). — Voir Doussineau (P.) Maurel (E.). — Voir Lapujoulade (J.) Maurice (F.), Froyer (G.), Miner (M.) and Gauneau (M.). — Correlation between D.C. bulk conductivity and impurities in undoped poly (p-phenylene).	L-473 L-401 L-305 L-267 L-463	Rivory (J.), Frigerio (J. M.) et Rebouillat (J. P.). — Contribution à la détermination de la structure électronique d'alliages métalliques amorphes Cu ₅₀ Zr ₅₀ par spectroscopie optique Robert (H.). — Voir Granier (W.) Roger (G.). — Voir Fournel (A.) Roger (G.). — Voir More (C.) Rolland (M.). — Voir Bernier (P.) Romagnan (J. P.). — Voir Laheurte (J. P.) Rondelez (F.). — Voir Vidakovic (P.) Rosenblatt (C.). — Magnetically induced nematic-smectic A tricritical point.	L-481 L-301 L-445 L-313 L-295 L-197 L-323
par photoémission angulaire Mamy (R.), Thomas (J.), Jezequel (G.) et Lemonnier (J. C.). — Anisotropie des transitions interbandes dans le nitrure de bore hexagonal. Manes (L.). — Voir Boeuf (A.) Martins (A. F.). — Voir Volino (F.). Matecki (M.). — Voir Doussineau (P.) Maurel (E.). — Voir Lapujoulade (J.) Maurice (F.), Froyer (G.), Miner (M.) and Gauneau (M.). — Correlation between D.C. bulk conductivity and impurities in undoped poly (p-phenylene). Mazaud (A.). — Voir Schulz (H. J.)	L-473 L-401 L-305 L-267 L-463 L-425 L-51	Rivory (J.), Frigerio (J. M.) et Rebouillat (J. P.). — Contribution à la détermination de la structure électronique d'alliages métalliques amorphes Cu ₅₀ Zr ₅₀ par spectroscopie optique	L-481 L-301 L-445 L-313 L-295 L-197 L-323
par photoemission angulaire. Mamy (R.), Thomas (J.), Jezequel (G.) et Lemonnier (J. C.). — Anisotropie des transitions interbandes dans le nitrure de bore hexagonal. Manes (L.). — Voir Boeuf (A.). Martins (A. F.). — Voir Volino (F.). Matecki (M.). — Voir Doussineau (P.) Maurel (E.). — Voir Lapujoulade (J.) Maurice (F.), Froyer (G.), Miner (M.) and Gauneau (M.). — Correlation between D.C. bulk conductivity and impurities in undoped poly (p-phenylene). Mazaud (A.). — Voir Schulz (H. J.). Meerschaut (A.). — Voir Mihaly (G.)	L-473 L-401 L-305 L-267 L-463	Rivory (J.), Frigerio (J. M.) et Rebouillat (J. P.). — Contribution à la détermination de la structure électronique d'alliages métalliques amorphes Cu ₅₀ Zr ₅₀ par spectroscopie optique	L-481 L-301 L-445 L-313 L-295 L-197 L-323
par photoemission angulaire. Mamy (R.), Thomas (J.), Jezequel (G.) et Lemonnier (J. C.). — Anisotropie des transitions interbandes dans le nitrure de bore hexagonal. Manes (L.). — Voir Boeuf (A.). Martins (A. F.). — Voir Volino (F.). Matecki (M.). — Voir Doussineau (P.) Maurel (E.). — Voir Lapujoulade (J.) Maurice (F.), Froyer (G.), Miner (M.) and Gauneau (M.). — Correlation between D.C. bulk conductivity and impurities in undoped poly (p-phenylene). Mazaud (A.). — Voir Schulz (H. J.) Meerschaut (A.). — Voir Mihaly (G.) Mihaly (G.), Housseau (N.), Mutka (H.), Zuppiroli (L.),	L-473 L-401 L-305 L-267 L-463 L-425 L-51	Rivory (J.), Frigerio (J. M.) et Rebouillat (J. P.). — Contribution à la détermination de la structure électronique d'alliages métalliques amorphes Cu ₅₀ Zr ₅₀ par spectroscopie optique	L-481 L-301 L-445 L-313 L-295 L-197 L-323
par photoemission angulaire. Mamy (R.), Thomas (J.), Jezequel (G.) et Lemonnier (J. C.). — Anisotropie des transitions interbandes dans le nitrure de bore hexagonal. Manes (L.). — Voir Boeuf (A.). Martins (A. F.). — Voir Volino (F.). Matecki (M.). — Voir Doussineau (P.) Maurel (E.). — Voir Lapujoulade (J.) Maurice (F.), Froyer (G.), Miner (M.) and Gauneau (M.). — Correlation between D.C. bulk conductivity and impurities in undoped poly (p-phenylene). Mazaud (A.). — Voir Schulz (H. J.) Meerschaut (A.). — Voir Mihaly (G.) Mihaly (G.), Housseau (N.), Mutka (H.), Zuppiroli (L.), Pelissier (J.), Gressier (P.), Meerschaut (A.) and	L-473 L-401 L-305 L-267 L-463 L-425 L-51	Rivory (J.), Frigerio (J. M.) et Rebouillat (J. P.). — Contribution à la détermination de la structure électronique d'alliages métalliques amorphes Cu ₅₀ Zr ₅₀ par spectroscopie optique	L-481 L-301 L-445 L-313 L-295 L-197 L-323
par photoemission angulaire. Mamy (R.), Thomas (J.), Jezequel (G.) et Lemonnier (J. C.). — Anisotropie des transitions interbandes dans le nitrure de bore hexagonal. Manes (L.). — Voir Boeuf (A.). Martins (A. F.). — Voir Volino (F.). Matecki (M.). — Voir Doussineau (P.) Maurel (E.). — Voir Lapujoulade (J.) Maurice (F.), Froyer (G.), Miner (M.) and Gauneau (M.). — Correlation between D.C. bulk conductivity and impurities in undoped poly (p-phenylene). Mazaud (A.). — Voir Schulz (H. J.) Meerschaut (A.). — Voir Mihaly (G.) Mihaly (G.), Housseau (N.), Mutka (H.), Zuppiroli (L.), Pelissier (J.), Gressier (P.), Meerschaut (A.) and Rouxel (J.). — Disorder in the quasi-one-dimensional	L-473 L-401 L-305 L-267 L-463 L-425 L-51 L-263	Rivory (J.), Frigerio (J. M.) et Rebouillat (J. P.). — Contribution à la détermination de la structure électronique d'alliages métalliques amorphes Cu ₅₀ Zr ₅₀ par spectroscopie optique	L-481 L-301 L-445 L-313 L-295 L-197 L-323
par photoémission angulaire. Mamy (R.), Thomas (J.), Jezequel (G.) et Lemonnier (J. C.). — Anisotropie des transitions interbandes dans le nitrure de bore hexagonal. Manes (L.). — Voir Boeuf (A.). Martins (A. F.). — Voir Volino (F.). Matecki (M.). — Voir Doussineau (P.). Maurel (E.). — Voir Lapujoulade (J.). Maurice (F.), Froyer (G.), Miner (M.) and Gauneau (M.). — Correlation between D.C. bulk conductivity and impurities in undoped poly (p-phenylene). Mazaud (A.). — Voir Schulz (H. J.). Meerschaut (A.). — Voir Mihaly (G.). Mihaly (G.), Housseau (N.), Mutka (H.), Zuppiroli (L.), Pelissier (J.), Gressier (P.), Meerschaut (A.) and Rouxel (J.). — Disorder in the quasi-one-dimensional conductor monoclinic tantalum trisulfide TaS ₃	L-473 L-401 L-305 L-267 L-463 L-425 L-51 L-263	Rivory (J.), Frigerio (J. M.) et Rebouillat (J. P.). — Contribution à la détermination de la structure électronique d'alliages métalliques amorphes Cu ₅₀ Zr ₅₀ par spectroscopie optique	L-481 L-301 L-445 L-313 L-295 L-197 L-323 L-9 L-263 L-401
par photoémission angulaire. Mamy (R.), Thomas (J.), Jezequel (G.) et Lemonnier (J. C.). — Anisotropie des transitions interbandes dans le nitrure de bore hexagonal. Manes (L.). — Voir Boeuf (A.). Martins (A. F.). — Voir Volino (F.). Matecki (M.). — Voir Doussineau (P.). Maurel (E.). — Voir Lapujoulade (J.). Maurice (F.), Froyer (G.), Miner (M.) and Gauneau (M.). — Correlation between D.C. bulk conductivity and impurities in undoped poly (p-phenylene). Mazaud (A.). — Voir Schulz (H. J.). Meerschaut (A.). — Voir Mihaly (G.). Mihaly (G.), Housseau (N.), Mutka (H.), Zuppiroli (L.), Pelissier (J.), Gressier (P.), Meerschaut (A.) and Rouxel (J.). — Disorder in the quasi-one-dimensional conductor monoclinic tantalum trisulfide TaS ₃ Miner (M.). — Voir Maurice (F.)	L-473 L-401 L-305 L-267 L-463 L-425 L-51 L-263 L-263 L-263	Rivory (J.), Frigerio (J. M.) et Rebouillat (J. P.). — Contribution à la détermination de la structure électronique d'alliages métalliques amorphes Cu ₅₀ Zr ₅₀ par spectroscopie optique	L-481 L-301 L-445 L-313 L-295 L-197 L-323 L-9 L-263 L-401
par photoémission angulaire. Mamy (R.), Thomas (J.), Jezequel (G.) et Lemonnier (J. C.). — Anisotropie des transitions interbandes dans le nitrure de bore hexagonal. Manes (L.). — Voir Boeuf (A.). Martins (A. F.). — Voir Volino (F.). Matecki (M.). — Voir Doussineau (P.) Maurel (E.). — Voir Lapujoulade (J.) Maurice (F.), Froyer (G.), Miner (M.) and Gauneau (M.). — Correlation between D.C. bulk conductivity and impurities in undoped poly (p-phenylene). Mazaud (A.). — Voir Schulz (H. J.). Meerschaut (A.). — Voir Mihaly (G.). Mihaly (G.), Housseau (N.), Mutka (H.), Zuppiroli (L.), Pelissier (J.), Gressier (P.), Meerschaut (A.) and Rouxel (J.). — Disorder in the quasi-one-dimensional conductor monoclinic tantalum trisulfide TaS ₃ Miner (M.). — Voir Maurice (F.). Miranday (J. P.). — Voir Velu (E.)	L-473 L-401 L-305 L-267 L-463 L-425 L-51 L-263 L-263 L-425 L-237	Rivory (J.), Frigerio (J. M.) et Rebouillat (J. P.). — Contribution à la détermination de la structure électronique d'alliages métalliques amorphes Cu ₅₀ Zr ₅₀ par spectroscopie optique Robert (H.). — Voir Granier (W.) Roger (G.). — Voir Fournel (A.) Roger (G.). — Voir More (C.) Rolland (M.). — Voir Bernier (P.) Romagnan (J. P.). — Voir Laheurte (J. P.) Rondelez (F.). — Voir Vidakovic (P.) Rosenblatt (C.). — Magnetically induced nematic-smectic A tricritical point. Rouxel (J.). — Voir Mihaly (G.) Rustichelli (F.). — Voir Boeuf (A.) Sadoc (J. F.) et Laridjani (M.). — Evolution structurale d'alliages Pd-Si amorphes en fonction de la température. Salin (D.) and Schon (W.). — Acoustics of water saturated packed glass spheres	L-481 L-301 L-445 L-313 L-295 L-197 L-323 L-9 L-263 L-401
par photoemission angulaire. Mamy (R.), Thomas (J.), Jezequel (G.) et Lemonnier (J. C.). — Anisotropie des transitions interbandes dans le nitrure de bore hexagonal. Manes (L.). — Voir Boeuf (A.). Martins (A. F.). — Voir Volino (F.). Matecki (M.). — Voir Doussineau (P.) Maurel (E.). — Voir Lapujoulade (J.) Maurice (F.), Froyer (G.), Miner (M.) and Gauneau (M.). — Correlation between D.C. bulk conductivity and impurities in undoped poly (p-phenylene). Mazaud (A.). — Voir Schulz (H. J.). Meerschaut (A.). — Voir Mihaly (G.) Mihaly (G.), Housseau (N.), Mutka (H.), Zuppiroli (L.), Pelissier (J.), Gressier (P.), Meerschaut (A.) and Rouxel (J.). — Disorder in the quasi-one-dimensional conductor monoclinic tantalum trisulfide TaS3 Miner (M.). — Voir Maurice (F.). Miranday (J. P.). — Voir Velu (E.). Mishonov (T.). — Voir Sariisky (Tz.)	L-473 L-401 L-305 L-267 L-463 L-425 L-51 L-263 L-263 L-263	Rivory (J.), Frigerio (J. M.) et Rebouillat (J. P.). — Contribution à la détermination de la structure électronique d'alliages métalliques amorphes Cu ₅₀ Zr ₅₀ par spectroscopie optique	L-481 L-301 L-445 L-313 L-295 L-197 L-323 L-9 L-263 L-401
par photoemission angulaire. Mamy (R.), Thomas (J.), Jezequel (G.) et Lemonnier (J. C.). — Anisotropie des transitions interbandes dans le nitrure de bore hexagonal. Manes (L.). — Voir Boeuf (A.). Martins (A. F.). — Voir Volino (F.). Matecki (M.). — Voir Doussineau (P.) Maurice (E.). — Voir Lapujoulade (J.) Maurice (F.), Froyer (G.), Miner (M.) and Gauneau (M.). — Correlation between D.C. bulk conductivity and impurities in undoped poly (p-phenylene). Mazaud (A.). — Voir Schulz (H. J.). Meerschaut (A.). — Voir Mihaly (G.). Mihaly (G.), Housseau (N.), Mutka (H.), Zuppiroli (L.), Pelissier (J.), Gressier (P.), Meerschaut (A.) and Rouxel (J.). — Disorder in the quasi-one-dimensional conductor monoclinic tantalum trisulfide TaS3 Miner (M.). — Voir Maurice (F.). Miranday (J. P.). — Voir Velu (E.). Mishonov (T.). — Voir Sariisky (Tz.) More (C.), Roger (G.), Sorbier (J. P.), Jérome (D.), Ribault	L-473 L-401 L-305 L-267 L-463 L-425 L-51 L-263 L-263 L-425 L-237	Rivory (J.), Frigerio (J. M.) et Rebouillat (J. P.). — Contribution à la détermination de la structure électronique d'alliages métalliques amorphes Cu ₅₀ Zr ₅₀ par spectroscopie optique	L-481 L-301 L-445 L-313 L-295 L-197 L-323 L-9 L-263 L-401 L-485 L-477
par photoémission angulaire. Mamy (R.), Thomas (J.), Jezequel (G.) et Lemonnier (J. C.). — Anisotropie des transitions interbandes dans le nitrure de bore hexagonal. Manes (L.). — Voir Boeuf (A.). Martins (A. F.). — Voir Volino (F.). Matecki (M.). — Voir Doussineau (P.) Maurice (E.). — Voir Lapujoulade (J.) Maurice (F.), Froyer (G.), Miner (M.) and Gauneau (M.). — Correlation between D.C. bulk conductivity and impurities in undoped poly (p-phenylene). Mazaud (A.). — Voir Schulz (H. J.) Meerschaut (A.). — Voir Mihaly (G.) Mihaly (G.), Housseau (N.), Mutka (H.), Zuppiroli (L.), Pelissier (J.), Gressier (P.), Meerschaut (A.) and Rouxel (J.). — Disorder in the quasi-one-dimensional conductor monoclinic tantalum trisulfide TaS3 Miner (M.). — Voir Maurice (F.) Mishonov (T.). — Voir Velu (E.) Mishonov (T.). — Voir Velu (E.) Mishonov (T.). — Voir Scriisky (Tz.) More (C.), Roger (G.), Sorbier (J. P.), Jérome (D.), Ribault (M.) and Bechgaard (K.). — One-dimensional organic	L-473 L-401 L-305 L-267 L-463 L-425 L-51 L-263 L-263 L-425 L-237	Rivory (J.), Frigerio (J. M.) et Rebouillat (J. P.). — Contribution à la détermination de la structure électronique d'alliages métalliques amorphes Cu ₅₀ Zr ₅₀ par spectroscopie optique	L-481 L-301 L-445 L-313 L-295 L-197 L-323 L-9 L-263 L-401 L-485 L-477 L-503
par photoémission angulaire. Mamy (R.), Thomas (J.), Jezequel (G.) et Lemonnier (J. C.). — Anisotropie des transitions interbandes dans le nitrure de bore hexagonal Manes (L.). — Voir Boeuf (A.). Martins (A. F.). — Voir Volino (F.). Matecki (M.). — Voir Doussineau (P.) Maurel (E.). — Voir Lapujoulade (J.) Maurice (F.), Froyer (G.), Miner (M.) and Gauneau (M.). — Correlation between D.C. bulk conductivity and impurities in undoped poly (p-phenylene) Mazaud (A.). — Voir Schulz (H. J.) Meerschaut (A.). — Voir Mihaly (G.) Mihaly (G.), Housseau (N.), Mutka (H.), Zuppiroli (L.), Pelissier (J.), Gressier (P.), Meerschaut (A.) and Rouxel (J.). — Disorder in the quasi-one-dimensional conductor monoclinic tantalum trisulfide TaS3 Miner (M.). — Voir Maurice (F.) Miranday (J. P.). — Voir Velu (E.) Mishonov (T.). — Voir Sariisky (Tz.) More (C.), Roger (G.), Sorbier (J. P.), Jérome (D.), Ribault (M.) and Bechgaard (K.). — One-dimensional organic superconductivity: an investigation through electron	L-473 L-401 L-305 L-267 L-463 L-425 L-51 L-263 L-263 L-425 L-237	Rivory (J.), Frigerio (J. M.) et Rebouillat (J. P.). — Contribution à la détermination de la structure électronique d'alliages métalliques amorphes Cu ₅₀ Zr ₅₀ par spectroscopie optique	L-481 L-301 L-445 L-312 L-295 L-197 L-323 L-401 L-485 L-477 L-503 L-155 L-477
par photoémission angulaire. Mamy (R.), Thomas (J.), Jezequel (G.) et Lemonnier (J. C.). — Anisotropie des transitions interbandes dans le nitrure de bore hexagonal. Manes (L.). — Voir Boeuf (A.). Martins (A. F.). — Voir Volino (F.). Matecki (M.). — Voir Doussineau (P.). Maurel (E.). — Voir Lapujoulade (J.). Maurice (F.), Froyer (G.), Miner (M.) and Gauneau (M.). — Correlation between D.C. bulk conductivity and impurities in undoped poly (p-phenylene). Mazaud (A.). — Voir Schulz (H. J.). Meerschaut (A.). — Voir Mihaly (G.). Mihaly (G.), Housseau (N.), Mutka (H.), Zuppiroli (L.), Pelissier (J.), Gressier (P.), Meerschaut (A.) and Rouxel (J.). — Disorder in the quasi-one-dimensional conductor monoclinic tantalum trisulfide TaS3 Miner (M.). — Voir Maurice (F.). Miranday (J. P.). — Voir Velu (E.). Miranday (J. P.). — Voir Sariisky (Tz.). More (C.), Roger (G.), Sorbier (J. P.), Jérome (D.), Ribault (M.) and Bechgaard (K.). — One-dimensional organic superconductivity: an investigation through electron Schottky tunnelling in N/GaSb-(TMTSF)2PF6 junc-	L-473 L-401 L-305 L-267 L-463 L-425 L-51 L-263 L-263 L-425 L-237 L-503	Rivory (J.), Frigerio (J. M.) et Rebouillat (J. P.). — Contribution à la détermination de la structure électronique d'alliages métalliques amorphes Cu ₅₀ Zr ₅₀ par spectroscopie optique	L-481 L-301 L-445 L-312 L-295 L-197 L-323 L-401 L-461 L-475 L-503 L-153
par photoémission angulaire. Mamy (R.), Thomas (J.), Jezequel (G.) et Lemonnier (J. C.). — Anisotropie des transitions interbandes dans le nitrure de bore hexagonal. Manes (L.). — Voir Boeuf (A.). Martins (A. F.). — Voir Volino (F.). Matecki (M.). — Voir Doussineau (P.) Maurice (E.). — Voir Lapujoulade (J.) Maurice (F.), Froyer (G.), Miner (M.) and Gauneau (M.). — Correlation between D.C. bulk conductivity and impurities in undoped poly (p-phenylene). Mazaud (A.). — Voir Schulz (H. J.). Meerschaut (A.). — Voir Mihaly (G.) Mihaly (G.), Housseau (N.), Mutka (H.), Zuppiroli (L.), Pelissier (J.), Gressier (P.), Meerschaut (A.) and Rouxel (J.). — Disorder in the quasi-one-dimensional conductor monoclinic tantalum trisulfide TaS3 Miner (M.). — Voir Maurice (F.). Mishonov (T.). — Voir Sariisky (Tz.). More (C.), Roger (G.), Sorbier (J. P.), Jérome (D.), Ribault (M.) and Bechgaard (K.). — One-dimensional organic superconductivity: an investigation through electron Schottky tunnelling in N/GaSb-(TMTSF) ₂ PF ₆ junctions under pressure.	L-473 L-401 L-305 L-267 L-463 L-425 L-51 L-263 L-263 L-425 L-237 L-503	Rivory (J.), Frigerio (J. M.) et Rebouillat (J. P.). — Contribution à la détermination de la structure électronique d'alliages métalliques amorphes Cu ₅₀ Zr ₅₀ par spectroscopie optique	L-481 L-301 L-445 L-312 L-295 L-197 L-323 L-401 L-485 L-477 L-503 L-155 L-477
par photoemission angulaire. Mamy (R.), Thomas (J.), Jezequel (G.) et Lemonnier (J. C.). — Anisotropie des transitions interbandes dans le nitrure de bore hexagonal. Manes (L.). — Voir Boeuf (A.). Martins (A. F.). — Voir Volino (F.). Matecki (M.). — Voir Doussineau (P.) Maurice (F.), Froyer (G.), Miner (M.) and Gauneau (M.). — Correlation between D.C. bulk conductivity and impurities in undoped poly (p-phenylene). Mazaud (A.). — Voir Schulz (H. J.). Meerschaut (A.). — Voir Mihaly (G.) Mihaly (G.), Housseau (N.), Mutka (H.), Zuppiroli (L.), Pelissier (J.), Gressier (P.), Meerschaut (A.) and Rouxel (J.). — Disorder in the quasi-one-dimensional conductor monoclinic tantalum trisulfide TaS3 Miner (M.). — Voir Maurice (F.). Miranday (J. P.). — Voir Velu (E.). Mishonov (T.). — Voir Sariisky (Tz.) More (C.), Roger (G.), Sorbier (J. P.), Jérome (D.), Ribault (M.) and Bechgaard (K.). — One-dimensional organic superconductivity: an investigation through electron Schottky tunnelling in N/GaSb-(TMTSF) ₂ PF ₆ junctions under pressure. More (C.). — Voir Fournel (A.)	L-473 L-401 L-305 L-267 L-463 L-425 L-51 L-263 L-263 L-425 L-237 L-503	Rivory (J.), Frigerio (J. M.) et Rebouillat (J. P.). — Contribution à la détermination de la structure électronique d'alliages métalliques amorphes Cu ₅₀ Zr ₅₀ par spectroscopie optique	L-481 L-301 L-445 L-312 L-295 L-197 L-323 L-401 L-485 L-477 L-503 L-155 L-477
par photoémission angulaire. Mamy (R.), Thomas (J.), Jezequel (G.) et Lemonnier (J. C.). — Anisotropie des transitions interbandes dans le nitrure de bore hexagonal. Manes (L.). — Voir Boeuf (A.). Martins (A. F.). — Voir Volino (F.). Matecki (M.). — Voir Doussineau (P.) Maurice (E.). — Voir Lapujoulade (J.) Maurice (F.), Froyer (G.), Miner (M.) and Gauneau (M.). — Correlation between D.C. bulk conductivity and impurities in undoped poly (p-phenylene). Mazaud (A.). — Voir Schulz (H. J.). Meerschaut (A.). — Voir Mihaly (G.). Mihaly (G.), Housseau (N.), Mutka (H.), Zuppiroli (L.), Pelissier (J.), Gressier (P.), Meerschaut (A.) and Rouxel (J.). — Disorder in the quasi-one-dimensional conductor monoclinic tantalum trisulfide TaS3 Miner (M.). — Voir Maurice (F.). Miranday (J. P.). — Voir Velu (E.). Mishonov (T.). — Voir Sariisky (Tz.). More (C.), Roger (G.), Sorbier (J. P.), Jérome (D.), Ribault (M.) and Bechgaard (K.). — One-dimensional organic superconductivity: an investigation through electron Schottky tunnelling in N/GaSb-(TMTSF)2PF6 junctions under pressure. More (C.). — Voir Fournel (A.). Moret (R.). — Voir Pouget (J. P.).	L-473 L-401 L-305 L-267 L-463 L-425 L-51 L-263 L-263 L-425 L-237 L-503	Rivory (J.), Frigerio (J. M.) et Rebouillat (J. P.). — Contribution à la détermination de la structure électronique d'alliages métalliques amorphes Cu ₅₀ Zr ₅₀ par spectroscopie optique	L-481 L-301 L-445 L-312 L-295 L-197 L-322 L-263 L-401 L-485 L-477 L-501 L-131
par photoémission angulaire. Mamy (R.), Thomas (J.), Jezequel (G.) et Lemonnier (J. C.). — Anisotropie des transitions interbandes dans le nitrure de bore hexagonal. Manes (L.). — Voir Boeuf (A.). Martins (A. F.). — Voir Volino (F.). Matecki (M.). — Voir Doussineau (P.) Maurel (E.). — Voir Lapujoulade (J.) Maurice (F.), Froyer (G.), Miner (M.) and Gauneau (M.). — Correlation between D.C. bulk conductivity and impurities in undoped poly (p-phenylene). Mazaud (A.). — Voir Schulz (H. J.). Meerschaut (A.). — Voir Mihaly (G.). Mihaly (G.), Housseau (N.), Mutka (H.), Zuppiroli (L.), Pelissier (J.), Gressier (P.), Meerschaut (A.) and Rouxel (J.). — Disorder in the quasi-one-dimensional conductor monoclinic tantalum trisulfide TaS3 Miner (M.). — Voir Maurice (F.). Mishonov (T.). — Voir Velu (E.). Mishonov (T.). — Voir Sariisky (Tz.). More (C.), Roger (G.), Sorbier (J. P.), Jérome (D.), Ribault (M.) and Bechgaard (K.). — One-dimensional organic superconductivity: an investigation through electron Schottky tunnelling in N/GaSb-(TMTSF)2PF6 junctions under pressure. More (C.). — Voir Fournel (A.) Moret (R.). — Voir Pouget (J. P.). Moussa (F.). — Voir Benattar (J. J.).	L-473 L-401 L-305 L-267 L-463 L-425 L-51 L-263 L-263 L-425 L-237 L-503	Rivory (J.), Frigerio (J. M.) et Rebouillat (J. P.). — Contribution à la détermination de la structure électronique d'alliages métalliques amorphes Cu ₅₀ Zr ₅₀ par spectroscopie optique	L-481 L-301 L-445 L-312 L-295 L-197 L-323 L-96 L-401 L-485 L-477 L-503 L-155 L-477 L-131
par photoémission angulaire. Mamy (R.), Thomas (J.), Jezequel (G.) et Lemonnier (J. C.). — Anisotropie des transitions interbandes dans le nitrure de bore hexagonal. Manes (L.). — Voir Boeuf (A.). Martins (A. F.). — Voir Volino (F.). Matecki (M.). — Voir Doussineau (P.) Maurice (E.). — Voir Lapujoulade (J.) Maurice (F.), Froyer (G.), Miner (M.) and Gauneau (M.). — Correlation between D.C. bulk conductivity and impurities in undoped poly (p-phenylene). Mazaud (A.). — Voir Schulz (H. J.). Meerschaut (A.). — Voir Mihaly (G.). Mihaly (G.), Housseau (N.), Mutka (H.), Zuppiroli (L.), Pelissier (J.), Gressier (P.), Meerschaut (A.) and Rouxel (J.). — Disorder in the quasi-one-dimensional conductor monoclinic tantalum trisulfide TaS3 Miner (M.). — Voir Maurice (F.). Miranday (J. P.). — Voir Velu (E.). Mishonov (T.). — Voir Sariisky (Tz.). More (C.), Roger (G.), Sorbier (J. P.), Jérome (D.), Ribault (M.) and Bechgaard (K.). — One-dimensional organic superconductivity: an investigation through electron Schottky tunnelling in N/GaSb-(TMTSF)2PF6 junctions under pressure. More (C.). — Voir Fournel (A.). Moret (R.). — Voir Pouget (J. P.).	L-473 L-401 L-305 L-267 L-463 L-425 L-263 L-263 L-263 L-251 L-263 L-425 L-237 L-503	Rivory (J.), Frigerio (J. M.) et Rebouillat (J. P.). — Contribution à la détermination de la structure électronique d'alliages métalliques amorphes Cu ₅₀ Zr ₅₀ par spectroscopie optique	L-481 L-301 L-445 L-312 L-295 L-197 L-323 L-90 L-401 L-485 L-477 L-503 L-155 L-477 L-131 L-51 L-51 L-35
par photoemission angulaire. Mamy (R.), Thomas (J.), Jezequel (G.) et Lemonnier (J. C.). — Anisotropie des transitions interbandes dans le nitrure de bore hexagonal. Manes (L.). — Voir Boeuf (A.). Martins (A. F.). — Voir Volino (F.). Matecki (M.). — Voir Doussineau (P.) Maurice (E.). — Voir Lapujoulade (J.) Maurice (F.), Froyer (G.), Miner (M.) and Gauneau (M.). — Correlation between D.C. bulk conductivity and impurities in undoped poly (p-phenylene). Mazaud (A.). — Voir Schulz (H. J.). Meerschaut (A.). — Voir Mihaly (G.). Mihaly (G.), Housseau (N.), Mutka (H.), Zuppiroli (L.), Pelissier (J.), Gressier (P.), Meerschaut (A.) and Rouxel (J.). — Disorder in the quasi-one-dimensional conductor monoclinic tantalum trisulfide TaS3 Miner (M.). — Voir Maurice (F.). Miranday (J. P.). — Voir Velu (E.). Mishonov (T.). — Voir Sariisky (T.). More (C.), Roger (G.), Sorbier (J. P.), Jérome (D.), Ribault (M.) and Bechgaard (K.). — One-dimensional organic superconductivity: an investigation through electron Schottky tunnelling in N/GaSb-(TMTSF)2PF6 junctions under pressure. More (C.). — Voir Fournel (A.). Moret (R.). — Voir Pouget (J. P.) Moussa (F.). — Voir Benattar (J. J.). Mutka (H.). — Voir Mihaly (G.).	L-473 L-401 L-305 L-267 L-463 L-425 L-263 L-263 L-263 L-313 L-445 L-543 L-67 L-263	Rivory (J.), Frigerio (J. M.) et Rebouillat (J. P.). — Contribution à la détermination de la structure électronique d'alliages métalliques amorphes Cu ₅₀ Zr ₅₀ par spectroscopie optique	L-481 L-301 L-445 L-312 L-295 L-197 L-323 L-96 L-401 L-485 L-477 L-503 L-155 L-477 L-131
par photoémission angulaire. Mamy (R.), Thomas (J.), Jezequel (G.) et Lemonnier (J. C.). — Anisotropie des transitions interbandes dans le nitrure de bore hexagonal. Manes (L.). — Voir Boeuf (A.). Martins (A. F.). — Voir Volino (F.). Matecki (M.). — Voir Doussineau (P.) Maurice (E.). — Voir Lapujoulade (J.) Maurice (F.), Froyer (G.), Miner (M.) and Gauneau (M.). — Correlation between D.C. bulk conductivity and impurities in undoped poly (p-phenylene). Mazaud (A.). — Voir Schulz (H. J.). Meerschaut (A.). — Voir Mihaly (G.). Mihaly (G.), Housseau (N.), Mutka (H.), Zuppiroli (L.), Pelissier (J.), Gressier (P.), Meerschaut (A.) and Rouxel (J.). — Disorder in the quasi-one-dimensional conductor monoclinic tantalum trisulfide TaS3 Miner (M.). — Voir Maurice (F.). Mishonov (T.). — Voir Velu (E.). Mishonov (T.). — Voir Sariisky (Tz.). More (C.), Roger (G.), Sorbier (J. P.), Jérome (D.), Ribault (M.) and Bechgaard (K.). — One-dimensional organic superconductivity: an investigation through electron Schottky tunnelling in N/GaSb-(TMTSF)2PF6 junctions under pressure. More (C.). — Voir Fournel (A.) Moret (R.). — Voir Pouget (J. P.) Moussa (F.). — Voir Benattar (J. J.) Mutka (H.). — Voir Mihaly (G.). Nedellec (P.). — Voir Sniadower (L.)	L-473 L-401 L-305 L-267 L-463 L-425 L-51 L-263 L-425 L-237 L-503	Rivory (J.), Frigerio (J. M.) et Rebouillat (J. P.). — Contribution à la détermination de la structure électronique d'alliages métalliques amorphes Cu ₅₀ Zr ₅₀ par spectroscopie optique	L-481 L-301 L-445 L-312 L-295 L-197 L-323 L-90 L-401 L-485 L-477 L-503 L-155 L-477 L-131 L-51 L-51 L-35
par photoémission angulaire. Mamy (R.), Thomas (J.), Jezequel (G.) et Lemonnier (J. C.). — Anisotropie des transitions interbandes dans le nitrure de bore hexagonal. Manes (L.). — Voir Boeuf (A.). Martins (A. F.). — Voir Volino (F.). Matecki (M.). — Voir Doussineau (P.) Maurice (E.). — Voir Lapujoulade (J.) Maurice (F.), Froyer (G.), Miner (M.) and Gauneau (M.). — Correlation between D.C. bulk conductivity and impurities in undoped poly (p-phenylene). Mazaud (A.). — Voir Schulz (H. J.). Meerschaut (A.). — Voir Mihaly (G.). Mihaly (G.), Housseau (N.), Mutka (H.), Zuppiroli (L.), Pelissier (J.), Gressier (P.), Meerschaut (A.) and Rouxel (J.). — Disorder in the quasi-one-dimensional conductor monoclinic tantalum trisulfide TaS3 Miner (M.). — Voir Maurice (F.). Mishonov (T.). — Voir Sariisky (Tz.). More (C.), Roger (G.), Sorbier (J. P.), Jérome (D.), Ribault (M.) and Bechgaard (K.). — One-dimensional organic superconductivity: an investigation through electron Schottky tunnelling in N/GaSb-(TMTSF)2PF6 junctions under pressure. More (C.). — Voir Fournel (A.). Moret (R.). — Voir Pouget (J. P.). Moussa (F.). — Voir Benattar (J. J.). Mutka (H.). — Voir Mihaly (G.). Nedellec (P.). — Voir Sniadower (L.). Nguyen Huu Tinh, Malthête (J.) and Destrade (C.). —	L-473 L-401 L-305 L-267 L-463 L-425 L-263 L-263 L-263 L-313 L-445 L-543 L-67 L-263	Rivory (J.), Frigerio (J. M.) et Rebouillat (J. P.). — Contribution à la détermination de la structure électronique d'alliages métalliques amorphes Cu ₅₀ Zr ₅₀ par spectroscopie optique	L-481 L-301 L-445 L-313 L-295 L-197 L-323 L-96 L-461 L-482 L-477 L-131 L-51 L-51 L-52 L-53 L-63
par photoémission angulaire. Mamy (R.), Thomas (J.), Jezequel (G.) et Lemonnier (J. C.). — Anisotropie des transitions interbandes dans le nitrure de bore hexagonal. Manes (L.). — Voir Boeuf (A.). Martins (A. F.). — Voir Volino (F.). Matecki (M.). — Voir Doussineau (P.) Maurel (E.). — Voir Lapujoulade (J.) Maurice (F.), Froyer (G.), Miner (M.) and Gauneau (M.). — Correlation between D.C. bulk conductivity and impurities in undoped poly (p-phenylene). Mazaud (A.). — Voir Schulz (H. J.). Meerschaut (A.). — Voir Mihaly (G.). Mihaly (G.), Housseau (N.), Mutka (H.), Zuppiroli (L.), Pelissier (J.), Gressier (P.), Meerschaut (A.) and Rouxel (J.). — Disorder in the quasi-one-dimensional conductor monoclinic tantalum trisulfide TaS3 Miner (M.). — Voir Maurice (F.). Mishonov (T.). — Voir Sariisky (Tz.). More (C.), Roger (G.), Sorbier (J. P.), Jérome (D.), Ribault (M.) and Bechgaard (K.). — One-dimensional organic superconductivity: an investigation through electron Schottky tunnelling in N/GaSb-(TMTSF) ₂ PF ₆ junctions under pressure. More (C.). — Voir Fournel (A.) Moret (R.). — Voir Fournel (A.) Moret (R.). — Voir Fournel (A.) Moret (R.). — Voir Benattar (J. J.) Mutka (H.). — Voir Mihaly (G.). Nedellec (P.). — Voir Sniadower (L.) Nguyen Huu Tinh, Malthête (J.) and Destrade (C.). — Reentrant nematic and columnar phases in disc-like	L-473 L-401 L-305 L-267 L-463 L-425 L-51 L-263 L-425 L-237 L-503 L-4313 L-445 L-543 L-67 L-263 L-13	Rivory (J.), Frigerio (J. M.) et Rebouillat (J. P.). — Contribution à la détermination de la structure électronique d'alliages métalliques amorphes Cu ₅₀ Zr ₅₀ par spectroscopie optique	L-481 L-301 L-445 L-312 L-295 L-197 L-323 L-96 L-461 L-477 L-503 L-477 L-131 L-51 L-51 L-52 L-477 L-131
par photoémission angulaire. Mamy (R.), Thomas (J.), Jezequel (G.) et Lemonnier (J. C.). — Anisotropie des transitions interbandes dans le nitrure de bore hexagonal. Manes (L.). — Voir Boeuf (A.). Martins (A. F.). — Voir Volino (F.). Matecki (M.). — Voir Doussineau (P.) Maurice (E.). — Voir Lapujoulade (J.) Maurice (F.), Froyer (G.), Miner (M.) and Gauneau (M.). — Correlation between D.C. bulk conductivity and impurities in undoped poly (p-phenylene). Mazaud (A.). — Voir Schulz (H. J.). Meerschaut (A.). — Voir Mihaly (G.). Mihaly (G.), Housseau (N.), Mutka (H.), Zuppiroli (L.), Pelissier (J.), Gressier (P.), Meerschaut (A.) and Rouxel (J.). — Disorder in the quasi-one-dimensional conductor monoclinic tantalum trisulfide TaS3 Miner (M.). — Voir Maurice (F.). Mishonov (T.). — Voir Sariisky (Tz.). More (C.), Roger (G.), Sorbier (J. P.), Jérome (D.), Ribault (M.) and Bechgaard (K.). — One-dimensional organic superconductivity: an investigation through electron Schottky tunnelling in N/GaSb-(TMTSF)2PF6 junctions under pressure. More (C.). — Voir Fournel (A.). Moret (R.). — Voir Pouget (J. P.). Moussa (F.). — Voir Benattar (J. J.). Mutka (H.). — Voir Mihaly (G.). Nedellec (P.). — Voir Sniadower (L.). Nguyen Huu Tinh, Malthête (J.) and Destrade (C.). —	L-473 L-401 L-305 L-267 L-463 L-425 L-263 L-263 L-263 L-313 L-445 L-543 L-67 L-263	Rivory (J.), Frigerio (J. M.) et Rebouillat (J. P.). — Contribution à la détermination de la structure électronique d'alliages métalliques amorphes Cu ₅₀ Zr ₅₀ par spectroscopie optique	L-481 L-301 L-445 L-313 L-295 L-197 L-323 L-96 L-461 L-482 L-477 L-131 L-51 L-51 L-52 L-53 L-63

Subtil (J. L.). — Voir Laporte (P.)	L-339	Walgraef (D.). — Voir Dewel (G.)	L-361
Thiry (P.). — Voir Mamy (R.)	L-79	Wiechert (H.). — Voir Tiby (C.)	L-155
	L-473	Zgonik (M.). — Voir Čopič (M.)	L-405
Tiby (C.), Wiechert (H.), Lauter (H. J.) and Schaab (J.). —		Zidler (B.). — Voir Chaillou (D.)	L-343
Absence of continuous growth for bee and hep solid		Zuppiroli (L.). — Voir Mihaly (G.)	L-263
³ He on Grafoil	L-155		
Trebin (HR.) and Kutka (R.). — Restricted homotopies			
and focal conic textures			
Vachier (G.). — Voir Mamy (R.)			
Vautier (C.). — Voir Viger (C.)	L-495	PHYSIQUE APPLIQUÉE	
Velu (E.), Renard (J. P.) and Miranday (J. P.). — Remanent		THISIQUE AFFLIQUEE	
magnetization and anisotropic susceptibility in the			
spin-glass phase of amorphous PbMnFeF ₇	L-237		
Veyssie (J. J.). — Voir Garoche (P.)		Cherpak (N. T.). — Voir Mishenko (V. V.)	
Vial (J. C.). — Voir Buisson (R.)	L-115	Fabre (G.). — Voir Zepp (G.)	L-175
Vidakovic (P.), Allain (C.) and Rondelez (F.). — Dynamic		Mishenko (V. V.) and Cherpak (N. T.). — Inversion ratio	
correlation length in polymer solutions at the theta		in trivalent-iron doped and alusite in the Q and V	
temperature	L-323	bands	L-145
Viger (C.), Fleury (G.) et Vautier (C.). — Fatigue photo-	Y 10.0	Saissy (A.). — Diffusion Raman stimulée d'une onde infra-	
	L-495	rouge à 1,06 μm dans une fibre optique	L-163
Volino (F.), Martins (A. F.), Blumstein (R. B.) and Blum-		Wick (A.). — Voir Zepp (G.)	L-175
stein (A.). — Evidence by proton NMR of the align-		Zepp (G.), Wick (A.) et Fabre (G.). — Atténuation d'une	
ment of a thermotropic nematic polymer by a magnetic	1 205	onde à la surface d'un conducteur entouré de diélec-	I 176
field of ~ 1 tesla	L-303	triques	L-175

.



REVUE DE PHYSIQUE APPLIQUÉE

Tome 16 — 1981

ARTICLES DE REVUE

Chilaya (G.). — Induction of chirality in nematic phases	193	Porteseil (J. L.). — Voir Vergne (R.)	449
Cotillard (J. C.). — Voir Vergne (R.).	449	Vedrine (J. C.). — Méthodes expérimentales de caractérisa-	
Dormann (J. L.). — Le phénomène de superparamagnétisme.	275	tion des catalyseurs industriels	373
Fauchais (P.). — Voir Kassabji (F.).	549	Vergne (R.), Cotillard (J. C.) et Porteseil (J. L.). — Quelques aspects statistiques des processus d'aimantation dans	
Kassabji (F.) et Fauchais (P.). — Les générateurs à plasma	549	les corps ferromagnétiques. Cas du déplacement d'une	
Martin (G. A.). — Détermination des tailles de particules métalliques et de leur distribution en catalyse hétéro-		seule paroi de Bloch à 180° dans un milieu monocristallin	
gènegène	181	aléatoirement perturbé	449
gene	101	1	
PHYSIQ	UE	APPLIQUÉE	
PHYSIQUE ET MÉCANIQUE		Gourrier (S.). — Voir Dimitriou (P.)	419
DES MATÉRIAUX		Hage-Ali (M.) and Siffert (P.). — Electrostatic analysis of	
		back-scattered heavy ions for semiconductor surface	
Allay (L.). — Voir Bosca (G.)	387	investigation	165
Attolini (G.). — Voir Franchi (S.)	1	Hallais (J.). — Voir Laurence (G.)	
Bonnet (R.). — Voir Gaymard (P.)	145	Hottier (F.). — Voir Laurence (G.)	
Bosca (G.), David (J.), Allay (L.), Darmon (J. M.) et Victor (Y.).		Ishra (M.). — Voir Kamal (M.)	491
- Piézorésistivité du manganin et de l'ytterbium dépo-		Kamal (M.) and Ishra (M.). — Crystallization and mechanical properties of some Fe-Ni base metallic glass	491
sés par pulvérisation cathodique	387	Khalfallah (O.) et Priester (L.). — Interaction entre les dislo-	771
Bussière (P.). — The application of Mössbauer spectroscopy	477	cations extrinsèques du joint de grains et l'impureté car-	
in heterogeneous catalysis research	477	bone dans le fer α et ses alliages cubiques centrés	153
Cadoz (J.), Castaing (J.) et Philibert (J.). — Glissement prismatique de Al ₂ O ₃ déformé par compression	135	Kripal (R.). — Voir Misra (B. N.)	411
Casenave (D.). — Voir Nodet (H.).	591	Laurence (G.), Hottier (F.) and Hallais (J.). — Qualitative and	
Castaing (J.). — Voir Cadoz (J.).	135	quantitative assessments of the growth of (Al, Ga)	
Coelho (R.) et Goffaux (R.). — Dissipation et claquage ther-	100	As-GaAs heterostructures by in situ ellipsometry	579
mique dans les solides non métalliques soumis à un		Menez (R.). — Voir Drifford (M.)	19
champ électrique intense	67	Misra (B. N.) and Kripal (R.). — Study of ESR absorption and	
Dalbiez (J. P.) Voir Drifford (M.)	19	bonding character of copper (II) complex with L-Hydro-	
Darmon (J. M.). — Voir Bosca (G.)	387	xyproline	411
David (J.). — Voir Bosca (G.)	387	Nectoux (P.) — Voir Drifford (M.)	19
Dimitriou (P.) and Gourrier (S.). — Plasma anodization of		Nodet (H.), Casenave (D.), Gauthier (R.) et Pinard (P.). — Méthode de mesure de la longueur de diffusion dans des	
silicon at room temperature	419	rubans de silicium polycristallin destinés à la conversion	
Drifford (M.), Menez (R.), Tivant (P.), Nectoux (P.) et Dal-		photovoltaïque	591
biez (J. P.). — Diffusion quasiélastique de la lumière		Pelosi (C.). — Voir Franchi (S.)	1
sous champ électrique : mobilité électrophorétique et	19	Phan Pham (T.). — Voir Gamboa (M.)	597
charge apparente des macromolécules	19	Philibert (J.). — Voir Cadoz (J.)	135
Franchi (S.), Pelosi (C.) and Attolini (G.). — Equilibrium	1	Pinard (P.). — Voir Nodet (H.)	591
calculations for VPE-InGaAsP	- '	Priester (L.) — Voir Khalfallah (O.)	153
Gamboa (M.), Phan Pham (T.), Tranduc (H.) et Rossel (P.). — Influence de la température sur la vitesse limite des		Rossel (P.) Voir Gamboa (M.)	597
trous dans un transistor MOS à canal court de type P	597	Siffert (P.). — Voir Hage-Ali (M.)	165
Gauthier (R.). — Voir Nodet (H.)	591	Tivant (P.). — Voir Drifford (M.)	19
(in) mard (P.) et Bonnet (R.) Glissements aux joints de		Tranduc (H.). — Voir Gambon (M.)	597
phases de l'eutectique lamellaire Al-Al ₂ Cu déformé à	1.45	Victor (Y.). — Voir Bosca (G.)	387
chaud par cisaillement	145	Walma (A. A.). — Statistical limits of carrier density related	77

PHYSIQUE DES DISPOSITIFS ÉLECTRONIQUES		Rosant (J. M.). — Voir Fitremann (J. M.)	93
Abou-Aly (A.). — Voir El-Raey (M.)	35	tiques	217
Azizi (C.), Graffeuil (J.) et Rossel (P.). — Le transistor à effet de champ à grille Schottky au GaAs : analyse et modèle mathématique du fonctionnement avec la grille	33	Snabre (P.). — Voir Mills (P.)	523
en polarisation directe	303		
Bielle-Daspet (D.). — Voir Roux (M.)	497	PHYSIQUE DE LA PRODUCTION	
Bourdon (B.). — Voir Squelard (S.)	657	ET DE L'UTILISATION DE L'ÉNERGIE	
Bresse (JF.). — Voir Joullie (A.).	209		
Brini (J.). — Etude expérimentale des fluctuations de courant dans une diode à double base au voisinage de son point		Alonso (N.), Beley (V. M.), Chartier (P.) and Ern (V.). — Dye	
critique	517 209	sensitization of ceramic semiconducting electrodes for photoelectrochemical conversion	5
El-Raey (M.) and Abou-Aly (A.). — On the amorphous silicon	209	Beley (V. M.). — Voir Alonso (N.)	5
on crystalline silicon heterojunctions	35	Benard (C.). — Voir Boch (G.)	237
Gamboa (M.). — Voir Rossel (P.)	509	Berger (X.). — Chauffage solaire et stratégies de chauffage.	225
Germain (P.). — Voir Squelard (S.)	657	Boch (G.), Boileau (E.) et Benard (C.). — Modélisation de	
Graffeuil (J.). — Voir Azizi (C.)	303	l'irradiation solaire au pas de temps de l'heure	237
Hamar-Thibault (S.). — Comportement d'îlots de silicium		Boileau (E.). — Voir Boch (G.)	237
épitaxié sur corindon (01.2)	317	Camanzi (A.). — Voir Romeo (N.).	11
Jalil (A.). — Voir Joullie (A.).	209	Canevari (V.). — Voir Romeo (N.) Chartier (P.). — Voir Alonso (N.)	
Joullie (A.), Rossi (A.), Bresse (JF.), Jalil (A.), Thierry-Mieg (V.) et Chevallier (J.). — Hétérojonctions ZnTe/		Cheron (Y.), Foch (H.) et Roux (J.). — Etude d'une nouvelle	7
GaAlSb préparées par épitaxie en phase liquide	209	utilisation des transistors de puissance dans les convertis-	
Phan Pham (T.). — Voir Rossel (P.).	509	seurs haute tension à fréquence élevée	333
Rossel (P.), Tranduc (H.), Gamboa (M.) et Phan Pham (T.).		Chéron (R.) et Leroy (M.). — Etude de la pression exercée	
Limitation fondamentale dans les transistors MOS de		par un champ magnétique sur un conducteur plan.	
puissance : le compromis entre la résistance à l'état pas-		Déformations produites	105
sant R_{ON} et tension de claquage V_{DBR}	509	Ern (V.). — Voir Alonso (N.)	5
Rossel (P.). — Voir Azizi (C.)	303	Foch (H.). — Voir Cheron (Y.). Joubert (A.) et Pécheux (J.). — Etude du comportement	333
Rossi (A.). — Voir Joullie (A.)	209	d'un système énergétique fonctionnant à partir du	
perties in P-type layers of silicon on sapphire	497	couplage des énergies solaire et éolienne	397
Squelard (S.), Zellama (K.), Germain (P.) and Bourdon (B.):	177	Leroy (M.). — Voir Chéron (R.)	105
Crystallization of phosphorus doped amorphous silicon		Logé (B.), Rialland (J. F.) et Robert (J.). — Etude d'un générateur électrochimique fonctionnant selon un cycle	
films prepared by glow discharge decomposition of silane	657	bitherme	37
Thierry-Mieg (V.). — Voir Joullie (A.)	209	Morand (A.) et Robert (J.). — Détermination analytique de	
Tranduc (H.). — Voir Rossel (P.)	509	la résistance électrique de barreaux rectangulaires.	
Walma (A. A.). — Statistical limits of carrier density related		Première partie : dispositifs à deux électrodes latérales	113
low frequency theories in M.O.S. devices	77	Morand (A.) et Robert (J.). — Détermination analytique de	
Zellama (K.). — Voir Squelard (S.)	657	la résistance électrique de barreaux rectangulaires.	
		2º partie : Dispositifs cylindriques infinis à trois électrodes latérales :	249
		Pécheux (J.). — Voir Joubert (A.).	
PHYSIQUE DES FLUIDES ET MÉCANIQUE	J E	Rialland (J. F.). — Voir Logé (B.).	
~		Robert (J.). — Voir Logé (B.)	
Bellet (D.). — Voir Ly (D. P.)	323	Robert (J.). — Voir Morand (A.)	
Bellet (D.). — Voir De Saint-Victor (X.).	217	Romeo (N.), Canevari (V.), Sberveglieri (G.), Tosi (A.) and	
Bitbol (M.) and Quemada (D.). — The entry flow of a dilute		Camanzi (A.). — Quasi-rheotaxy a new technique to grow	
suspension through a tube : concentration redistribution effects	663	large grain thin films on low cost amorphous substrates	11
Bousquet (A.). — Voir Ly (D. P.)	323	Roux (J.). — Voir Chéron (Y.). Sberveglieri (G.). — Voir Romeo (N.).	
Boyer (P.). — Voir Ly (D. P.)	323	Tosi (A.). — Voir Romeo (N.)	
Denat (A.), Gosse (B.) et Gosse (J. P.). — Conductivité électrique de solutions de surfactants dans les hydrocarbures.	673	200 (A)	
Fitremann (J. M.) and Rosant (J. M.). — A model for stratified	0,0		
gas-liquid turbulent flow in ducts of arbritory cross-	00	PHYSIQUE DES PARTICULES	
Section	93	DANS LE VIDE ET LES PLASMAS	
Gosse (B.). — Voir Denat (A.). Gosse (J. P.). — Voir Denat (A.).	673 673	Amount (I) Vois Bonch (D)	2.42
Quemada (O.). — Voir Bitbol (M.).	663	Amouroux (J.). — Voir Rapakoulias (D.). Berger (C.). — Voir Dard (J.)	343 435
Quemada (D.). — Voir Mills (P.)	523	Besson (J. L.). — Voir Vardelle (M.)	425
Ly (D. P.), Bellet (D.), Bousquet (A.) et Boyer (P:). — Ecou-		Chun-Sing (O.) and Schuessler (H. A.). — Storage and injec-	,2,0
lements pulsés de fluides inélastiques en conduite tron-		tion of charged particles in an electrostatic trap	441
conique ou déformable	323	Dard (J.) et Berger (C.) Expressions de potentiel électro-	
Ly (D. P.). — Voir De Saint Victor (X.)	217	statique pour des systèmes à symétrie plane et charge	
Mills (P.), Snabre (P.) and Quemada (D.). — Comportement non newtonien des suspensions de particules brownien-		d'espace nulle	435
nes	523	Domange (M.). — Voir Rapakoulias (D.). Fauchais (P.). — Voir Vardelle (M.).	343

Deistenkoin (5.), Late (1.) and vetter (R.). — Excitation		Comby (G.), Chalot (J. F.), Anger (N.), Beauval (J. J.) et	
spectrum of the iodine molecule induced by laser radia-		Quidort (J.). — Contribution à l'étude des détecteurs	
tion in the 15 780-15 815 cm ⁻¹ region : « complement		multipointes à focalisation cathodique	539
to the Atlas of the iodine molecule spectrum »	529		
Kudryavtsev (N. N.) and Novikov (S. S.), - A study of		Erratum	656
infrared radiation of vibrationally excited CO in the		Fig. 41. (NI) of Dec. 4 (NE)	
4.7 μ m band and CO ₂ in the 4.3 μ m and 2.7 μ m bands.	49	Courtier (N.) et Dumont (M.). — Asservissement de la fré-	
Luc (P.). — Voir Gerstenkorn (S.)	529	quence d'un laser à colorant sur une référence atomique	(0)
Miniscloux (D.). — Voir Taine (J.)	615	balayable par un champ magnétique	601
Novikov (S. S.). — Voir Kudryavtsev (N. N.).	49	Cuvellier (J.), Mestdagh (J. M.), Berlande (J.), De Pujo (P.) et	
Rapakoulias (D.), Domange (M.) et Amouroux (J.). — Décom-	77	Binet (A.). — Dispositif expérimental destiné à l'étude	
position des molécules polyatomiques dans un réacteur		des collisions atomiques et moléculaires aux énergies ther-	
à plasma hors d'équilibre I. NH ₃	343	miques et suprathermiques	679
Schuessler (H. A.). — Voir Chun-Sing (O.)	441	Della Negra (S.), Jacquet (D.) et Le Beyec (Y.) Analyse	
Taine (J.) et Miniscloux (D.). — Calcul d'un spectre syn-	441	de spectres d'énergie β^+ . Méthode et applications à la	
thétique d'absorption à basse résolution pour un mélange		détermination d'énergie totale de transition β pour des	
CO. CO gas pouts à boute terméseture	F15	noyaux identifiés par la technique de temps de vol dans	
CO ₂ -CO-gaz neutre à haute température	615	la région $N \simeq Z \simeq 40$	633
Vardelle (A.). — Voir Vardelle (M.)	425	Dehaes (J. C.) et Carmeliet (J.). — Installation contrôlée par	
Vardelle (M.), Vardelle (A.), Besson (J. L.) et Fauchais (P.).		microprocesseur pour la mesure des quatre paramètres	
Corrélations entre les propriétés des dépôts et les condi-		de Stokes en spectroscopie faisceau-lame	26
tions de fonctionnement d'une installation de projec-		De Pujo (P.). — Voir Cuvellier (J.)	679
tion plasma: un exemple l'alumine γ	425	Dumont (M.). — Voir Courtier (N.)	601
Vetter (R.). — Voir Gerstenkorn (S.)	529	Fournier (A. M.) et Gaune (P.). — Viscosité du chlorure de	
		sodium entre 1 100 et 1 200 K	629
		Gaune (P.). — Voir Fournier (A. M.)	629
TATOOTRIA COATOAT		Gildwarg (P.). — Voir Cahen (C.)	353
INSTRUMENTATION		Gonnard (P.). — Voir Ohanessian (H.)	647
		Jaquet (D.). — Voir Della Negra (S.)	633
Anger (N.). — Voir Comby (G.)	539	Jegou (J. P.). — Voir Cahen (C.)	353
Audoin (C.). — Fast cavity auto-tuning systems for hydrogen		Joubert (F.) Voir Nouillant (M.)	259
masers	125	Lamalle (B.), Malherbe (J. C.) et Suzeau (P.) Perfec-	
Baril (M.). — Voir Berger (C.)	175	tionnement d'un asservissement de fréquence, pour un	
Bazemlal (O.). — Voir Boquillon (J. P.)	365	spectromètre microonde utilisé dans l'étude de la largeur	
Beauval (J. J.). — Voir Comby (G.)	539	de raie de SO ₂	119
Berger (C.) et Baril (M.). — Caractéristiques optiques de		Le Beyec (Y.). — Voir Della Negra (S.)	633
miroirs électrostatiques à symétrie de révolution compo-		Malherbe (J. C.). — Voir Lamalle (B.)	119
sés de trois cylindres de même rayon	173	Mestdagh (J. M.). — Voir Cuvellier (J.)	679
Berlande (J.). — Voir Cuvellier (J.)	679	Morlais (M.). — Monochromateur de haute résolution pour	
Binet (A.). — Voir Cuvellier (J.)	679	le domaine spectral de 40-200 nanomètres	359
Blavette (D.), Bostel (A.) et Sarrau (J. M.). — Exploitation		Nouillant (M.), Tournerie (B.) et Joubert (F.). — Dispositif	207
en temps réel de la sonde atomique	405	de mesure des défauts de surfaces par émission exoélec-	
Boquillon (J. P.) et Bazemlal (O.). — Laser à colorant à		tronique photo-stimulée	259
cavité en anneau pompé par flash et de grande qualité		Ohanessian (H.) et Gonnard (P.). — Optimisation du trans-	20)
spatiale	365	fert sur capacité de l'énergie libérée par un élément piézo-	
Bostel (A.), - Voir Blavette (D.)	405	électrique en compression	647
Cahen (C.), Jegou (J. P.), Pelon (J.), Gildwarg (P.) and Porte-		Pelon (J.) Voir Cahen (C.)	353
neuve (J.). — Wavelength stabilization and control of		Porteneuve (J.). — Voir Cahen (C.).	353
the emission of pulsed dye lasers by means of a multi-		Quidort (J.). — Voir Comby (G.).	539
beam Fizeau interferometer	353	Sarrau (J. M.). — Voir Blavette (D.)	405
Carmeliet (J.). — Voir Dehaes (J. C.)	267	Suzeau (P.). — Voir Lamalle (B.)	119
Chalot (J. F.). — Voir Comby (G.).	539	Tournerie (B.). — Voir Nouillant (M.)	

The second of Market and Market a

green and the second se

e ayrı yazı elekti. At

. (1 unterest) (1

Conférence Internationale sur les Matériaux pour la Conversion Photothermique de l'Energie Solaire

Njaccio (France), 1980

optiques des surfaces rugueuses	(C.)
bouchacra (G.), Chassagne (G.) et Delmas (A.) Ma-	Battuello (M.) Voir Ricolfi (T.)
tériaux absorbants sélectifs obtenus par implantation ionique C1-327	Behaghel (J.M.) Voir Amblard (J.)
bramovich (M.) Voir Fantini (M.C.A.)	Behaghel (J.M.), Berthier (S.) et Lafait (J.) Propriétés optiques des surfaces sélec- tives : définitions et métho-
llred (D.D.) Voir Seraphin (B.O.)	des de caractérisation. C1-355
mblard (J.), Froelicher (M.), Blondeau (G.), Lafait (J.),	Behaghel (J.M.) Voir Lafait (J.) C1-133
Behaghel (J.M.) and Herrera (R.) Rough nickel selecti- ve coatings obtained by elec-	Benavente (R.) Voir Ariès (L.) C1-213
trochemical processes C1-147	Berthier (S.) Voir Behaghel (J.M.)
ndretta (A.), Bartoli (B.), Coluzzi (B.) and Cuomo (V.) Selective surfaces for natu- ral cooling devices C1-423	Berthier (S.) Voir Lafait (J.) C1-133
ranovitch (E.) Material properties and performance in photothermal solar ener- gy conversion C1-1	Berthier (S.) et Lafait (J.) Modélisation des propriétés optiques des milieux inhomo- gènes à structure com- plexe
riès (L.), Bonino (J.P.), Benavente (R.), Laaouini (A.)	Beucherie (P.) Voir Schneiders (A.M.)
et Traverse (J.P.) Ela- boration d'absorbeurs sélec- tifs de l'énergie solaire à partir d'acier inoxydable.C1-213	Bliek (P.J.), Deleuil (R.), Papini (F.) et Papini (M.) Etude et préparation de sur- faces rugueuses destinées à
rmand (L.) Voir Sigrist (M.) C1-453	la conversion photothermique de l'énergie solaire C1-105
ubert (A.) Mesure du fac- teur d'émission des surfaces sélectives pour la conver-	Blondeau (G.) Voir Amblard . (J.) C1-147
sion thermique de l'énergie solaire	Blower (R.) Voir Mason (J.J.)
alme (A.) Voir Chevalier (J.L.)	Bonino (J.P.) Voir Ariès (L.) C1-213
artoli(B.) Voir Andretta	Booth (D.C.) Voir Seraphin

Bouchut (P.) et Chenevas-Paule (A.) Influence de l'hydro- gène sur les propriétés op-	et Sacadura (J.F.) Influ- ence de l'oxydation et de la rugosité sur les caracté- ristiques radiatives des aciers inoxydables C1-16
tiques du silicium amor- phe	Driver (P.M.) and McCormick (P.G.) Photothermal degradation of black chrome - A possible contributing factor
tance	El Golli (S.) et Grenier (Ph.) Effet de serre inverse- Application à une cave so- laire
Chassagne (G.) Voir Abouchacra (G.)	Fantini (M.C.A.), Moro (J.R.) and Abramovich (M.)Thin films of gas-evaporated Co for use in photothermal conversion
Chenevas-Paule (A.) Voir Bouchut (P.) C1-439	François (J.C.) Voir Sigrist (M.)
Chevalier (J.L.), Balme (A.) et Sallee (H.) Durabili- té des matériaux plastiques transparents utilisés en couverture de capteurs solai- res plans	Froelicher (M.) Voir Amblard (J.)
Chevallier (J.) Voir Sigrist (M.)	Gravier (P.) Voir Sigrist (M.) C1-45
Coluzzi (B.) Voir Andretta (A.)	Grenier (Ph.) Voir El Golli (S.) C1-43
Craig (S.) Voir Harding (G.L.)	Harding (G.L.), Craig (S.), Curmi (P.) and Lake (M.) Selective properties of
Cuomo (V.) Voir Andretta (A.)	rough sputtered films C1-87 Harding (G.L.) and Window (B.)
Curmi (P.) Voir Harding (G.L.)	Materials problems in eva- cuated solar energy collec- tors
Cutileiro Indias (M.A.) Grandeurs optiques carac- téristiques en énergie so- laire. Uniformisation de la	Herrera (R.)Voir Amblard
terminologie C1-339 Deleuil (R.) Voir Bliek	Isetti (C.) Voir Pisoni (C.)
(P.J.)	Kayoun (P.) Voir Sigrist (M.) C1-45
Delmas (A.) Voir Abouchacra (G.)	Kulišić (P.) Voir Vuletin

aaouini (A.) Voir Ariès (L.)	Ricolfi (T.) and Battuello (M.) On the use of integrating sphere reflectometers for measuring the so-
(J.)	lar absorptance of selecti- ve surfaces
afait (J.) Voir Behaghel (J.M.)	Ricolfi (T.) Voir Pisoni (C.) C1-195
afait (J.), Behaghel (J.M.), Berthier (S.) et Rivory (J.). Surfaces sélectives rugueuses de nitrure de titane: propriétés opti- ques et modélisation C1-133	Ritchie (I.T.) and Spitz (J.) Thermal degradation of cer- met solar selective absor- bers
afait (J.) Voir Berthier (S.) C1-285	Ritchie (I.) Voir Valignat (J.)
ake (M.) Voir Harding (G.L.)	Rivory (J.) Voir Lafait (J.) C1-133
cCormick (P.G.) Voir Driver (P.M.)	Roux (L.) Voir Sigrist (M.)
ason (J.J.) and Blower (R.) Selective conversion coa-	Sacadura (J.F.) Voir Demont (Ph.)
tings on nickel and stain- less steel C1-231	Sallee (H.) Voir Chevalier (J.L.)
ason (J.J.) Voir Whittle (J.E.)	Schneiders (A.M.) et Beucherie (P.) Sélectivité spectrale par effet de structure des
attox (D.M.) and Sowell (R.R.) A survey of selective solar absorbers and their limi-	couches minces déposées de nickel et de chrome C1-123
tations	Seraphin (B.O.), Booth (D.C) and Allred (D.D.) Amorphous silicon in photothermal conversion
ailharey (D.) Voir Sigrist (M.) C1-453	Sigrist (M.), Chassaing (G.), François (J.C.), Gravier (P.), Armand (L.), Pierrisnard
apini (F.) Voir Bliek (P.J.)	(R.), Roux (L.), Pailharey (D.), Kayoun (P.) et Chevallier (J.) Propriétés optiques
apini (M.) Voir Bliek (P.J.)	des nitrures et carbonitrures de titane - Sélectivité spectrale C1-453
etit (R.) Sur l'étude élec- tromagnétique de l'absorp- tion de la lumière par les surfaces rugueuses C1-43	Sikkens (M.) Spectrally selective absorption in nickel carbide and nickel nitride films C1-465
ierrisnard (R.) Voir Sigrist (M.)	Sowell (R.R.) Voir Mattox (D.M.)
isoni (C.), Isetti (C.), Battuello (M.) and Ricolfi (T.) Procedures for ob- taining black chrome solar	Spitz (J.) Voir Ritchie (I.T.)
selective coatings with re- producible radiation proper-	Tran Nguyen (H.) Voir Demont (Ph.)

Tran Van (D.) Voir Valignat (J.)	Whittle (J.E.) and Mason (J.J.) The use of porta- ble instruments for opti- cal property measurement of selective surfaces C1-36
Valignat (J.), Tran Van (D.) et Ritchie (I.) Caracté- risation et stabilité de	Window (B.) Voir Harding (G.L.) C1-17
couches antireflets C1-393	Wirgin (A.) Sur la réponse spectrale de certaines
Vuletin (J.), Kulišić (P.) and Bosanac (M.) Selecti- velv absorbing coatings C1-191	surfaces texturées C1-57

Les Implications Conceptuelles de la Physique Quantique

Paris (France), 1980

les inégalités de Bell	C2-63
Bell (J.S.) Bertlmann's socks and the nature of reality	C2-41
D'Espagnat (B.) A la recherche du réel	C2-99
Laloë (F.) Cadre général de la mé- canique quantique. Les objections de Einstein, Podolsky et Rosen	C2-1
Shimony (A.) Reflexions sur la phi- losophie de Bohr, Heisenberg et Schrödinger	C2-81

and the second second in the Se

idastera, intitist, na sa sa santa

y (E.) i stayen om Spase de li cy hy i en gy

The Interactions between Dislocations and Defects in Oxides

Lyon (France), 1980

Blanchin (M.G.), Bursill (L.A.), Hutchison (J.L.) and Gai (P.L.) Direct observation of disorder within crystallographic shear defects in defor- med and non-stoichiome- tric rutiles	C3-95	Fontaine (G.). See Blanchin (M.G.)	C3-1
Blanchin (M.G.), Castaing (J.), Fontaine (G.), Heuer (A.H.), Hobbs (L.W.) and Mitchell (T.E.) Introductory report: " The interactions bet- ween dislocations and de- fects in oxides"	C3-1	tion	C3-31 C3-95 cd C3-113
Blanchin (M.G.) See Vri- nat (M.)	C3-79	(W.)	C3-119
Bursill (L.A.) See Blan-	C3-95	(M.G.)	C3-1
chin (M.G.)Cadoz (J.) See Castaing		Heuer (A.H.) See Keller (R.J.)	C3-73
Castaing (J.), Cadoz (J.) and Kirby (S.H.) Deformation of Al ₂ O ₃ single	C3-43	Hirsch (P.B.) Plastic deformation and electronic mechanisms in semiconductors and insulators	
crystals between 25°C and 1800°C: basal and prismatic slip	C3-43	Hobbs (L.W.) See Blanchin (M.G.)	
Castaing (J.) See Blanchin (M.G.)	C3-1	Hutchison (J.L.) See Blanchin (M.G.)	C3-95
Castaing (J.) See Dominguez- Rodriguez (A.)	- C3-67	Keller (R.J.), Mitchell (T.F and Heuer (A.H.) Slip systems in UO _{2+x}	
Dominguez-Rodriguez (A.), Sanchez (M.), Marquez (R.), Castaing (J.) and Monty	,	Kirby (S.H.) See Castaing (J.)	
(C.) Point defects and high temperature creep of CoO single crystals	C3-67	Marquez (R.) See Dominguez Rodriguez (A.)	C3-67
Duclos (R.) High temperature deformation of MgO·1.8Al ₂ O ₂	9	Mitchell (T.E.) See Blanch (M.G.)	
spinel single crystals of <111> axial orientation.	C3-49	Mitchell (T.E.) See Keller (R.J.)	C3-73
Fantozzi (G.) See Gabbay (M.)	C3-31	Monty (C.) See Dominguez- Rodriguez (A.)	C3-67

slip planes.....

Puls (M.P.) Atomistic models of point-defect/dislocation interactions in oxides C3-13	Thevenard (P.) and Guermazi (M.) Extended defects and precipitates in implan- ted TiO2
Routbort (J.L.) High-tempe- rature deformation of cubic	Vrinat (M.) and Blanchin
oxides	(M.G.) Plastic deforma- tion of slightly reduced
Sanchez (M.) See Dominguez- Rodriguez (A.) C3-67	rutile
Skrotzki (W.) and Haasen (P.) Hardening mechanisms of ionic crystals on {110} and {100}	

.... C3-119

COLLOQUE Nº 4 - 1981 (Tome I et Tome II)

Ninth International Conference on Amorphous and Liquid Semiconductors

Grenoble (France), 1981

Abdulgafarov (E.E.) See Sarsembinov (Sh.Sh.)	C4-911	Andriesh (A.M.), Gerasimenko (V.S. Ivaschenko (Yu.N.), Iovu (M.A.	·),
Abelès (B.) See Cody (G.D.)	C4-301	Iovu (M.S.), Mironos (A.V.), Smirnov (V.L.), Chernii (M.R.) and Shutov (S.D.) Effect of	
Abelès (B.) See Wronski (C.R.)	C4-475	composition and structure on transport properties of chalco-	
Abkowitz (M.) and Enck (R.C.) Xerographic spectroscopy of		genide glasses	C4-963
localized electronic states in amorphous semiconductors	C4-443	Andro (P.) See Peyre (J.F.)	C4-163
Adler (D.) Defects in amorphous		Aoki (H.) Anderson localisation in two-band systems	C4-51
chalcogenides and silicon	C4-3	Assumpção (R.) See Chambouley-	0/ 1000
Adler (D.) See Senturia (S.D.)	C4~503	ron (I.)	C4-1009
Adler (D.) See Shiraishi (T.).	C4-579	Atari (S.) See Sawan (Y.)	C4-511
Adler (D.) See Walsh (P.J.)	C4-317	Austin (I.G.) See Depinna (S.P.)	C4-323
Adriaenssens (G.J.), Michiel (H.) and Hammam (M.). Isothermal relaxation currents in arsenic chalcogenides	C4-939	Austin (I.G.) See Nashashibi (T.S.)	C4-467
Agarwal (S.C.) See Kumar (S.).	C4-527	Azoury (S.J.) See Senturia (S.D.)	C4-503
Akagi (Y.) See Kawamura (H.)	C4-357	Baixeras (J.) See Peyre (J.F.)	C4-163
Al Dallal (S.)See Kalem (S.)	C4-361	Balberg (I.), Hanak (J.J.), Weakli (H.A.) and Gal (E.) Fundament	
Aleksiejúnas (A.), Bareikienė (R.) Bondarenko (V.) and Gečiauskas	,	tunneling processes in MOS _a solar cells	
(S.) Conductivity relaxation in structure Pt-liquid V ₂ 0 ₅ -Pt	C4-1059	Balkan (N.) See Long (A.R.)	C4-107
Alimoussa (L.), Carchano (H.) and Thomas (J.P.) Influence of		Bali (K.) See Szörényi (T.)	C4-997
hydrogen partial pressure on deposition and properties of		Bandow (T.) See Suzuki (M.)	C4-623
sputtered amorphous gallium arsenide	C4-683	Bareikienė (R.) See Aleksiejunas (A.)	
Allison (J.) See Mirza (A.R.).	C4-659	Bässler (H.) See Schönherr (G.)	C4-111
Alvarez (F.) See Dalal (V.)	C4-491	Belin (E.) See Dufour (G.)	C4-877
Amer (N.M.) See Jackson (W.B.)	C4-293	Belu (A.) See Devenyi (A.)	C4-1089
Andrew (R.) See Wautelet (M.).	C4-395	Bendali (M.) See Mackowski (J.M.)	C4-975

Berger (J.M.) See Donnadieu (A.)	C4-655	(W.J.)	C4-193
Bernholc (J.) See DiVincenzo (D.P.)	C4-137	Booth (D.C.) and Voss (K.J.) The optical and structural properties of CVD germanium	
Beyer (W.), Mell (H.) and Overhof (H.) Transport properties of		300000000000000000000000000000000000000	04-1033
compensated a-Si films	C4-103	Bordure (G.) See Giuntini (J.C.)	04-95
Beyer (W.) and Wagner (H.) Hy- drogen evolution from plasma- deposited amorphous silicon		Bösch (M.A.) See Murayama (K.)	C4-343
films	C4-783	Bouchut (P.) See Olivier (M.).	C4-305
Bhatia (K.L.) and Hunklinger (S.) Interaction of ultra- sonic phonons with donor impuri	i –	Boulitrop (F.) See Dunstan (D.J.)	C4-331
ties in hydrogenated amorphous	C4-479	Bourdon (B.) See Zellama (K.).	C4-815
Bhatia (K.L.), v. Haumeder (M.) and Hunklinger (S.)- Study of defects in a-Ge and a-GeH _x		Bourne (L.C.), Rowland (S.C.) and Bienenstock (A.).— Radical dis- tribution function studies of glassy germanium-silver-chalcoge	en
using surface acoustic wave technique	C4-365	-	C4-951
Bienenstock (A.) See Bourne		Bourneix (J.) See Kalem (S.)	C4-361
(L.C.)	C4-951 ·	Brassard (C.), Groleau (R.), L'Ecuyer (J.), Martin (J.P.), Currie (J.F.), Depelsenaire (P.), Wertheimer (M.) and Yelon (A.). Nuclear scattering measurements of composition profi-	
talline counterparts	C4-383	les in a-Si:H multilayer structures	C4-795
Bishop (S.G.) See Shanabrook (B.V.)	C4-865	Breitschwerdt (K.G.) See Schmid	
Blake (J.) See Viktorovitch (P.)	C4-455	Brenzikofer (R.) See Chambouleyro	on C4-1009
Blake (J.) See Oguz (S.)	C4-679	Bresser (W.J.), Boolchand (P.), Suranyi (P.) and De Neufville	
Blanchard (C.) See Deville (A.)	C4-1001	(J.P.) Intrinsically broken chalcogen chemical order in	C4-193
Bogomolova (L.D.), Jachkin (V.A.), Glassova (M.P.) and Spasibkina (S.N.) Vanadate semiconducting glasses doped with oxides of other transition metals		Brinkmann (R.), Elwenspoek (M.), von Hartrott (M.), Novak (A.), Quitmann (D.) Nuclear quadru- polar relaxation in liquid	
Bonch-Bruevich (V.L.) and Mironov			C4-105
(A.G.) Some problems of the theory of electron energy spec-		Brodsky (M.H.). See Hvam (J.M.).	C4-551
trum of a disordered semiconductor	C4-33	Brodsky (M.H.) See DiVincenzo (D.P.)	C4-137
Bondarenko (V.) See Aleksiejunas	3	Brooks (B.) See Cody (G.D.)	C4-301

	C4-849	Chakraverty (B.K.) Imperfect amorphous solid and bipola- ronic ground-state	C4-749
Bruyère (J.C.) See Deneuville (A.)	C4-733	Chamberlain (J.M.) and Moseley	
Bruyère (J.C.) and Deneuville (A.) Photoluminescence ver- sus boron doping and prepara-		(A.J.) First observations of photoconductivity in bulk doped Ge _Y Se _{1-Y} glasses	C4-309
tion conditions	C4-339	Chambouleyron (I.), Constantino	
Bruyère (J.C.) See Toulemonde (M.)	C4-799	(C.), Jousse (D.), Assumpção (R.) and Brenzikofer (R.) Electrical and optical proper-	
Buroff (A.) Arsenic sulphide as		ties of amorphous tin oxide	C4-1009
evaporated dry photoresist	C4-967	Chao (S.S.) See Tsu (R.)	C4-269
Butcher (P.N.) and McInnes (J.A.) The Hall mobility of		Chao (Y.Q.) See Liao (X.B.)	C4-663
hopping carriers	C4-91	Chen (I.) and Lee (S.) Theoretical analyses of a-Si:H dio-	
Caffier (D.), Le Contellec (M.)		de characteristics	C4-499
and Richard (J.). Photocon- ductivity in a-Si:H and		Chen (Z.M.) See Liao (X.B.)	C4-663
a-Si _x C _{1-x} :H, correla- tion with photoluminescence		Chenevas-Paule (A.) and Dijon	
results	C4-1037	(J.) Thermal and optical	
	0/ 1/7	space charge spectroscopy of gap states in a-Si:H	C4-605
Cagna (A.) See Pistoulet (B.).	C4-147	Set coaces in a primitivity	04 003
Capezzuto (P.) See Evangelisti (F.)	C4-849	Chenevas-Paule (A.).— See Lamotte (B.)	C4-839
Carchano (H.) See Alimoussa		Chernii (M.R.) See Andriesh	
(L.)	C4-683	(A.M.)	C4-963
Cardona (M.) See Shen (S.C.)	C4-349	Chevallier (J.) See Kalem (S.)	C4-361
Carius (R.), Fischer (R.) and Holzenkämpfer (E.). The band		Chianelli (R.R.) See Chien (F.Z.)	C4-273
gap of glow-discharge-produced amorphous SiO _X	C4-1025	Chien (F.Z.), Moss (S.C.), Liang (K.S.) and Chianelli (R.R.).	
Carlos (W.E.) and Taylor (P.C.)		Computer calculation of scatte-	
High temperature H NMR in a-Si:H	C4-725	ring intensity for disordered molydenum disulfide	C4-273
		Chik (K.P.) See John (P.K.)	C4-639
Cavenett (B.C.).— See Depinna (S.P.)	C4-323	Chopra (K.L.) See Rasagopalan	
C (C D) Observe (V) and		(S.)	C4-911
Ceasar (G.P.), Okumura (K.) and Grimshaw (S.F.) Preparation		Cody (G.D.), Tiedje (T.), Abelès	
and properties of ion beam deposited a-SiH _X	C4-627	(B.), Moustakas (T.D.), Brooks (B.) and Goldstein (Y.) Disor der and the optical absorption	-
Cebulka (J.M.) See Tiedje (T.)	C4-155	edge of hydrogenated amorphous	C4-301
Cetina (E.A.) See Sansores			
(L.E.)	C4-823	Cohen (J.D.), Lang (D.V.), Harbiso	n

(J.P.) and Sergent (A.M.) Stu-	Davis (E.A.) See Fasol (G.) C4-5/1
dy of light induced changes in	Delegative (C) and Tourse (D)
a-Si:H by detailed computer mode- ling of admittance and DLTS C4-371	Deleonibus (S.) and Jousse (D.) Electron transport in hydroge-
Collins (R.W.) and Paul (W.) Time	nated amorphous silicon. Schottky barriers and deep localized sta-
resolved photoluminescence spectra	tes kinetics C4-487
of sputtered a-Si:H C4-591	tes kinetics
	Demond (F.J.), Müller (G.), Damjant-
Collins (R.W.) See Oguz (S.) C4-679	schitsch (H.), Mannsperger (H.), Kalbitzer (D.), Lecomber (P.G.)
Constantino (C.) See Chambouleyron	and Spear (W.E.) Hydrogen pro-
(I.)	filing in gas phase doped and
Cramarossa (F.) See Evangelisti	ion implanted amorphous silicon
(F.)	films C4-779
(11)	Den Boer (W.) Determination of
Crandall (R.S.) Electronic pro-	midgap density of states in
perties of a-Si:H from measure-	a-Si:H using space-charge-
ments on devices C4-413	limited current measurements. C4-451
Cros (A.), Salvan (F.) and Derrien	De Neufville (J.P.) See Bresser
(J.) AES and EELS measurements	(W.J.)
on the Si-Au-Ag interface C4-1081	
Currie (I E) - See Present (C) C/-705	Deneuville (A.), Bruyère (J.C.)
Currie (J.F.) See Brassard (C.) C4-795	and Hamdi (H.) Multiphases description of a-Si:H C4-733
Cutler (M.) Bond equilibrium	description of a-Si:H C4-733
theory in covalently bonded	Deneuville (A.) See Bruyère
alloys	(J.C.)
•	
Cutler (M.) See Radscheit (H.) C4-1051	Deneuville (A.) See Toulemonde
	(M.)
Czaja (W.) and Kinmond (S.)	Deneuville (A.) See Magarino
Photoluminescence decay in	(J.)
a-Si: influence of excitation density and n-type doping C4-543	
density and n-type doping 04-545	Depelsenaire (P.) See Brassard
Czubatyj(W.) See Madam (A.) C4-463	(C.)
(101) 101	Depinna (S.P.), Cavenett (B.C),
Dalal (V.) and Alvarez (F.)	Searle (T.M.) and Austin (I.G.)
Minority carrier transport	Optically detected magnetic re-
in depletion layer of n-i-p	sonance (ODMR) in a-Si C4-323
Si:H solar cells C4-491	
	Derrien (J.) See Cros (A.) C4-108
Damjantschitsch (H.) See Demond	
(F.J.)	Dersch (H.) See Schweitzer (L.) C4-827
Dandoloff (P) Zowhow (P)	Day 1 (4) D (6) D 1 (4)
Dandoloff (R.), Zeyher (R.) The third low of thermodyna-	Devenyi (A.), Rusu (C.), Belu (A.)
mics and the low temperature	and Manaila (R.) Mo-modified amorphous Ge : electrical trans-
properties of disordered	port and structure C4-108
solids C4-205	
	Deville (A.), Gaillard (B.),
Datta (T.) and Silver (M.) Tran-	Blanchard (C.) and Livage (J.)
sient photoconductivity and	Electron spin-lattice relaxa-
Schottky barrier profile deter-	tion: a tool for the investi-
mination in a-Si:H C4-563	gation of paramagnetic centers
Davis (E.A.) Review of group-V	in amorphous semiconductors. C4-1001
amorphous semiconductors C4-855	Dias (A.G.) See Guimaraes (L.) C4-609

Dijon (J.) See Chenevas-Paule (A.) C4-605	Elwenspoek (M.) See Brinkmann (R.)	C4-1055
Dimaria (D.J.) Insulator physics and engineering: Electrically- Alterable Read-Only-Memory applications	Emin (D.) Temporal decay of the optically induced properties of a small-polaronic solid	C4-535
DiVincenzo (D.P.), Bernholc (J.)	Enck (R.C.) See Abkowitz (M.).	C4-443
and Brodsky (M.H.) The electronic structure of a model defect in hydrogenated amorphous silicon C4-137	Evangelisti (F.), Fiorini (P.), Fortunato (G.), Frova (A.), Bruno (G.), Capezzuto (P.) and Cramarossa (F.) Photo- luminescence studies on	
Dixmier (J.), Mosseri (R.) and Sadoc (J.F.) Diffraction study of	a-Si:H:Cl films	C4-849
the a-Si:H structure evolution during annealing treatments	Ewen (P.J.) See Firth (A.P.)	C4-903
performed directly on the X ray goniometer C4-237	Failly-Lovato (M.) See Wautelet (M.)	C4-395
Dixmier (J.) See Wu Zhong Yan. C4-515	Fang (C.J.) See Vardeny (Z.)	C4-539
Döhler (G.H.) Theory of tempe- rature and intensity dependence of photoconductivity in amor- phous semiconductors C4-115	Fasol (G.) and Davis (E.A.) Ti- me-resolved luminescence mea- surements on amorphous phos- phorus	C4-571
Donnadieu (A.), Yous (B.), Berger (J.M.), Ferraton (J.P.) and Robin (J.) Effect of temperature on optical properties of glow discharge hydrogenated	Feltz (A.), Kahnt (H.) and Schirrmeister (F.). Evidence for charged defects in chalco- genides from high field conduc- tivity	C4-935
amorphous silicon films C4-655	Ferrari (L.) and Russo (G.) On	
Dufour (G.), Belin (E.), Senemaud (C.), Gheorghiu (A.) and Theye (M.L.) Disorder effects on GaP core levels studied by elec-	the relaxation time of certain two-level systems in amorphous structures	C4-1073
tron spectroscopies C4-877	Ferraton (J.P.) See Donnadieu (A.)	C4-655
Dunstan (D.J.) and Boulitrop (F.) A no-Stokes shift model for the	Ferrier (R.P.) See Long (A.R.)	C4-107
photoluminescence of a-Si:H C4-331 Dupin (J.P.) See Gheorghiu (A.)	Fiorini (P.) See Evangelisti (F.)	C4-849
Easton (B.C.) See Powell	Firth (A.P.), Owen (A.E.) and Ewen (P.J.) Laser-Raman stu- dy of the structure of silver	
(doped As ₄₀ S ₆₀ glass	C4-903
Economou (E.N.) See Pickett (W.E.)	Fischer (R.) See Carius (R.)	C4-1025
El-Den (M.B.) See Kotkata (M.F.)	Fischer (R.) See Radscheit (H.)	C4-1051
El-Gabaly (M.) See Sawan (Y.). C4-511	Fischer (R.) See Hoheisel (B.)	C4-819
Elliott (S.R.) Photo-induced	Fitzgerald (A.G.) See Spear (W.E.)	C4-257
E.S.R. in glassy sulphur C4-387		

Flachet (J.C.) - See Thomas		Gaspard (J.P.). See Lemaire	
(P.)	C4-151	(P.)	C4-765
Fleury (G.), Viger (C.), Vautier (C.) and Lhermitte (C.) Crystallization of amorphous Se	-	Gaspard (J.P.) See Mosseri (R.)	C4-245
films	C4-947	Geciauskas (S.). See Aleksiejunas	c4-1059
Flynn (B.W.) and Owen (A.E.) The electronic and optical proper-		Geiger (J.) See Kniffler (N.).	C4-811
ties of vanadium tellurite gla	C4-1005	Gelmont (B.L.), Kolomiets (B.T.), Mazets (T.F.) and Pavlov (S.K.)	
Fortuna (H.S.) See Scharp (A.C.)	C4-159	Polarized light electroabsorp- tion anisotropy of vitreous	
Fortunato (G.) See Evangelisti		chalcogenide semiconductors	C4-919
(F.)	C4-849	Gerasimenko (V·S·).— See Andriesh (A·M·)	C4-963
Fournier (Y.) See Nguyen Van Dong	C4-647	Germain (P.) See Zellama (K.).	C4-815
Friederich (A.) See Magarino (J.)	C4- 737	Gheorghiu (A.), Rappeneau (T.), Dupin (J.P.) and Theye (M.L.) Electronic properties of flash- evaporated amorphous GaSb	
The Hall effect due to hopping conduction in the localized		films	C4-881
states of amorphous semi- conductors	C4-87	Gheorghiu (A.) See Dufour (G.)	C4-877
Fritzsche (H.), Smid (V.), Ugur (H.) and Gaczi (P.J.) Plasma		Gibbs (G.V.) See Revesz (A.G.)	C4-217
deposited hydrogenated chalcogonide glasses		Gibson (R.A.) See Spear (W.E.)	C4-1143
Fritzsche (H.) See Tanielian	0/ 275	Gibson (R.A.) - See Nashashibi (T.S.)	C4-467
(M.H.)	C4-375	Giuntini (J.C.), Jacquemin (J.L.),	
Fritzsche (H.) See Persans (P.D.)	C4-597	Zanchetta (J.V.) and Bordure (G.) Modelling of the behavior of hopping conductivity in a so	
Frova (A.) See Evangelisti (F.)	C4-849	lid exhibiting a semiconductor- insulator transition, as a func- tion of frequency	
Fuhs (W.) See Welsch (H.M.)	C4-567		
Fuhs (W.) See Grünewald (M.)	C4-523	Glassova (M.P.). See Bogomolova (L.D.)	C4-993
Furukawa (S.) See Matsumura (H.)	C4-209	Godzik (K.) and Schirmacher (W.) Theory of dispersive transport in amorphous semiconductors	- C4-127
Füstöss-Wégner (M.) See Hajtó (J.)	C4-313	Gogna (P.K.) See John (P.K.)	C4-639
Gaczi (P.J.) See Fritzsche	C4-699	Goldstein (Y.) See Cody (G.D.)	C4-301
Gaillard (B.) See Deville (A.)	C4-1001	Goldstein (Y.) See Wronski (C.R.)	C4-475
Gal (E.) See Balberg (I.)	C4-459	Goodmann (N.B.) See Tanielian	
		(M.H.)	T Thomas of I be

Graudenz (W.) - See Mackinnon		Hamakawa (Y.) Recent progress	
(A.)	C4-63	of the amorphous silicon solar	
		cells and their technology	C4-1131
Greaves (G.N.) Sodium environ-			
ments in glass	C4-225	Hamakawa (Y.) See Tawada (Y.).	C4-471
Greeb (K.H.) See Welsch (H.M.)	C4-567	Hamakawa (Y.) See Hotta (S.)	C4-631
Grevendonk (W.) See Vanhuyse	C4-923	Hamakawa (Y.) See Okamoto (H.)	C4-507
		Hamakawa (Y.) See Nonomura	
Grigorovici (R.) and Vancu (A.) Structural modelling of rever-		(S.)	C4-761
sible photodarkening in		Hamanaka (H.), Tanaka (K.), Tsuji	
a-As _x Se	C4-391	(K.) and Minomura (S.). Detailed temperature-cycling studies	
0.1.1. (0.7.)		of photostructural change in	,
Grimshaw (S.F.) See Ceasar		a-As ₂ S ₃	C4-399
(G.P.)	C4-627	4 1223	04-355
Grob (J.J.) See Toulemonde		Hamasaki (T.), Hirose (M.) and Osaka (Y.) Plasma diagnostic	
(M.)	C4-799	during the growth of a-Si:H	C4-807
		during the growth of a 51.11.	C4-007
Groleau (R.) See Brassard	C4-795	Hamdi (H.) See Deneuville (A.)	C4-733
Grünewald (M.), Müller (H.),		Hamdi (H.) See Toulemonde (M.)	C4-799
Thomas (P.) and Wirtz (D.)		Hammam (M.) See Adriaenssens	
Temperature dependence of the		(G.J.)	C4-939
hopping Hall mobility in spa-		() () () () () () () () () ()	
tially and energetically disordered systems	C4-99	Hanak (J.J.) See Balberg (I.).	C4-459
Grünewald (M.), Weber (K.), Fuhs		Hänisch (N.) See Kniffler (N.)	C4-811
(W.) and Thomas (P.). Field		Harbison (J.P.) See Cohen	
effect studies on a-Si:H films	C4-523	(J.D.)	C4-371
		Harshavardhan (K.S.) See	
Grünewald (M.) See Schweitzer	0/ 007	Rajagopalan (S.)	C4-911
(L.)	C4-827		
Guha (S.) See Navkhandewala		Hasegawa (S.), Kurata (Y.), Imai	
(R.V.)	C4-803	(Y.) and Narikawa (S.) - RF	
d		power and temperature depen- dences in GD a-Si produced	
Guimaraes (L.), Martins (R.) and			C4-675
Dias (A.G.) Photoresponse		from heated SiH,	C4-073
time and density of localized states in the mobility gap of		Hata (N.) See Yamasaki (S.)	C4-297
a-Si:H alloys prepared by R.F. glow discharge	C4-609	Hattori (K.) See Kawamura (H.)	C4-357
		The Manual on (M.) Can Manual	
Hajtó (J.) and Füstöss-Wégner		v. Haumeder (M.) See Bhatia	C1-265
(M.) Laser induced oscilla-		(K.L.)	C4-365
tory phenomena in a-GeSe ₂		Thursday (T. T.) - Americkeus south	
films	C4-313	Hauser (J.J.) Amorphous semi-	C4-943
		conducting Ag-Te films	U4 743
Halpern (V.) A comparison of		V. Idana Ha and Haiana-Na Ida	
coherent potential and random		Yu-Liang He and Hsiang-Na Liu	
walk methods for calculating		Investigation into the conduc-	
the transport properties of		tion mechanism of CVD amorphous	
amorphous semiconductors	C4-119	and polycrystalline silicon	C4-831
		111111111111111111111111111111111111111	07 001

Hevesi (I.) See Szörényi (T.). C4-	Recombination of photogenera-	
Hirabayashi (I.), Morigaki (K.) and Nitta (S.).— Recombination processes and gap states in	ted carriers in doped hydroge- nated amorphous silicon	C4-551
hydrogenated amorphous silicon as elucidated by time-resolved	Ichimura (T.) See Uchida (Y.).	C4-265
	-587 Imai (Y.) See Hasegawa (S.)	C4-675
Hirabayashi (I.) See Morigaki (K.) C4-	Imgrund (H.) and Overhof (H.) 335 Dispersive transport in R- hopping systems	C4-83
Hiraki (A.), Imura (T.), Mogi (K.) and Tashiro (M.) Si-H vibra-		C4-277
tional properties in crystalli- zed hydrogenated silicon fabri-	Inoue (M.) See Shirafuji (J.).	C4-583
cated by reactive sputtering in H ₂ atmosphere		C4-1123
Hirata (K.) See Kitao (M.) C4-	-927 Inuishi (Y.) See Shirafuji	
Hirose (M.) Physical properties	(J.)	C4-583
of amorphous CVD silicon C4-	(A.M.)	C4-963
Hirose (M.) See Hamasaki (T.) C4-	-807 Iovu (M.S.) See Andriesh	
Hirose (M.) See Mishima (Y.) C4-	(A.M.)	C4-963
Hodges (C.H.) A simple inter- pretation of electron loca-	Iqbal (Z.) See Veprek (S.)	C4-251
	Ivaschenko (Yu.N.) See Andriesh (A.M.)	C4-963
Hogg (W.R.) See Long (A.R.) C4-	-107 Izu (M.) See Tsu (R.)	C4-269
Hoheisel ((B.), Fischer (R.) and Stuke (J.) Dak- and photocon-	Jaafar (A.) See Walsh (P.J.)	C4-317
ductivity of compensated a-Si:H C4-	Jachkin (V.A.) See Bogomolova (L.D.)	C4-993
(20)	Jackson (W.B.) and Amer (M.N.) Subgap and band edge optical absorption in a-Si:H by photo-	
Horigome (S.) See Taniguchi (Y.)	-971 thermal deflection spectros-	C4-293
	Jacquemin (J.L.) See Giuntini	C4-95
Hotta (S.), Tawada (Y.), Okamoto (H.) and Hamakawa (Y.).— effects of the substrate po—	Jan (G.J.) See Tsu (R.)	C4-269
tential on the incorporation manner of hydrogen and impu-	Janai (M.) See Weil (R.)	C4-643
rity in a-Si:H films C4-	Janai (M.) Photolithographic processes in amorphous semi-	
Hsiang-Na Liu See Yu-Liang He. C4-	-831 conductors	C4-1105
Hunklinger (S.) See Bhatia (K.L.)	-479 Jeffrey (F.R.) See Shanks	C4-773
Hunklinger (S.) See Bhatia	Ye Xian Jing See Peng Shao-Qi.	C4-791

John (P.), Odeh (I.M.), Thomas (M.J.K.) and Wilson (J.I.B.)Preparation and properties of		Kawamori (A.). See Kawamura (H.)	C4-357
polysilanes : model compounds for hydrogenated a-Si	C4-651	Kawamura (H.), Hattori (K.), Matsunaga (K.), Akagi (Y.) and Kawamori (A.) Order parame-	
John (P.K.), Wong (S.K.), Gogna (P.K.), Tong (B.Y.) and Chik (K.P.). Theta-pinch plasma		ter in the glass transition of vitreous S-Ge	C4-357
hydrogenation of evaporated amorphous silicon films	C4-639	Kawamura (T.) See Yamamoto	C4-495
Johnson (R.L.) See Ley (L.)	C4-753	Keiper (R.) See Van Der Meer	06 175
Johnson (N.M.) See Thompson		(M.)	C4-175
(M.J.)	C4-617	Kelemen (S.) See Wronski	C4-475
Jones (D.I.) and Stewart (A.D.)		(0.000)	04 47 5
Electrical and optical properties of hydrogenated amorphous		Kim (G.I.) See Shirafuji (J.).	C4-583
carbon	C4-1085	Kinmond (S.) See Czaja (W.)	C4-543
Jousse (D.) See Deleonibus	oi 107	Kitao (M.), Hirata (K.) and	
(S.)	C4-487	Yamada (S.) AC conductivity	0/ 007
Jousse (D.) See Chambouleyron		of amorphous As-Se-Ag system.	C4-927
(I.)	C4-1009	Kniffler (N.), Muller (W.W.),	
Kahnt (H.) See Feltz (A.)	C4-935	Pirrung (J.M.), Hänisch (N.), Schröder (B.) and Geiger (J.).	
Kakimoto (Y.) See Suzuki (M.).	C4-623	Annealing and hydrogenation behaviour of evaporated and	
Kalbitzer (S.) See Reinelt		sputtered high-purity amor- phous silicon films	C4-811
(M.)	C4-843	photo dalacon lalmovitiviti	01 011
		Kolomiets (B.T.) Studies of	
Kalbitzer (D.) See Demond	01 330	chalocogenide vitreous semi-	
(F.J.)	C4-779	conductors in the Ioffe physico-technical institute	C4-887
Kalbitzer (S.) See Spear	0/ 11/2	V-1	
(W.E.)	C4-1143	Kolomiets (B.T.).— See Gelmont (B.L.)	C4-919
Kalem (S.), Chevallier (J.), Al Dallal (S.) and Bourneix (J.).	-	Kondo (H.) See Nakamura (G.)	C4-483
Infrared vibrational spectra of chlorinated and hydrogena-	C4-361	Kondo (M.) See Tawada (Y.)	C4-471
ted amorphous silicon	C4-737	Kong (G.L.) See Liao (X.B.)	C4-663
Kaplan (D.) See Magarino (J.).	U4=737	Kotkata (M.F.) and El-Den (M.B.).	
Kärcher (R.) See Ley (L.)	C4-753	Hardness and thermal conductivity of As-Se-Te chalcoge-	
Kastner (M.) See Monroe (D.)	C4-559	nide glasses	C4-931
Katayama (Y.), Shimada (T.), Usami (K.) and Maruyama (E.)		Kramer (B.) See Mackinnon (A.)	C4-63
Chemical sites of rare gas atoms in amorphous silicon		Kramer (B.) See Leschke (H.)	C4-59
and the same of th		Kumar (S.) and Agarwal (S.C.)	
Kawakyu (Y.) See Matsumura		Photovoltaic effect in inter-	
(H ₂)	C4-209	stitially doped a-Si:H	C4-527

Kumeda (M.) See Veda (S.) Kumurdjian (P.) See Mackowski	C4-729	Lemaire (P.) and Gaspard (J.P.) Density of states of amorphous hydrogenated Si	C4-765
(J.M.)	C4-975	Le Ny (J.Y.) See Nguyen Van	04 703
Kuno (S.) See Matsumura (M.)	C4-519	Dong	C4-647
Kurata (Y.) See Hasegawa (S.).	C4-675	Leschke (H.) and Kramer (B.) Disorder induced density of	
Kuwano (Y.), Ohnishi (M.) In- dustrialization of a-Si solar		states within the band gap	C4-59
cells	C4-1155	Levin (K.) See Weil (R.)	C4-643
Lachter (A.) See Oguz (S.)	C4-679	Ley (L.), Richter (H.), Kärcher (R.), Johnson (R.L.), Reichardt	
Lambert (C.J.) See Weaire (D.)	C4-47	(J.) Surface properties of a-Si:H and a-Si:F investigated	
Lamotte (B.), Rousseau (A.), Chenevas-Paule (A.). High re- solution NMR of ²⁹ Si in amor-		by photoelectron spectrosco-	C4-753
phous hydrogenated silicon	C4-839	Ley (L.) See Richter (H.)	C4-261
Lang (D.V.) See Cohen (J.D.)	C4-371	Lhermitte (C.) See Fleury (G.).	C4-947
Lannin (J.S.) See Pomian (R.J.)	C4-873	Liang (K.S.) See Chien (F.Z.).	C4-273
Laude (L.D.) See Wautelet (M.)	C4-395	Liao (X.B.), Kong (G.L.), Yang (X.R.), Wang (P.D.), Chao	
Lauder (E.H.) See Sussmann (R.S.)	C4-1029	(Y.Q.), Chen (Z.M.) and Liu (C.L.). Electrical properties and photoluminescence of amor-	
Le Comber (P.G.) Snell (A.J.), Mackenzie (K.D.), Spear		phous silicon	C4-663
(W.E.) Applications of a-Si field effect transistors		Liu (C.L.) See Liao (X.B.)	C4-663
in liquid crystal displays and in integrated logic cir-		Livage (J.) Amorphous transition metal oxides	
cuits	C4-423	Livage (J.) See Deville (A.)	C4-1001
Le Comber (P.G.). See Demond (F.J.)	C4-779	Long (A.R.), Hogg (W.R.), Balkan	
Le Comber (P.G.) See Nashashibi		(N.) and Ferrier (R.P.) AC loss in amorphous germanium at	
(T.S.)	C4-467	low temperatures	C4-107
Le Comber (P.G.) See Spear (W.E.)	C4-257	Longeaud (C.) See Peyre (J.F.)	C4-163
Le Comber (P.G.) See Spear		Lowry (M.E.) See Shanks (H.R.)	C4-773
(W.E.)	C4-1143	Lucovsky (G.) Chemical bending of alloy and dopant atoms in	
Le Contellec (M.). See Caffier (D.)	C4-1037	amorphous silicon	C4-741
L'Ecuyer (J.) See Brassard (C.)	C4-795	Lucovsky (G.) See Pollard (W.B.)	C4-353
De Lin Lee See Wen Yuan Xu	C4-695	McGill (J.) See Madan (A.)	C4-463
Lee (S.) See Chen (I.)	C4-499	McInnes (J.A.) See Butcher (P.N.)	C4-91

Manhanda (V.D.) O. t. C.			
Mackenzie (K.D.). See Le Comber (P.G.)	C4-423	Mašek (J.), Velický (B.) Elec- tronic state in amorphous ger-	
Mackinnon (A.), Kramer (B.) and		manium chalcogenides	C4-133
Graudenz (W.) New numerical method for the electronic properties of disordered systems	C4-63	Matsuda (A.), Matsumura (M.), Nakagawa (K.), Yamasaki (S.) and Tanaka (K.) A simpli-	
Mackintosh (A.J.) See Phillips		fied model for the deposi- tion kinetics of GD a-Si:H	0/ (07
(R.T.)	C4-869	films	C4-687
Mackowski (J.M.), Bendali (M.),		Matsuda (A.) See Yamasaki (S.)	C4-297
Normandon (P.) and Kumurdjian (P.) Diffusion kinetics in gold-amorphous GeTe, thin		Matsui (H.) See Shirafuji (J.)	C4-583
films	C4-975	Matsumura (H.), Sakai (K.), Kawakyu (Y.) and Furukawa	
Madan (A.), Czubatyj (W.), Yang (J.), McGill (J.) and Ovshinsky (S.R.) High efficiency, large		(S.) Structural model of fluorinated amorphous-silicon (a-Si:F)	C4-209
area photovoltaic devices using amorphous Si:F:H alloy		Matsumura (M.), Kuno (S.) and Uchida (Y.) Current path in	
Maekawa (T.) See Suzuki (M.)	C4-623	amorphous-silicon field effect	C4-519
Magarino (J.), Friederich (A.),			
Kaplan (D.) and Deneuville (A.) Effect of hydrogena-		Matsumura (M.) and Uchida (Y.) Large-area deposition of amor- phous-silicon	C4-671
tion on doped a-Si prepared by CVD	C4-737	Matsumura (M.) See Matsuda (A.)	C4-687
Manaila (R.) Reflection SAXS investigation of amorphous		Matsunaga (K.) See Kawamura (H.)	C4-357
thin films	C4-213	Mazets (T.F.) See Gelmont	
Manaila (R.) See Devenyi (A.).	C4-1089	(B.L.)	C4-919
Mannsperger (H.) See Demond (F.J.)	C4-779	Meaudre (M.), Meaudre (R.), Tardy (J.) and tribollet (B.).— Correlation between conductivity,	:-
Marfaing (Y.) See Wu Zhong Yan	C4-515	electron spin resonance and optical absorption in RF sput-tered SiO ₂ films	C4-1013
Marshall (J.M.) See Scharp (A.C.)	C4-159	Meaudre (R.) See Meaudre (M.).	C4-1013
Martin (J.P.) See Brassard		Mell (H.) See Beyer (W.)	C4-103
(C.)	C4-795	Mell (H.) See Weller (D.)	C4-143
Martins (R.) See Guimaraes (L.)	C4-609	Mell (H.) See Welsch (H.M.)	C4-567
Martinez (E.) and Yndurain (F) Electronic structure of amor-		Mencaraglia (D.) See Peyre (J.F.)	C4-163
phous SiO _x	C4-1021	Michiel (H.) See Adriaenssens (G.J.)	C4-939
Maruyama (E.) See Katayama	C/-797		
(Ÿ.)	C4-787	Minomura (S.) Pressure-induced transitions in amorphous sili-	
Maruyama (E.) See Taniguchi (Y.)	C4-971	con and germanium	C4-181

Minomura (S.) See Tsuji (K.) C4-233	Mosseri (R.) See Dixmier (J.). C4-237
Minomura (S.) See Hamanaka (H.) C4-399	Mosseri (R.) See Sadoc (J.F.). C4-189
Minomura (S.) See Oyanagi (H.) C4-221	Mott (N.F.) Conduction near a mobility edge C4-27
Mironos' (A.V.) See Andriesh	
(A.M.)	Moustakas (T.D.) See Tiedje (T.) C4-155
Mironov (A.G.) See Bonch-Bruevich	
(V.L.)	Moustakas (T.D.) See Cody (G.D.)
Mirza (A.R.), Rhodes (A.J.),	Marcahar (P) - The theory of
Allison (J.) and Thompson (M.J.) R.F. magnetron sputte-	Movaghar (B.) The theory of
ring of a-Si-H C4-659	transport in amorphous semi- conductors
Ting of a bi necession of opposite	conductors
Mishima (Y.), Hirose (M.), Suemune (I.), Yamanishi (M.) and Osaka	Müller (G.) See Demond (F.J.). C4-779
(Y.) Observation of acoustic emission from a-Si:H pin junc-	Müller (G.) See Spear (W.E.) C4-114
tions C4-447	Müller (H.) See Grünewald (M.) C4-99
Moddel (G.) See Viktorovitch	Müller (W.W.) See Kniffler
(P.)	(N.)
Mogi (K.) See Hiraki (A.) C4-277	Murayama (K.) and Bösch (M.A.) Radiative recombination of
Monroe (D.), Orenstein (J.) and	localized excitons in amorphous
Kastner (M.) Density of states	and crystalline As ₂ S ₃ C4-343
in the gap of a-As ₂ Se ₃ by pho-	N1- (D.) D1 (M.)
tocurrent transient spectros-	Nagels (P.), Rotti (M.) and
. сору С4-559	Vikhrov (S.) Doping of chal-
Morigaki (K.), Hirabayashi (I.),	cogenide glasses in the Ge-Se and Ge-Te systems C4-907
Sano (Y.) and Nitta (S.).	and Ge-Te systems C4-907
Defect creation by optical	Naito (H.) See Okuda (M.) C4-601
exitation in hydrogenated	(11)
amorphous silicon : time-	Nakagawa (K.) See Matsuda (A.) C4-687
resolved luminescence and	
ODMR measurements C4-335	Nakamura (G.), Sato (K.), Kondo
	(H.), Yukimoto (Y.) and Shirahata
Morigaki (K.) See Hirabayashi	(K.) Tandem type solar cells
(I.)	using a-Si:H and a-SiGe:H
Mort (J.) Geminate and non-	films C4-483
geminate recombination in	Nakau (T.) See Okuda (M.) C4-601
amorphous semiconductors C4-433	(10)
Moseley (A.J.) See Chamberlain	Nakayama (H.) See Okuda (M.) C4-601
(J.M.)	Nakayama (Y.) See Yamamoto (N.) C4-495
	2000 2000 (11) (4.43)
Moser (F.) See Weil (R.) C4-643	Narasimhan (K.L.) See Navkhandewala (R.V.)
Moss (S.C.) See Chien (F.Z.) C4-273	
Manager (D) and Coopered (ID)	Narikawa (S.) See Hasegawa
Mosseri (R.) and Gaspard (J.P.) Growth sequence of Si-clusters:	(S.)
from a few atoms to the amor-	Nashashibi (T.S.), Austin (I.G.),
phous phase C4-245	Searle (T.M.), Gibson (R.A.),

Le Comber (P.G.) and Spear		Odeh (I.M.) See John (P.)	C4-651
(W.E.) Injection luminescenc	е	(20),,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,	01 031
in amorphous silicon p [†] -i-n [†] junctions	C4-467	Oguz (S.), Paul (D.K.), Blake (J.), Collins (R.W.), Lachter	
Navkhandewala (R.V.), Narasimhan		(A.), Yacobi (B.G.) and Paul (W.) Inferior electronic	
(K.L.) and Guha (S.) Electro irradiation in amorphous hydro		properties of rf-sputtered a-Si:H films with only the	
genated silicon		2000-cm-1 Ir absorption	0/ 670
Nemanich (R.J.) See Tsai	0/ 1077		C4-679
(C.C.)	C4-1077	Oguz (S.) See Viktorovitch	0/ /55
Ngai (K.L.) Temperature and		(P.)	C4-455
frequency dependences of NMR spin-lattice relaxation time		Oheda (H.) See Okushi (H.)	C4-613
T ₁ in a-Si:H	C4-835	Oheda (T.) See Yamasaki (S.)	C4-297
Ngai (K.L.) See Strom (U.)	C4-123	Ohnishi (M.) See Kuwano (Y.)	C4-1155
Nguyen Van Dong, Fournier (Y.) an Le Ny (J.Y.). A new type of	d	Ohsawa (M.) See Uchida (Y.)	C4-265
a-Si prepared by dc sputte-		Okamoto (H.), Yamaguchi (T.),	
ring : P, B and H doping		Nonomura (S.) and Hamakawa	
effect	C4-647	(Y.) Drift type photovol- taic effect in a-Si p-i-n	
Nicholls (D.H.) See Powell		junction	C4-507
(M.J.)	C4-379		
Nichino (T.) See Nonomura (S.)	C4-761	Okamoto (H.) See Nonomura (S.)	C4-761
Nitta (S.), Takahashi (Y.) and		Okamoto (H.) See Hotta (S.)	C4-631
Noda (M.) Existence of irreversibility in Staebler-		Okamoto (H.) See Tawada (Y.)	C4-471
Wronski effect and fatigue		Okano (S.), Suzuki (M.) and Suzuk: (M.) Contact effects of me-	L
effect of photoluminescence of GDa-Si:H	C4-403	tals on chalcogenide amorphous	
	04 403	films	C4-959
Nitta (S.) See Hirabayashi	C4-587	Okuda (M.), Naito (H.), Nakayama (H.) and Nakau (T.) Optical	
Nitta (S.) See Morigaki (K.)	C4-335	DLTS measurements of localized	1-
	01 100	states in amorphous chalcogenic	C4-601
Noda (M.) See Nitta (S.)	C4-403		
Nonomura (S.), Okamoto (H.), Nishino (T.) and Hamakawa		Okumura (K.) See Ceasar (G.P.)	C4-62/
(Y.) Electroreflectance study of electronic struc-		Okushi (H.), Tokumaru (Y.), Yama- saki (S.), Oheda (H.) and	
trure in a-Si:H	C4-761	Tanaka (K.) Isothermal capa-	
	-1 -0-	citance transient spectroscopy	•
Nonomura (S.) See Okamoto (H.)	C4-507	its application to the study of gap states of a-Si:H	C4-613
Normandon (P.) See Mackowski	-1		C/-207
(J.M.)	C4-971	, Okushi (H.) See Yamasaki (S.).	C4-297
Novak (A.) See Brinkmann (R.).	C4-1055	Olivier (M.) and Bouchut (P.) Enhancement of the photocurrent	
Oda (S.) See Shimizu (I.)	C4-1123	signal in a-Si:H by mean of a waveguiding technique. Applica-	
Odajima (A.) - See Heno (T.)	C4-899	tion to the optical absorption	

spectroscopy below the band	C4-305	Perron (J.C.) See Rabit (J.)	C4-1047
Orenstein (J.) See Monroe (D.)	C4-559	Persans (P.D.) and Fritzsche (H.) Dual light beam modu-	
Osaka (Y.) See Mishima (Y.)	C4-447	lation of photocarrier life- time in intrinsic a-Si:H	C4-597
Osaka (Y.) See Hamasaki (T.)	C4-807	Peyre (J.F.), Baixeras (J.),	
Oswald (H.R.) See Veprek (S.).	C4-251	Mencaraglia (D.), Andro (P.) and Longeaud (C.).— Electron	
Overhof (H.) See Imgrund (H.).	C4-83	drift mobility measurement in undoped triode DC sput- tered a-Si:H	C4-163
Overhof (H.) See Beyer (W.)	C4-103	Phillips (J.C.) Domain micros-	04-103
Ovshinsky (S.R.) Principles and applications of amorphicity, structural change, and optical		copy in chalcogenide alloy glass thin films	C4-197
information encoding	C4-1095	Phillips (R.T.), Mackintosh (A.J.) and Yoffe (A.D.)	
Ovshinsky (S.R.). See Madan (A.)	C4-463	Electrical and optical pro- perties of UHV sublimed a-As.	C4-869
Ovshinsky (S.R.) See Tsu (R.).	C4-269	Phillips (W.A.) See Wright (0.B.)	C4-1017
Owen (A.E.) See Firth (A.P.)	C4-903	Picard (C.) See Zellama (K.)	C4-815
Owen (A.E.) See Flynn (B.W.)	C4-1005	Pichard (J.L.) and Sarma (G.).	
Oyanagi (H.), Tanaka (K.), Hosoya (S.) and Minomura (S.). EXAFS studies on the local structure		Finite size scaling approach to Anderson localization	C4-37
in amorphous GeS and GeSe	C4-221	Pickett (W.E.), Papaconstantopoulo (D.A.), Economou (E.N.) Cal-	S
Pankove (J.I.) See Schade (H.)	C4-327	culations of transport proper- ties in a-Si:H	C4-769
Papaconstantopoulos (D.A.) See Pickett (W.E.)	C4-769	Pilione (L.J.) See Pomian (R.J.)	C4-873
Paul (W.) Closing Address	C4-1165		04-073
Paul (D.K.) See Oguz (S.)	C4-679	Pirrung (J.M.) See Kniffler (N.)	C4-811
Paul (W.) See Collins (R.W.)	C4-591	Pistoulet (B.), Roche (F.) and Cagna (A.) AC field and	
Paul (W.) See Oguz (S.)	C4-679	frequency dependence of a-Si:H conductivity at 4.2 K	C4-147
Paul (W.) See Viktorovitch (P.)	C4-455	Plecenik (R.M.) See Scott	
	C4-433	(B.A.)	C4-635
Pavlov (S.K.) See Gelmont (B.L.)	C4-919	Pollak (M.) See Friedman (L.).	C4-87
Peng Shao-Qi, Yu Bing Qai, Zhang Pei Xian and Ye Xian Jing		Pollak (F.H.) See Tsu (R.)	C4-269
Effects of Argon on the pro- perties of RF sputtered amor-	0/ 701	Pollak (M.) A scaling theory for hopping conduction	C4-141
Pepper (M.) Two dimensional	C4-791	Pollard (W.B.) and Lucovsky (G.) Phonon densities of states and	-
electron localization	C4-17	eigenvectors in hydrogenated	

	and fluorinated amorphous si-	C4-353	Rebouillat (J.P.) See Vérié	C4-667
or.	mian (R.J.), Pilione (L.J.) and Lannin (J.S.) Optical absorp-		Reichardt (J.) See Ley (L.)	c4-753
Por	tion structural order in sputte red amorphous phosphorus		Reimer (J.A.) NMR investiga- tions of hydrogenated amor- phous silicon	C4-715
	ling of the liquid state of tetrahedrally bended semi-conductors	C4-1043	Reinelt (M.) and Kalbitzer (S.) The diffusion of heavy alkali atoms in amorphous silicon	C4-843
	rtal (J.C.) See Ždanowicz	C4-1069	Revesz (A.G.), Wemple (S.H.), Gibbs (G.V.) Structural or- dering related to chemical	
701	well (M.J.), Easton (B.C.) and Nicholls (D.H.) The effect of		bonds in random networks	C4-217
	annealing and illumination on the field effect conductance of		Rhodes (A.J.) See Mirza (A.R.)	C4-659
D-10	amorphous silicon	C4-379	Richard (J.) See Caffier (D.).	C4-1037
	Bing Qai See Peng Shao-Qi	C4-643	Richter (H.) and Ley (L.) Pho- toemission studies of the tran- sition from amorphous to mi-	
Qu:	itmann (D.) See Brinkmann		crocrystalline silicon	C4-261
	(R.)	C4-1055	Richter (H.) See Ley (L.)	C4-753
Ra	bit (J.) and Perron (J.C.) Photoconductivity in liquid selenium	C4-1047	Robertson (J.) Theory of dopant level depths in a-Si	C4-745
Ra	dscheit (H.), Fischer (R.) and		Robin (J.) See Donnadieu (A.).	C4-655
	Cutler (M.) Electronic beha- vior of liquid semiconducting		Roche (F.) See Pistoulet (B.)	C4-147
	alloys My(Se.5 Te.5)1-y with monovalent doping ele-		Rochette (J.F.) See Vérié (C.)	C4-667
	ments	C4-1051	Rotti (M.) See Nagels (P.)	C4-907
Ra	dscheit (H.) Nonmetal to me- tal transition in liquid chal-		Rousseau (A.) See Lamotte (B.)	C4-839
	cogenides	C4-1063	Rowland (S.C.). See Bourne (L.C.)	C4-951
ĸa	<pre>jagopalan (S.), Harshavardhan (K.S.), Singh (B.) and Chopra (K.L.) Photo and electroche- mical doping in Ge-based chal-</pre>		Rubinstein (J.) See Senturia (S.D.)	C4-503
	cogenides	C4-911	Russo (G.) See Ferrari (L.)	C4-1073
Ra	ppeneau (T.) See Gheorghiu	C4-881	Rusu (C.) See Devenyi (A.)	C4-1089
Ra	tobylskaja (V.A.) and Simonova (L.A.) Ultrasonic studies of the polymer structure of the chalcogenide glasses	C4-955	Sadoc (J.F.), Mosseri (R.) A new approach of amorphous semiconductors structure using curved spaces	C4-189
-			Sadoc (J.F.) See Dixmier (J.).	C4-237
Ka	<pre>(y (S.), Vardeny (Z.) and Tauc (J.) Time resolved spectros- copy of the photoinduced ab-</pre>		Saito (K.) See Shimizu (I.)	C4-1123
	sorption in glow-discharge	C4-555	Saito (S.) See Taniguchi (Y.).	C4-971

Sakai (K.) See Matsumura (H.).	C4-209	Schröder (B.) See Kniffler	C4-811
Salvan (F.) See Cros (A.)	C4-1081	Schuchardt (R.) See Van Der Meer	
Sano (Y.) See Morigaki (K.)	C4-325	(M.)	
Sansores (L.E.), Cetina (E.A.) and Tagueña-Martínez (J.). Application of molecular orbital self-consistent-field method to the study of a-Si	d C4-823	Schweitzer (L.), Grünewald (M.) and Dersch (H.).— Influence of correlation effects on the electronic properties of amor- phous silicon	C4-827
Sarma (G.) See Pichard (J.L.).	C4-37	Schweitzer (L.)- See Weller (D.)	C4-143
Sarott (F.A.) See Veprek (S.). Sarsembinov (Sh. Sh.), Abdulgafare (E.E.) and Tolkanchinov (K.)	C4-251	Scott (B.A.), Plecenik (R.M.) and Simonyi (E.E.). Deposition of a-Si:H by homogeneous CVD	C4-635
Radiation-induced defects formation in chalcogenide glasses	C4-915	Searle (T.M.) See Depinna (S.P.)	C4-323
grasses	04 713	Searle (T.M.) See Nashashibi	
Sato (K.). See Nakamura (G.)	C4-483	(T.S.)	C4-467
Sawan (Y.), El-Gabaly (M.), Wakim (F.) and Atari (S.) A study		Senemaud (C.) See Dufour (G.).	C4-877
of a new heterojunction made of n-type silicon and p-type amorphous semiconductor	C4-511	Senturia (S.D.), Rubinstein (J.), Azoury (S.J.) and Adler (D.) Charge centroid determination in field-effect experiments	C4-503
Schade (H.) and Pankove (J.I.) Electron-beam induced centers in hydrogenated amorphous si- licon	C4-327	Sergent (A.M.) - See Cohen (J.D.)	
Scharp (A.C.), Marshall (J.M.) and Fortuna (H.S.) Transport properties of amorphous		Shah (J.) Time resolved optical spectroscopy in amorphous silicon and chalcogenides	-
As ₂ Se ₃	C4-159	Shanabrook (B.V.), Bishop (S.G.) and Taylor (P.C.). Photolumi-	
Scher (H.) Recombination in disordered solids	C4-547	nescence and ESR studies of lo- calized states in amorphous	
Shiraishi (T.) and Adler (D.)		phosphorus	C4-865
High-speed transient effects in chalcogenide glasses	C4-579	Shanabrook (B.V.) See Bishop (S.G.)	C4-383
Schirmacher (W.) See Godzik (K.)	C4-127	Shanks (H.R.), Jeffrey (F.R.), Lowry (M.E.). Bonding in hy- drogenated amorphous silicon.	C4-773
Schirrmeister (F.) See Feltz		drogenated amorphous silicon.	04-773
(A·)···································	C4-935	Shen (S.C.) and Cardona (M.) Infrared and far infrared ab-	
Schmidt (W.W.) and Breitschwerdt (K.G.) AC transport and structural relaxations in		sorption of B- and P-doped amorphous Si	C4-349
disordered solids	C4-171	Shimada (T.) See Katayama (Y.)	C4-787
Schönherr (G.), Bässler (H.) and Silver (M.) Monte Carlo simu- lation of variable range hop-		Shimakawa (K.) Contribution of single polaron hopping to AC conduction in amorphous chal-	
ping at the Fermi levels	C4-111	cogenides	C4-167

Shimizu (I.), Oda (S.), Saito (K.	,	of amorphous edition asi-	
Tomita (H.) and Inoue (E.) a-Si:H in electrophotography) ;	of amorphous silicon solar cells	C4-1143
and vidicon devices	C4-1123	Spear (W.E.), Willeke (G.), Le Comber (P.G.) and Fitzgerald	
Shimizu (T.) See Ueda (S.)	C4-729	(A.G.) Electronic proper- ties of microcrystalline si-	
Shirashi (T.) and Adler (D.) High-speed transient effects in chalcogenide glasses	C4-579	licon films prepared in a glow discharge plasma	C4-257
Shirafuji (J.), Kim (G.I.), Matsu:	Ĺ	Spear (W.E.) See Demond (F.J.)	C4-779
(H.), Inoue (M.), Yoshino (K.) and Inuishi (Y.) Non-dispers and dispersive transport in	ive	Spear (W.E.) See Le Comber (P.G.)	C4-423
amorphous germanium selenide and hydrogenated silicon	C4-583	Spear (W.E.).— See Nashashibi (T.S.)	C4-467
Shirahata (K.) See Nakamura (G.)	C4-483	Stewart (A.D.) See Jones (D.I.)	C4-1085
Shutov (S.D.) See Andriesh (A.M.)	C4-963	Stoica (T.) Light-induced changes in a-Si:H analysed by field	
Siffert (P.) See Toulemonde		effect measurements	C4-407
(M.)	C4-799	Street (R.A.) Photoconductivity and related measurements of	
Silver (M.) See Datta (T.)	C4-563	the conduction band tail in a-Si:H	C4-575
Silver (M.) See Schönherr (G.)	C4-111	Street (R.A.) Luminescence in	
Simonova (L.A.) See Ratobylskaja (V.A.)	C4-955	a-Si:H	C4-283
Simonyi (E.E.) See Scott	C4-635	Street (R.A.) See Thompson (M.J.)	C4-617
Singh (B.) See Rajagopalan		Strom (U.) and Ngai (K.L.) AC conductivity of disordered	
(S.)	C4-911	solids below 10 ¹¹ Hz	C4-123
Slade (M.L.) See Zallen (R.)	C4-241	Strom (U.) See Bishop (S.G.)	C4-383
Slechta (J.) On efficiency of scattering methods to determi-		Stuke (J.) See Hoheisel (B.)	C4-819
ne the structure of disordered materials	C4-229	Stuke (J.) See Weller (D.)	C4-143
Śmid (V.) See Fritzsche (H.)	C4-699	Suemune (I.) See Mishima (Y.).	C4-447
Smirnov (V.L.) See Andriesh		Zhong Lin Sun See Wen Yuan Xu	C4-695
(A.M.)	C4-963	Suranyi (P.) See Bresser (W.J.)	C4-193
Snell (A.J.) See Le Comber (P.G.)	C4-423	Sussmann (R.S.) and Lauder (E.H.). Luminescence in plasma deposi-	-
Spasibkina (S.N.) - See Bogomolova (L.D.)	a C4-993	ted amorphous Si _x C _{l-x}	C4-1029
Spear (W.E.), Gibson (R.A.), Yang (D.), Le Comber (P.G.), Muller (G.), Kalbitzer (S.) Some new developments in the field		Suzuki (M.), Maekawa (T.), Kakimot (Y.) and Bandow (T.) Proper- ties of pure silicon amorphous films prepared by rf-BIAS sput-	

Suzuki (M.) See Okano (S.)	C4-959	Theye (M.L.) See Dufour (G.)	C4-877
Suzuki (M.) See Okano (S.)	C4-959	Theye (M.L.).— See Gheorghiu (A.)	C4-881
Szörényi (T.), Bali (K.) and Hever (I.) Amorphous vanadium oxides by CVD: preparation, elec- trical magnetic properties		Thomas (P.) and Flachet (J.C.) Conductivity measurements on UHV deposited amorphous sili-	
Tagüena-Martinez (J.) See		con	C4-151
Sansores (L.E.)	C4-823	Thomas (P.) See Grünewald (M.)	C4-99
Takahashi (Y.) See Nitta (S.).	C4-403	Thomas (P.) See Grünewald (M.)	C4-523
Tanaka (K.) See Hamanaka (H.).	C4-399	Thomas (J.P.) See Alimoussa (L.)	C4-683
Tanaka (K.) See Matsuda (A.)	C4-687	Thomas (M.J.K.). See John (P.).	C4-651
Tanaka (K.) See Okushi (H.)	C4-613		
Tanaka (K.) See Oyanagi (H.)	C4-221	Thompson (M.J.), Johnson (N.M.) and Street (R.A.).— Measure— ment and analysis of current	
Tanaka (K.) See Yamasaki (S.).	C4-297	transients in well-characte- rized a-Si:H	C4-617
Tanielian (M.H.), Goodmann (N.B.)			
and Fritzsche (H.) Photo- creation of defects in plasma- deposited a-Si:H	C4-375	Thompson (M.J.).— See Mirza (A.R.)	C4-659
Taniguchi (Y.), Yamamoto (H.),		Thompson (M.J.) See Tsai (C.C.)	C4-1077
Horigome (S.), Saito (S.) and Maruyama (E.). A new chalco- genide photoconductor for a diode laser beam printer	~ ~71	Thompson (M.J.) See Walsh (P.J.)	C4-317
,	C4-971	Tiedje (T.), Moustakas (T.D.) and	
Tardy (J.) See Meaudre (M.)	C4-1013	Cebulka (J.M.) Temperature dependence of the electron	
Tashiro (M.) See Hiraki (A.)	C4-277	drift mobility in hydrogenated a-Si prepared by sputtering	C4-155
Tashiro (M.) See Hiraki (A.)	C4-277	Tiedje (T.) See Cody (G.D.)	C4-301
Tauc (J.) See Ray (S.)	C4-555	Tokumaru (Y.) See Okushi (H.).	C4-613
Tauc (J.) See Vardeny (Z.)	C4-539		
Tawada (Y.), Kondo (M.), Okamoto (H.) and Hamakawa (Y.) Valen	_	Tokunaga (A.) See Tsujino (K.) Tolkanchinov (K.) See Sarsembino	
cy electron control in a glow discharge produced a-SiC:H and		(Sh. Sh.)	C4-915
its application to a-Si solar cell	C4-471	Tomita (H.) See Shimizu (I.)	C4-1123
Tawada (Y.) See Hotta (S.)	C4-631	Tong (B.Y.) See John (P.K.)	C4-639
Taylor (P.C.) See Bishop (S.G.)	C4-383	Toulemonde (M.), Grob (J.J.), Bruyère (J.C.), Deneuville (A. Hamdi (H.) and Siffert (P.)),
Taylor (P.C.) See Carlos		Interaction between argon and dopants in sputtered a-Si:H	C4-799
(W-E.)	C4-725		0. 777
Taylor (P.C.) See Shanabrook		Tribollet (B.) See Meaudre	C4-1013
(B.V.)	C4-865		

Tsai (C.C.), Thompson (M.J.) and Nemanich (R.J.).— Structural and electrical properties of noble metal-hydrogenated amor- phous silicon interfaces	C4-1077	Vanhuyse (B.), Van Den Keybus (P.) and Grevendonk (W.) Faraday rotation in a-As ₂ Se ₃	C4-923
Tsu (R.), Chao (S.S.), Izu (M.), Ovshinsky (S.R.), Jan (G.J.)	04 1077	Vardeny (Z.), Tauc (J.) and Fang (C.J.). Picosecond dynamics of carriers in amorphous semi-	
and Pollak (F.H.) The nature of intermediate range order in		conductors	C4-539
Si:F:H:(P) alloy systems	C4-269	Vardeny (Z.). See Ray (S.)	C4-555
Tsuji (K.), Minomura (S.) X-ray diffraction study of atomic		Vautier (C.) See Fleury (G.) Velický (B.) See Mašek (J.)	C4-947
structure of plasma deposited a-Si:H alloys	C4-233	Veprek (S.), Iqbal (Z.), Oswald	04-133
Tsuji (K.) See Hamanaka (H.)	C4-399	(H.R.), Sarott (F.A.) and Wagner (J.J.) Parameters	
Tsujino (K.), Yamamoto (M.), Tokunaga (A.) and Yonezawa (F.) Localization of elec-		controlling the deposition of amorphous and microcystal- line silicon in Si:H discharge	
trons in two-dimensional		plasmas	C4-251
random networks	C4-55	Vérié (C.), Rochette (J.F.) and Rebouillat (J.P.) New amor-	
Uchida (Y.), Ichimura (T.), Ueno (M.) and Ohsawa (M.). Proper- ties of microcrystalline P-dopo	ed	phous alloy semiconductors : a-Si _{1-x} Sn _x	C4-667
glow discharge Si:H films	C4-265	Viger (C.) See Fleury (G.)	C4-947
Uchida (Y.) See Matsumura (M.)	C4-519	Vikhrov (S.) See Nagels (P.)	C4-907
Uchida (Y.) See Matsumura (M.)	C4-671	Viktorovitch (P.), Moddel (G.),	
<pre>Ueda (S.), Kumeda (M.) and Shimize (T.) NMR and ESR studies on annealing effects in a-Si:F:H</pre>	u	Blake (J.), Oguz (S.), Weisfield (R.L.) and Paul (W.) Density of states stu-	
and a-Si:H	C4-729	dy in sputtered a-Si:H effect of impurities and H related	
Ueno (T.) and Odajima (A.) Silver-doped amorphous As ₂ Se ₃		defects	C4-455
films studied by XPS	C4-899	Viscor (P.), Yoffe (A.D.) Sta- ble amorphous germanium films	
Ueno (M.) See Uchida (Y.)	C4-265	prepared in ultra high vacuum and measured in-situ structure	C4-691
Ugur (H.) See Fritzsche (H.)	C4-699	and electronic properties	04-091
Usami (K.) See Katayama (Y.)	C4-787	Von Hartrott (M.) See Brinkmann (R.)	C4-1055
Vancu (A.) See Grigorovici (R.)	C4-391	Voss (K.J.) See Booth (D.C.)	C4-1033
Van Den Keybus (P.). See Vanhuyse	C4-923	Wagner (H.) See Beyer (W.)	C4-783
		Wagner (J.J.) See Veprek (S.).	C4-251
Van Der Meer (M.), Keiper (R.) and Schuchardt (R.) Non- ohmic hopping conduction-a		Wakim (F.) See Sawan (Y.)	C4-511
treatment by directed perco-	C4-175	Wakita (K.) See Yamamoto (N.).	C4-495

Walsh (P.J.), Jaafar (A.), Thompso	n	Witzke (H.) See Wronski (C.R.)	24-475
(M.J.) and Adler (D.) Active and passive studies of amor-		Wong (S.K.) See John (P.K.)	24-639
phous chalcogenide IR emit- ters	C4-317	Wright (0.B.) and Phillips (W.A.) Thermoelastic effects in vi-	
Wang (P.D.) See Liao (X.B.)	C4-663	treous silica	24-1017
Zhong Pang Wang See Wen Yuan	C4-695	Wronski (C.R.), Goldstein (Y.), Kelemen (S.), Abeles (B.) and Witzke (H.) Correlation of	
Wautelet (M.), Andrew (R.), Failly-Lovato (M.) and Laude (L.D.) Laser-induced meta-		electrical properties of a-SiH _X MIS solar cell struc- tures with the state of oxida- tion of the interface	C4-475
stable states in amorphous semiconductors	C4-395		24-99
Weaire (D.) and Lambert (C.J.) The decimation method and Andersen localization	C4-47	Zhang pei Xian See Peng Shao-	C4-791
Weakliem (H.A.) See Balberg	C4-459	Wen Yuan Xu, Zhong Lin Sun, Zhong Pang Wang and De Lin Lee The enhancement of the dark conduc-	
Weber (K.) See Grünewald (M.).	C4-523	tivity and photoconductivity of GD a-Si introducing O_2 , N_2	C4-695
Weclewicz (Cz.) See Zdanowicz	C4-1069		C4-679
Weil (R.), Janai (M.), Pratt (B.),		Yamada (S.) See Kitao (M.)	C4-927
Levin (K.) and Moser (F.) Glow-discharge a-Si:F prepared from SiF ₂ gas	C4-643	Yamaguchi (T.) See Okamoto (H.) (Yamamoto (H.) See Taniguchi	C4-507
Weinstein (B.A.) See Zallen (R.)	C4-241	(Y.)	C4-971 C4-55
Weisfield (R.L.) See Viktorovito			J4-33
(P.)		Yamamoto (N.), Wakita (K.), Nakayama (Y.) and Kawamura	
Weller (D.), Mell (H.), Schweitzer (L.) and Stuke (J.) Magnetic field dependence of the photo-		(T.) Photoelectric proper- ties of GD a-Si :H monolayer films for electrophotographic	
and dark conductivity in do- ped a-Si:H films	C4-143	applications	C4-495
Welsch (H.M.), Fuhs (W.), Greeb	0, 2,0	Yamanishi (M.) See Mishima (Y.)	C4-447
(K.H.) and Mell (H.) Prima- ry and secondary photocurrents		Yamasaki (S.), Hata (N.), Yoshida	
in n-type and p-type a-Si:H films	C4-567	(T.), Oheda (H.), Matsuda (A.), Okushi (H.) and Tanaka (K.) Annealing studies on low opti-	
Wemple (S.H.) See Revesz (A.G.)	C4-217	cal absorption of GD a-Si:H using photoacoustic spectros- copy	C4-297
Wertheimer (M.) See Brassard (C.)	C4-795		24-687
Willeke (G.) See Spear (W.E.).	C4-257	Yamasaki (S.) See Okushi (H.).	24-613
Wilson (J.I.B.) See John (P.).	C4-651	Wu Zhong Yan, Marfaing (Y.) and	

in transport properties in a-Si p-n- junctions	C4-515	Yukimoto (Y.) See Nakamura	C4-483
Yang (D.) See Spear (W.E.)	C4-1143	Zallen (R.), Weinstein (B.A.) and Slade (M.L.). Network dimension) —
Yang (J.) See Madan (A.)	C4-463	nality of amorphous GeS ₂ : op- tical high-pressure experiments	
Yang (X.R.) See Liao (X.B.)	C4-663	on a-GeS ₂ , 2d-GeS ₂ , and	C4-241
Yelon (A.) See Brassard (C.)	C4-795	Zanchetta (J.V.) See Giuntini	0/ 05
Ynduráin (F.) See Martínez		(J.C.)	C4-95
(E.)	C4-1021	Zdanowicz (L.), Żdanowicz (W.), Weclewicz (Cz.), Portal	
Yoffe (A.D.) See Phillips (R.T.)	C4-869	(J.C.) Transport properties of amorphous films of cadmium	
Yoffe (A.D.) See Viscor (P.)	0/ (01	arsenide	C4-1069
ioffe (A.D.) See viscor (P.)	C4-691	Ždanowicz (W.) See Ždanowicz	
Yonezawa (F.) See Tsujino		(L.)	C4-1069
(K.)	C4-55	Zellama (K.), Germain (P.), Picard (C.) and Bourdon (B.)	
Yoshida (T.) See Yamasaki (S.)	C4-297	A theoretical study of hydro- gen exodiffusion in a-Si:H	
Yoshino (K.) See Shirafuji		comparison with conductivity	
(J.)	C4-583	measurements	C4-815
Yous (B.) See Donnadieu (A.)	C4-655	Zeyher (R.) See Dandoloff (R.)	C4-205

.

Contract to the second of the second

:

. b

production of the contract of

c alla (1) in som in 1, 1, 1.

the state of the s

COLLOQUE Nº 5 - 1981 (Tome I et Tome II)

Seventh International Conference on Internal Friction and Ultrasonic Attenuation in Solids

Lausanne (Suisse), 1981

Agyeman (K.), Armbruster (E.), Künzi (H.U.), Das Gupta (A.)			C5-569
and Güntherodt (HJ.) Hydrogen related internal		Atrens (A.) See Ritchie (I.G.)	C5-319
friction peaks in metallic glasses	C5-535	Atsumi (T.) See Tanaka (K.)	C5-139
Ahlers (M.) See Ghilarducci de Salva (A.)	C5-1001	Augustyniak (B.) and Chomka (W.) AG effect in equiatomic Co-Pt alloy	C5-621
Akao (F.) See Miyasato (T.)	C5-665	Azou (P.) See Carrard (M.)	C5-187
Akao (F.) See Togushi (M.)	C5-701	Baudraz (F.), Tirbonod (B.) and Gotthardt (R.) Influence of	
Alefeld (G.) See Qi (Zh.)	C5-763	remaining impurity concentra- tion on the internal friction	
Ali (A.R.), Farid (Z.), Kamel (R.) and Said (G.). On the role of Mg atoms on the reco-		after cyclic deformation in pure aluminium	C5-229
very on cold-worked A1-5.5	C5-313	Benoit (W.) See Diallo (C.)	C5-957
Al'Shits (V.I.), Kobelev (N.P.)		Benoit (W.) See Gremaud (G.)	C5-163
and Soifer (Ya.M.) Dynamic dislocation damping in me-		Benoit (W.) See Gremaud (G.)	C5-369
tals	C5-345	Benoit (W.) See De Lima (F.)	C5-85
Amano (S.) See Okuda (S.)	C5-67	Benoit (W.) See Millet (P.)	C5-929
Amirault (J.P.) See Riviere (A.)	C5-439	Benoit (W.) See Schaller (R.).	C5-881
Anderson (M.P.) and Nowick		Beretz (D.), Hillairet (J.) and Halbwachs (M.) Mobility pa-	
(A.S.) Relaxation peaks		rameters for the vacancies and	
produced by defect complexes in cerium dioxide doped with		the self-interstitials in con- centrated α-AgZn alloys	C5-747
trivalent cations	C5-823	constitution a region and a region of the re	03 / 4/
Armbruster (E.) See Agyeman		Berrisford (R.A.) and Leak (G.M.). An internal friction study of	
(K.)	C5-535	phase transformations in magne- sium-indium alloys	
Astie (B.) and Degauque (J.) Influence of the dislocation		Berry (B.S.) and Pritchet (W.C.)	
structures on the behaviour of magnetic domain walls in		Defect studies of thin layers by the vibrating-reed techni-	
high purity iron	C5-627	que	C5-1111
Astie (P.), Peyrade (J.P.) and Groh (P.). Thermally activa-	•	Bertin (Y.A.), Gacougnolle (J.L.), Sarrazin (S.) and De Fouquet	
ted evolution of the disloca-		(J.) Internal friction tran-	
tion structure generated by		sient effects of titanium hy- drogen solid solution	C5-175
low temperature deformation and associated damping pheno-			35 113
mena in pure iron	C5-91	Birnbaum (H.K.) See Mazzolai (F.M.)	C5-769

Boch (P.) See Fargeot (D.)	C5-899	Cavaille (J.Y.) See Etienne (S.)	C5-1129
Bonjour (E.) See Khoder (A.F.)	C5-1085	Chamberod (A.) See Chambron	
Bonetti (E.), Castellani (L.), Evangelista (E.) and Gondi		(W.)	C5-511
(P.) Dynamic internal friction in Al	C5-433	Chambron (W.) and Chamberod (A.) Structural relaxation and atomic mobility by magnetic	3
Britton (W.G.B.) See Wang (Y.T.)	C5-387	anisotropy measurements in some metallic glasses	C5-511
Bronsveld (P.M.) See De Rooy	C5-923	Chang (Y.K.) Kojima (K.)	C5-989
• •	03 720	Chatain (D.) See Demont (P.)	C5-563
Bussière (J.F.) See Faucher (B.)	C5-1091	Cheeke (J.D.N.), Madore (G.) and Hikata (A.) Magnetic field	
Cannelli (G.) and Cantelli (R.) A study of hydrogen accumula- tion around alloying titanium		dependent relaxation time in p-InSb at low temperatures	C5-689
in niobium by the internal friction technique	C5-793	Chen (H.B.) See Igata (N.)	C5-193
Cannelli (C.) and Cantelli (R.)	03 773	Chen (T.K.) See Wu (Z.L.)	C5-841
The influence of thermal cy- cling on the acoustic emission during hydride formation in niobium	C5-947	Chicois (J.), Hamel (A.), Fougères (R.), Esnouf (C.), Fantozzi (G.) and Perez (J.).— Internal stress measurement in aluminium by "Dip-test" method and	3
Cannelli (G.) See Koiwa (M.)	C5-787	correlation with the Bordoni relaxation evolution	C5-169
Cantelli (R.) See Cannelli (G.)	C5-793	Chih-Hsiou Tang, Mei Su and Ting-	
Cantelli (R.) See Cannelli (C.)	C5-947	Chuan Lei Relationship bet- ween internal friction and temper brittleness of alloy	017
Cantelli (R.) See Koiwa (M.)	C5-787	steels	C5-917
Capelle (E.) See Lebienvenu (M.)	C5-905	Chomka (W.) and Denga (E.). The influence of thin surface layers of mineral oils on the damping capacity of metal sam-	
Caro (J.A.) and Mondino (M.A.) Peaking effect, experiments		ples	C5-1165
and theory	C5-295	Chomka (W.) See Augustyniak (B.)	C5÷621
Carrard (M.), Prioul (C.), Plusquellec (J.) and Azou (P.) Behaviour of Fe-Ni-C alloy doped with hydrogen in		Chou (L.S.) and Letcher (S.V.) Ultrasonic propagation in the liquid crystal phases of p-	
the γ phase	C5-187	alkoxybenzoic acids	C5-587
Carrard (M.) See Prioul (C.)	C5-1067	Claus (R.O.) Attenuation of ultrasonic interface waves on	
Castellani (L.) See Bonetti (E.)	C5-433	metal-polymer-metal bounda- ries	C5-1159
Cavaille (J.Y.) See Etienne	C5_575	Clauss (A.) See Heulin (B.)	C5-829

Cou	mach (M.) See Khoder (A.F.).	C5-1085	Diallo (C.), Mondino (M.) and	
Dao	ou (J.N.) See Vajda (P.)	C5-799	Benoit (W.) Precipitation and recrystallization in Al	05 057
Das	Gupta (A.) See Agyeman		Mn alloys	C5-957
	(K.)	C5-535	Diehl (J.) See Weller (M.)	C5-817
De	Batist (R.) and Eersels (L.) Internal friction and modulus		Drescher (E.) See Schmidt (H.)	C5-339
	effects at the phase transi-		Dubois (B.) See Lebienvenu	
	tions in ferroelastic lead orthovanadate	C5-1103		C5-905
Do	Patiet (D) - Con Wissala		Dubois (B.) See Lebienvenu	05 011
De	Batist (R.) See Miyada- Naborikawa (L.T.)	C5-283		C5-911
De	Fouquet (J.) See Bertin		Dufresne (J.F.), Magalas (L.B.) and Moser (P.) Amplitude	
	(Y.A.)	C5-175	dependence of internal fric-	
	,,		tion in iron at low tempe-	
De	Fouquet (J.).— See Woirgard (J.)	C5-407	rature	C5-103
			Dufresne (J.F.) See Magalas	
Deg	<pre>gauque (J.) and Zarembowitch (A.) Introduction to the</pre>	4	(L.B.)	C5-127
	study of damping effects due		Du Jia Ju A new internal fric-	
	to magnetic, electronic pro- perties and to phonons inter-		tion peak associated with tra- ce hydrogen in Ti-6Al-4V	
	actions	C5-607	alloy	C5-775
	gauque (J.) See Astie (B.)	C5-627	Eersels (L.) See De Batist	C5-1103
De	Hosson (J.Th.M.) See De Rooy	as 022	Parala (T.) Can Marala	
	(A.)	C5-923	Eersels (L.) See Miyada- Naborikawa (L.T.)	C5-283
Del	laey (L.) See Sugimoto (K.).	C5-1019	Elbaum (C.) Ultrasonic studies	
Del	laey (L.) See Vandeurzen		of phase transitions	C5-855
	(U.)	C5-1169		
Del	laey (L.) See Van Humbeeck		Elbaum (C.) See Park (G.)	C5-517
	(J.)	C5-1007	Elsayed (T.) See Demont (P.)	C5-563
De	Lima (F.) and Benoit (W.)		Ersoy (Y.) Attenuations of plane	:
	Internal friction study of		waves in deformable, thermo-	
	the y peak in cold-worked	C5-85	electric solids subjected to a primary magnetic field in	
	niobium		an arbitrary direction	C5-653
рег	mont (P.), Chatain (D.), Elsayed (T.), Lacabanne (C.), Ronarc'h	1	Esnouf (C.) and Fantozzi (G.)	
	(D.) and Moura (J.L.). Thermo-	- 1.	Medium temperature internal	
	stimulated creep for the study		friction in high purity f.c.c.	
	of mechanical energy losses in		and h.c.p. metals	C5-445
	polymers	C5-563	Esnouf (C.) See Chicois (J.)	C5-169
Dei	nga (E.) See Chomka (W.)	C5-1165		
		*	Esnouf (C.) See Gabbay (M.)	C5-289
De	Rooy (A.), Bronsveld (P.M.)		Esnouf (C.) See San Juan (J.).	C5-43
	and De Hosson (J.Th.M.) Internal friction in ternary			
	α-CuNiZn alloys	C5-923		

Esnouf (C.) See Seyed Reihani	C5-145	temperature	C3-10/3
(S.M.) Etienne (S.), Cavaille (J.Y.),	03-143	Fernandez (A.) See Parisot	C5-569
Fouquet (F.) and Perez (J.) Inelastic properties of Se		Fiore (N.F.) See Lunarska (E.)	C5-121
based glasses in the very low frequency range	C5-575	Fiore (N.F.) See Lunarska (E.)	C5-1147
Etienne (S.), Cavaille (J.Y.), Perez (J.) and Salvia (M.)		Fiore (N.F.) See Zielinski (A.)	
Automatic system for micro- mechanical properties analy- sis	C5-1129	Fossheim (K.) See Holt (R.M.).	C5-1097
Evangelista (E.) See Bonetti		Fossum (J.O.) See Holt (R.M.).	C5-1097
(E.)	C5-433	Fougères (R.) See Chicois (J.)	C5-169
Ewert (S.), Hof (A.), Lenz (D.) and Schmidt (H.) Ultrasonic		Fouquet (F.) See Etienne (S.).	C5-575
attenuation and sound veloci- ty anomalies below 5K in		Frank (W.) See Moser (N.)	C5-647
LaAl ₂	C5-695	Fujimoto (K.) See Iwasaki (Y.)	C5-475
Fantozzi (G.) and Ritchie (I.G.). Internal friction caused by	-	Fujimoto (K.) See Iwasaki (Y.)	C5-493
the intrinsic properties of dislocations	C5-3	Gabbay (M.), Esnouf (C.), Vincent (A.) and Fantozzi (G.). Point defect-dislocation interaction	
Fantozzi (G.) See Chicois (J.)	C5-169	in Mg 0	C5-289
Fantozzi (G.) See Esnouf (C.).	C5-445	Gacougnolle (J.L.) See Bertin (Y.A.)	C5-175
Fantozzi (G.) See Gabbay (M.).	C5-289	Gault (C.) See Fargeot (D.)	C5-899
Fantozzi (G.).— See San Juan (J.)	C5-43	Girard (P.) See Lauzier (J.)	C5-247
Fantozzi (G.) See Seyed Reihani (S.M.)	C5-145	Ghilarducci de Salva (A.) and Ahlers (M.) Anelastic effects in single crystals of β-Cu-Zn	3
Fantozzi (G.) See Vincent (A.)	C5-393	and β-Cu-Zn-Al alloys	C5-1001
Fargeot (D.), Gault (C.), Platon (F.) and Boch (P.) Elastic		Gobin (P.F.) See Morin (M.)	C5-1013
and anelastic effects associa- ted with precipitation pheno-		Gobran (N.K.) See Kamel (R.)	C5-505
mena in non stoichiometric spinels	C5-899	Golosovski (M.A.) and Soifer (Ya.M.) The investigation of dislocation photodamping	
Farid (Z.) See Ali (A.R.)	C5-313	in NaCl with F-centers	C5-301
Faucher (B.), Bussière (J.F.), Snead, Jr. (C.L.) and Suenaga (M.) Internal friction and Young's modulus of Nb ₃ Sn between 6 and 300 K	C5-1091	Golosovski (M.A.), Kobelev (N.P.) and Soifer (Ya.M.) Mecha- nism of the N-S transition effect on the amplitude de-	
Feltham (P.) Internal friction	33 1371	pendent damping and modulus defect in lead	C5-1079
at kHz-frequencies in super- conductors near the critical		Gondi (P.) See Bonetti (E.)	C5-433

Gotthardt (R.) and Mercier (0.) Relation between the damping		Halbwachs (M.) See Beretz (D.)	C5-747
spectrum and the dislocation		Hamel (A.) See Chicois (J.)	C5-169
motion in martensitic alloys.	C5-995		
Gotthardt (R.) See Baudraz	C5-229	Hammerschmid (F.), Kirchner (H.O.K and Schoeck (G.) Internal)
(10)	CJ-229	friction in the alkali me-	C5-31
Granato (A.V.), Schwarz (R.B.) and Kneezel (G.A.) A model		Hanada (R.), Shinohara (M.), Sado	
for ultrasonic effects during an FCC-HCP martensitic transformation	OF -10FF	(Y.) and Kimura (H.) An internal friction study of	
	C5-1055	N-H and O-H pairs in single crystal of niobium at low	
Granato (A.V.) See Hultman (K.L.)	C5-753	temperatures	C5-757
Granato (A.V.) See Kneezel	C5-1061	Haneczok (G.), Poloczek (T.) and Moroń (J.W.) Internal fric-	
	C3-1001	tion in molybdenum containing nitrogen in solid solution	
Granato (A.V.) See Lücke (K.).	C5-327	form	C5-811
Grau (R.) and Schultz (H.) Dis- location relaxation peaks in		Hanna (B.F.) See Kamel (R.)	C5-505
molybdenum	C5-49	Harangozó (I.Z.) See Kiss (S.)	C5-529
Gremaud (G.) A new technique of measurement by coupling of a		Harangozó (I.Z.) See Kiss (S.)	C5-963
cyclic low frequency stress with an ultrasonic wave	C5-1141	Hatano (K.) See Yamane (T.)	C5-481
0 1 (0) 1 D (1 / 1)		Hausch (G.) and Török (E.) In-	
Gremaud (G.) and Benoit (W.) The interactions between		fluence of the thermoelastic martensitic transformation on	
dislocations and point defects		the elastic and anelastic pro-	
studied by ultrasonic measu- rements during a cyclic de-		perties of β_1 -CuAlNi alloys	C5-1031
formation	C5-369	Hausch (G.) See Schneider (W.)	C5-635
Gremaud (G.) and Benoit (W.) Thermally activated double		Hayakawa (H.) See Yoshida (I.)	C5-1123
kink generation studied by		Heulin (B.) and Clauss (A.) In-	
ultrasonic measurements du-		fluence of ageing on the in-	
ring a cyclic deformation	C5-163	ternal friction of Nb-Ti-N alloys	C5-829
Groh (P.) and Schultz (H.)			
Dislocation relaxation pro- cesses in metals	C5-25	He Yizhen See Mo Chi-Mei	C5-523
Cesses In metals	05 25	Hikata (A.) See Cheeke	
Groh (P.) See Astie (P.)	C5-91	(J.D.N.)	C5-689
Guenin (G.) See Morin (M.)	C5-1013	Hikata (A.) See Park (G.)	C5-517
Guntherodt (HJ.) See Agyeman	05525	Hiki (Y.), Kosugi (T.) and Kino	
(K.)	C5-535	(T.) Pressure dependence of damping constant of disloca-	05 262
Haga (K.) Magnetomechanical damping and domain structures			C5-363
in grain oriented silicon		Hiki (Y.) and Tamura (J.) - Ultra-	
steel sheets	C5-615	sonic attenuation in ice crystals near the melting tempera-	
Hakomori (K.) See Igata (N.)	C5-61	ture	C5-547

Hillairet (J.) See Beretz (D.)	C5-747	Jacques (H.) and Robrock (KH.)
	ar (0r	Elastic after effect studies
Hof (A.) See Ewert (S.)	C5-695	of molybdenum after electron irradiation at 4.7 K C5-723
Holder (J.) See Hultman (K.L.)	C5-753	
Holt (R.M.), Fossheim (K.) and		Jacques (H.) See Kollers (G.). C5-729
Fossum (J.O.) Dynamic cri-		Johnson (M.P.) See Trivisonno
tical behaviour in sound pro-	C5-1097	(J.)
		Kamel (R.), Gobran (N.K.), Saleh
Hoshino (K.) See Otani (T.)	C5-935	(S.) and Hanna (B.F.) Anelas- ticity and structural relaxa-
Hultman (K.L.), Holder (J.) and		tion of the metallic glass
Granato (A.V.) Tunneling of mixed dumbbells in Al-Zn	C5-753	Fe ₄₀ Ni ₄₀ B ₂₀ C5-505
		Kamel (R.) and Najeeb (E.) Re-
Hunklinger (S.) Dynamic proper-		laxation of molecular confor-
ties of disordered solids at low temperatures	C5-595	mation of plasticized poly (vinyl chloride) (PVC) C5-557
10w temperatures	03 373	(vinyi emoriae) (ivo)
Igata (N.), Chen (H.B.), Miyahara (K.) and Uba (T.).— Relaxation		Kamel (R.) See Ali (A.R.) C5-313
peaks of cold worked 304L		Kaufmann (H.J.), Pal-Val (P.P.),
stainless steel	C5-193	Schulze (D.) and Startsev
Toots (N) Minchara (V)		(V.I.) Amplitude-dependent internal friction in high pu-
Igata (N.), Miyahara (K.), Hakomori (K.), Miyagi (T.)		rity molybdenum single crys-
and Yoshida (H.) α and β		tals in the temperature range
peaks of cold worked and hy-		5.9-300 K C5-55
drogen charged molybdenum	C5∸61	Kawabe (H.) and Kuwahara (K.)
Ilczuk (J.) See Zgadzaj (J.)	C5-835	High damping and modulus cha-
		racteristics in a super plastic
Ishioka (S.) See Koiwa (M.)	C5-787	Zn-Al alloy
Ito (K.), Kobayashi (M.) and		Kawai (Y.) and Ogawa (T.) Inter-
Tsukishima (M.) High damping		nal stress and ΔE -effect of a Mn ferrite single crystal C5-633
capacity on Mn-base γ-phase alloys	C5-641	rm leffice single crystal CJ-033
		Kê (T.S.) Low frequency inter-
Iwasa (I.), Saito (N.) and Suzuki		nal-friction peaks as a func-
(H.) Amplitude-dependent sound attenuation in bcc ³ He		tion of strain amplitude in cold-worked dilute aluminium
crystals with ⁴ He impurities.	C5-37	alloys
Iwasaki (K.) An anomalous in-		Kê (T.S.) Internal friction
	C5-951	associated with high-tempera-
Township (W.) and Walderson (W.)		ture dislocation mobility C5-421
Iwasaki (Y.) and Fujimoto (K.) Effect of grain boundary segre-	_	Kê (T.S.) See Kong (Q.H.) C5-265
gation of antimony on relaxa-		ne (1000). Dee nong (40m).
tion at grain boundaries in silicon-iron alloys	C5-475	Kê (T.S.) See Sun (Z.Q.) C5-451
	03 473	Kê (T.S.) See Weller (M.) C5-817
Iwasaki (Y.) and Fujimoto (K.) Kinetics of secondary recrys-		Kedves (F.J.) See Kiss (S.) C5-529
tallization in grain-oriented		1100 Page 100 Page 10
silicon steel studied by high-		Kedves (F.J.) See Kiss (S.) C5-963
temperature background	C5-493	

Rhoder (A.F.), Couach (M.) and Bonjour (E.) Ultrasonic at-		crystals	C5-989
tenuation and occurence of		Volleys (C) Jacques (N) Pake	
the martensitic transforma-		Kollers (G.), Jacques (H.), Rehn (L.E.) and Robrock (KH.).	
tion in the high T _c Al5 com-		Trapping of self-interstitials	
pound V ₃ Si	C5-1085	at oversized impurities in	
	03 1003	Al-Mg and Cu-In and undersized	
Kimura (H.) See Hanada (R.)	C5-757	impurities in Al-Fe	C5-729
		Tapor It It I te	03-129
Kino (T.) See Hiki (Y.)	C5-363	Kong (Q.H.), Li (G.Y.) and Kê	
		(T.S.) - A low temperature	
Kirchner (H.O.K.) See		cold work internal-friction	
Hammerschmid (F.)	C5-31	peak in Al-0.5 wt % Ga	C5-265
W1 (0) W-1 (B 7)			
Kiss (S.), Kedves (F.J.) and		Kong (Q.P.), Lücke (K.) and	
Harangozó (I.Z.) Grain boun-		Sokolowski (G.) - A study of	
dary and precipitation effects in different AlMn alloys	OF 062	the internal friction in con-	05 071
In different Almi alloys	C3-963	centrated copper alloys	C5-271
Kiss (S.), Posgay (G.), Harangozó		Koshimizu (S.) See Tirbonod	
(I.Z.) and Kedves (F.J.)		(B.)	C5-1043
Structural relaxation and crys-		,	
tallization of FeB and NiP me-		Kosugi (T.) See Hiki (Y.)	C5-363
tallic glasses followed by in-			
ternal friction and modulus		Kronmüller (H.) See Moser (N.)	C5-647
measurements	C5-529		
7 1 (0) 1 0		Künzi (H.U.) See Agyeman (K.).	C5-535
Kneezel (G.A.) and Granato (A.V.)	, -	Vinnerachi (N.) - Can Chila (C.)	OF 67
Ultrasonic attenuation and ve-		Kuramochi (N.) See Okuda (S.).	C5-67
locity changes during the FCC- HCP martensitic transformation		Kurosawa (T.) See Otani (T.)	C5-935
in cobalt-nickel	C5-1061	Ruiosawa (1.). See Otani (1.).	03-333
IN CODULE HICKCIPPOPPE	03 1001	Kuwahara (K.) See Kawabe (H.).	C5-941
Kneezel (G.A.) See Granato			
(A.V.)	C5-1055	Lacabanne (C.) See Demont (P.)	C5-563
Kobayashi (M.) See Ito (K.)	C5-641	Lauzier (J.), Girard (P.) and	
T 1 1 (Y D) 0 . A11014.		Minier (C.) A study of the	
Kobelev (N.P.) See Al'Shits	OF 245	stability of irradiation in- duced point defects on various	
(V.I.)	C5-345	dislocation classes in copper	
Kobelev (N.P.) See Golosovski		and silver	C5-247
(M.A.)	C5-1079		
(, , , , , , , , , , , , , , , , , ,		Lauzier (J.) and Minier (C.)	
Koiwa (M.), Ishioka (S.), Cannelli	L	A proposal to identify the	
(G.) and Cantelli (R.) Me-		dislocation classes giving	
chanical relaxation of substi-		rise to different Bordoni	
tutional-interstitial (tetra-		relaxation processes in cop-	
hedral) atom complexes in the		per	C5-133
bcc lattice	C5-787		
	or 701	Leak (G.M.) See Berrisford	CE . 007
Koiwa (M.) See Tanaka (S.)	C5-781	(R.A.)	C5-887
Voice (M) - Con Vachinari (0.)	C5-893	Lebienvenu (M.), Capelle (E.),	
Koiwa (M.) See Yoshinari (0.).	33 373	Ocampo (G.) and Dubois (B.).	
Kojima (K.), Shimomura (K.) and		Internal friction measurements	
Chang (Y.K.) Dispersion		during the tempering of the	
anomaly of ultrasonic wave		1000°C as quenched Cu-10.1	
in Cu-Zn crystals and V ₃ Si		wt % Al alloy	C5-905
,			

Lebienvenu (M.) and Dubois (B.) Influence of the nitro-		Lücke (K.) See Marx (F.P.)	C5-241
gen content on the internal friction of Fe-16.6 wt % Cr		Lücke (K.) See Schmidt (H.)	C5-339
alloys	C5-911	Lücke (K.) See Schmidt (H.)	C5-351
Lenz (D.), Lücke (K.) and Schmidt (H.). Evaluation of dislocation pinning effects from si-		Lunarska (E.) and Fiore (N.F.) Rayleigh wave studies of ca- thodic H-charging of Fe	C5-121
multaneous measurements of attenuation and sound velo- city in copper	C5-381	Lunarska (E.) and Fiore (N.F.) Development of a surface ul- trasonic wave technique for	:
Lenz (D.) See Ewert (S.)	C5-695	the study of H entry into	C5-1147
Lenz (D.) See Schmidt (H.)	C5-339	Lunarska (E.) See Zielinski	
Lenz (D.) See Schmidt (H.)	C5-351	(A.)	C5-181
Lenz (D.) See Schrey (P.)	C5-671	Madore (G.) See Cheeke	
Lenz (D.) See Schulz (J.)	C5-151	Magalas (L.B.), Dufresne (J.F.)	
Letcher (S.V.) See Chou (L.S.)	C5-587	and Moser (P.). The Snoek- köster relaxation in iron	
Li (G.Y.) See Kong (Q.H.)	C5-265	Magalas (L.B.) and Moser (P.)	
Li Cheng-Hsiu See Li Wern-Bin.	C5-463	Internal friction in cold worked iron	C5-97
Li Cheng-Hsiu See Li Wern-Bin.	C5-469	Magalas (L.B.) See Dufresne	
Liu Zheng-Qun See Li Wern-Bin.	C5-463	(J.F.)	C5-103
Liu Zheng-Qun See Li Wern-Bin.	C5-469	Marx (F.P.), Sokolowski (G.) and Lücke (K.) Influence of oxi-	
Li Wern-Bin, Liu Zheng-Qun, Yang Quo-Ping, Li Cheng-Hsiu and Zhang Bin Investigation of		dizing and reducing pre-treat- ments on the internal fric- tion of electron irradiated	
the distribution and the state of small amount of rare earth		copper	C5-241
elements in pure iron by the method of internal friction	C5-463	Matsui (H.) and Schultz (H.) The effect of hydrogen char-	
Li Wern-Bin, Yang Guo-Ping, Li		ging on the dislocation re-	C5-115
Cheng-Hsiu and Liu Zheng-Qun		laxation in iron	
An internal friction study of the mechanism of improvement		Matsumoto (K.). See Yamane (T.)	C5-805
of the properties of Fe-Cr-Al alloys by the addition of a		Maul (M.) and Schultz (H.) Hydrogen-dislocation inter-	
small amount of rare earth elements	C5-469	cations and intrinsic dislo-	C5-73
Lücke (K.) and Granato (A.V.) The rigid rod model of dislo- cation resonance including applications to point defect		Mazzolai (F.M.) Contribution to the characterization of the anelastic properties of multi-phase systems	C5-867
drag	C5-327		33 007
Lücke (K.) See Kong (Q.P.)	C5-271	Mazzolai (F.M.) and Birnbaum (H.K.) Ultrasonic attenua- tion and velocity studies in	
Lücke (K.) See Lenz (D.)	C5-381	the $\alpha-\alpha$ ' solid solution phases of the Nb-H system	C5-769

McCrum (N.G.) Unresolved pro- blems in polymeric anelasti-		Mondino (M.) See Diallo (C.)	C5-957
city	C5-501	Morin (M.), Guenin (G.) and Gobin (P.F.). Internal friction mea-	
Mazot (Ph.) See Woirgard (J.).	C5-1135	surements related to the two way memory effect in Cu-Zn-Al	
Mei Su See Chih-Hsiou Tang	C5-917	alloy exhibiting thermoelastic martensitic transformation	C5-1013
Mercier (0.), Tirbonod (B.) and Török (E.) The internal fric- tion spectrum of premartensi-		Moroń (J.W.) See Haneczok (G.)	C5-811
tic transformations	C5-1037	Moroń (J.W.) See Zgadzaj (J.).	C5-835
Mercier (0.) See Gotthardt	C5-995	Moser (P.) Introduction for the point defect relaxation session	C5-721
Michard (F.) and Plicque (F.) Ultrasonic absorption in some highly attenuating ionic-mole- cular crystals	C5-683	Moser (N.), Kronmüller (H.) and Frank (W.) Magnetic relaxa- tions in amorphous ferromagne-	
Millet (P.), Schaller (R.) and		tic Fe _{80-x} Ni _x P _{20-y} B _y alloys	C5-647
Benoit (W.) Characteristic internal friction spectrum of		Moser (P.) See Dufresne (J.F.)	C5-103
grey cast iron	C5-929	Moser (P.) See Magalas (L.B.).	C5-97
Minamida (K.) See Yoshida (I.)	C5-1123	Moser (P.) See Magalas (L.B.).	C5-127
Minier (C.) See Lauzier (J.)	C5-133	Moser (P.) See Vajda (P.)	C5-799
Minier (C.) See Lauzier (J.)	C5-247	Moser (P.) See Weller (M.)	C5-741
Miyada-Naborikawa (L.T.), De Batis (R.) and Eersels (L.) Inter-		Motegi (M.) See Yoshida (I.)	C5-1123
nal friction in zirconium		Moura (J.L.) See Demont (P.)	C5-563
Miyagi (T.) See Igata (N.)	C5-61	Naithani (U.C.) and Semwal (B.S.). Sound attenuation in ferro-	
Miyahara (K.) See Igata (N.)	C5-61	electric solids	C5-659
Miyahara (K.) See Igata (N.)	C5-193	Najeeb (E.) See Kamel (R.)	C5-557
Miyasato (T.), Tokumura (M.) and Akao (F.) Acoustic attenua- tion by shallow donors in ger-		Nakamura (Y.) See Sugimoto (K.)	C5-1019
manium in a magnetic field	C5-665	No (L.) See San Juan (J.)	C5-43
Mizubayashi (H.) and Okuda (S.) Anelastic study on low ampli- tude fatigue of copper		Nowick (A.S.) See Anderson (M.P.)	C5-823
Mizubayashi (H.) See Okuda	C5-67	Ocampo (G.) See Lebienvenu	C5-905
Mo Chi-Mei, Shui Jiapeng and He		Ogawa (T.) See Kawai (Y.)	C5-633
Yizhen Internal friction peaks in amorphous Pd ₈₀ Si ₂₀ alloys	C5-523	Ogurtani (T.Ö.) Unified theory of dislocation damping with a special reference to point-	05-025
Manding (M A) - See Caro (T.A.)	C5-295	defect dragging	C5-235

Ohsone (H.) See Yoshida (I.)	C5-1153	Peyrade (J.P.) See Astie (P.).	C5-91
Okuda (S.), Mizubayashi (H.), Kuramochi (N.), Amano (S.), Shimada (M.) and Tatekawa		Platkov (V.Ya.) See Pal-Val (L.N.)	C5-253
(H.) Dislocation motion in Va metals studied by internal friction	C5-67	Platkov (V.Ya.) See Pal-Val (P.P.)	C5-259
Okuda (S.) See Mizubayashi	05 07	Platon (F.) See Fargeot (D.)	C5-899
(H.)	C5-157	Plicque (F.) See Michard (F.).	C5-683
Otani (T.), Hoshino (K.) and Kurosawa (T.) Damping		Plicque (F.) See Vacher (R.)	C5-553
capacity and mechanical properties of Zn-Al alloy castings	C5-935	Plusquellec (J.) See Carrard (M.)	C5-187
Pal-Val (L.N.), Platkov (V.Ya.) and Roshchupkin (A.M.) dis-		Poloczek (T.). – See Haneczok	C5-811
location amplitude dependent ultrasound absorption in nor-		Posgay (G.) See Kiss (S.)	C5-529
mal and superconducting lead containing diverse valency impurities	C5-253	Prioul (C.) and Carrard (M.) The influence of athermal transformation and cold-wor-	
Pal-Val (P.P.), Platkov (V.Ya.) and Startsev (V.I.). On the internal friction in antimony		king on the low temperature (77-300)K internal friction behaviour of Fe-Ni-C alloy	C5-106
caused by unpinning of over- damped and underdamped dislo-		Prioul (C.) See Carrard (M.)	C5-187
cations	C5-259	Pritchet (W.C.) See Berry (B.S.)	C5-111
Pal-Val (P.P.) See Kaufmann (H.J.)	C5-55	Qi (Zh.), Völkl (J.), Wipf (H.) and Alefeld (G.) The influ-	
Parisot (J.), Astier (J.), Fernandez (A.) and Rafi (O.) Internal friction of glassy		ence of N-impurities on the diffusion coefficient of H and D in Nb and Ta	C5-763
polycarbonate during plastic flow	C5-569	Rafi (0.) See Parisot (J.)	C5-569
Park (G.), Hikata (A.) and Elbaum	03 307	Rehn (L.E.) See Kollers (G.)	C5-729
(C.) Ultrasonic studies of metallic glasses at low tempe-			03-729
ratures	C5-517	Revel (G.) See Seyed Reihani (S.M.)	C5-145
Pelous (J.) See Vacher (R.)	C5-553	Ritchie (I.G.), Atrens (A.), SO (C.B.) and Sprungmann (K.W.)	
Perez (J.) See Chicois (J.)	C5-169	Dislocation relaxation in a random array of solutes	C5-319
Perez (J.) See Etienne (S.)	C5-575	Ritchie (I.G.) and Sprungmann	03 317
Perez (J.) See Etienne (S.)	C5-1129	(K.W.) Dislocation core diffusion and unpinning in	
Perez (J.) See Tatibouet (J.).	C5-541	α-Ζr	C5-427
Perez (J.) See Vincent (A.)	C5-393	Ritchie (I.G.) See Fantozzi	C5-3
Perrin (B.) Ultrasonic relaxa-	C5-677		

Riviere (A.), Amirault (J.P.) and Woirgard (J.) High tem- perature internal friction and dislocation motion in poly and		Schaller (R.) and Benoit (W.) Precipitation stages in Al-Ag alloys determined by the evo- lution of the internal fric-	
single crystals of F.C.C. me- tals	C5-439	tion spectrum	C5-881
Riviere (A.) See Woirgard (J.)	C5-407	Schaller (R.) See Millet (P.).	C5-929
Rivière (A.) See Woirgard (J.)	C5-1135	Schindlmayr (R.).— See Schlipf (J.)	C5-357
Robrock (KH.) Mechanical re- laxation studies of point defects in metals	C5-709	Schlipf (J.) and Schindlmayr (R.) On the dislocation resonance peak in deformed crystals	C5-357
Robrock (KH.) and Schober (H.R.) Models for frozen-free-split phenomena in e-irradiated Al and Al-Fe	c5-735	Schmidt (H.), Lenz (D.), Drescher (E.) and Lücke (K.).— Ultrasonic investigation of the dislocation structure in plastically	
Robrock (KH.) See Jacques (H.)	C5-723	deformed copper	
Robrock (KH.) See Kollers	C5-729	Schmidt (H.), Lenz (D.) and Lücke (K.). Temperature dependence of dislocation resonance dam-	
Rodrian (U.) and Schultz (H.)		ping	C5-351
Dislocation relaxation peaks in tantalum, intrinsic and	C5-70	Schmidt (H.) - See Ewert (S.)	C5-695
impurity dependent effects	C5-79	Schmidt (H.) See Lenz (D.)	C5-381
Ronarc'h (D.) See Demont (P.).	C5-563	Schmidt (H.) See Schrey (P.)	C5-671
Roshchupkin (A.M.). See Pal-Val	C5-253	Schneider (W.), Schrey (P.), Hausch (G.) and Török (E.) Damping capacity of Fe-Cr and	
Sado (Y.) See Hanada (R.)	C5-757	Fe-Cr based high damping alloys	C5-635
Said (G.) See Ali (A.R.)	C5-313	Schober (H.R.) See Robrock	
Saito (N.) See Iwasa (I.)	C5-37	(KH.)	C5-735
Sakamoto (K.) and Shimada (M.) Internal friction of deformed iron containing hydrogen	C5-109	Schoeck (G.). See Hammerschmid (F.)	C5-31
Saleh (S.) See Kamel (R.)	C5-505	Schrey (P.), Schulz (J.), Schmidt (H.) and Lenz (D.). Influence of impurities and dislocations	
Salvia (M.) See Etienne (S.)	C5-1129	on ultrasonic attenuation in copper and dilute copper alloys	
San Juan (J.), No (L.), Fantozzi (G.), Esnouf (C.) and Vanoni		at low temperatures	C5-671
(F.) Microplasticity and internal friction in pure iron.	C5-43	Schrey (P.) See Schneider (W.)	C5-635
Sarrazin (S.) See Bertin	*	Schultz (H.) See Grau (R.)	UJ-47
/V A \	C5-175	Schultz (H.) See Groh (P.)	C5-25

Schultz (H.) See Matsui (H.)	C5-115	(B.)	C5-109
Schultz (H.) See Maul (M.)	C5-73	Snoeys (R.) See Vandeurzen	
Schultz (H.) See Rodrian (U.).	C5-79	(U.)	C5-116
Schulz (J.) and Lenz (D.) In- fluence of plastic deformation		So (C.B.) See Ritchie (I.G.)	C5-319
on the temperature dependence of ultrasonic attenuation in copper single crystals	C5-151	Soifer (Ya.M.) See Al'Shits (V.I.)	C5-345
Schulz (J.) See Schrey (P.)	C5-671	Soifer (Ya.M.) See Golosovski (M.A.)	C5-301
Schulze (D.) See Kaufmann (H.J.)	C5-55	Soifer (Ya.M.) See Golosovski (M.A.)	C5-107!
Schulze (K.) See Weller (M.)	C5-817	Sokolowski (G.) See Kong (Q.P.)	C5-271
Schwarz (R.B.) See Granato	05_1055		1
(A.V.) Seeger (A.) The kink picture of	C5-1055	Sokolowski (G.) See Marx (F.P.)	C5-241
dislocation mobility and dis- location-point-defect inter- actions	C5-201	Spears (C.J.) Internal friction in concentrated alloys	C5-277
		Spencer, Jr. (J.W.) Stress re-	
Semwal (B.S.) See Naithani (U.C.)	C5-659	laxations at low frequencies in fluid-saturated rocks	C5-1175
Seyed Reihani (S.M.), Esnouf (C.), Fantozzi (G.) and Revel (G.) Effect of impurities		Sprungmann (K.W.) See Ritchie (I.G.)	C5-319
on the internal friction spectrum of magnesium after plastic deformation	C5-145	Sprungmann (K.W.) See Ritchie (I.G.)	C5-427
Seyed Reihani (S.M.) See Vincent (A.)	C5-393	Startsev (V.I.) See Kaufmann (H.J.)	C5-55
		Startsev (V.I.) See Pal-Val	CE 250
Shen Huimin See Wang Yening	C5-1049	(P.P.)	C5-259
Shestopal (V.O.) On diffusional mechanisms of relaxation. Pure diffusion and drift	- C5-847	Stephens (R.W.B.).— See Wang (Y.T.)	C5-387
Shimada (M.) See Okuda (S.)	C5-67	Suenaga (M.) See Faucher (B.).	C5-1091
Shimada (M.) See Sakamoto (K.)	C5-109	Sugai (T.) See Yoshida (I.)	C5-1123
Shimomura (K.) See Kojima (K.)	C5-989	Sugimoto (K.) Internal friction phenomena associated with dif-	
Shinohara (M.) See Hanada (R.)	C5-757	fusionless phase transformations in alloys	C5-971
Shui Jiapeng See Mo Chi-Mei	C5-523	Sugimoto (K.), Nakamura (Y.) and	
Slotwinski (A.R.) See Trivisonn (J.)	o C5-983	Delaey (L.) Pseudoelastic behavior of coil spring made of a high-damping Cu-Zn-Al	05.131
		alloy	C5-1019

tinuous-distribution disloca-		(T·)·······	C5-665
tion model of internal fric- tion associated with the in-		Török (E.). See Hausch (G.)	C5-1031
homogeneous sliding along high-angle grain boundaries	C5-451	Török (E.) See Mercier (0.)	C5-1037
Suzuki (H.) See Iwasa (I.)	C5-37	Török (E.) See Schneider (W.).	C5-635
Suzuki (T.) See Wuttig (M.)	C5-1025	Trivisonno (J.), Slotwinski (A.R.) and Johnson (M.P.). Influence of the martensitic phase trans-	
Takahashi (Y.) See Yamane (T.)	C5-481	formation on the ultrasonic attenuation and velocity in	
Tamura (J.) See Hiki (Y.)	C5-547	single crystal lithium	C5-983
Tanaka (K.), Atsumi (T.) and Yamada (M.) An internal fric		Tsukishima (M.) See Ito (K.)	C5-641
tion peak due to hydrogen-dis- location interaction in ni-		Uba (T.) See Igata (N.)	C5-193
ckel	C5-139	Vacher (R.), Pelous (J.), Plicque (F.) and Zarembowitch (A.)	
Tanaka (S.) and Koiwa (M.) An internal friction study of		Frequency dependence of sound absorption of vitreous silica	
V-Ti-H(D) alloys	C5-781	between 10 MHz and 35 GHz	C5-553
Tani (S.) See Yoshida (I.)		Vajda (P.), Daou (J.N.) and Moser (P.) Anelastic effects in	
Tatekawa (H.) See Okuda (S.)	C5-67	lutetium containing hydrogen (or deuterium) in solid solu-	
Tatibouet (J.), Perez (J.) and Vassoille (R.) Very low		tion	C5-799
frequencies internal friction measurements of ice I _h	C5-541	Vandeurzen (U.), Verelst (H.), Snoeys (R.) and Delaey (L.) Dynamic properties of high	
Ting-Chuan Lei Relationship between internal friction		damping metals	C5-1169
and creep rupture behavior of aluminum alloys	C5-487	Van Humbeeck (J.) and Delaey (L.). The influence of strain-rate, amplitude and temperature on	
Ting-Chuan Lei See Chih-Hsiou	C5-917	the hysteresis of a pseudo- elastic Cu-Zn-Al single crys-	
Tirbonod (B.) and Koshimizu (S.).	-		C5-1007
Dislocation relaxation in the martensitic phase of the ther-		Vanoni (F.). See San Juan (J.).	
moelastic NiTi and NiTiCu	C5-1043	Vassoille (R.).— See Tatibouet	
Tirbonod (B.) See Baudraz (F.)	C5-229	Verelst (H.) See Vandeurzen	C5-1169
Tirbonod (B.) See Mercier (0.)	C5-1037	Vincent (A.), Seyed Reihani	
Togushi (M.) and Akao (F.) Acoustic attenuation of sur-		(S.M.), Fantozzi (G.) and Perez (J.) Dislocations	
face acoustic waves in super- conducting thin films of amor-		point defects interactions in magnesium	C5-393
phous Bi	C5-701	Vincent (A.) See Gabbay (M.)	C5-289
Tokumura (M.) See Miyasato		Völkl (J.) See Qi (Zh.)	C5-763

Wang (Y.T.), Britton (W.G.B.) and Stephens (R.W.B.).— Non-linear interaction between ultrasonic shear waves and dislocations		Wu (Z.L.) and Chen (T.K.) Lat- tice parameter change in Fe- Ti-N alloys in relation to substitutional-interstitial	
in aluminium	C5-387	interaction peak	C5-841
Wang Yening, Shen Huimin, Xu Zirar Zou Yifeng, Zhu Jinsong, Zhang Zhifang and Yang Zhaojin In-	1,	Wuttig (M.) and Suzuki (T.) Nonlinear resonance in CuAlNi alloys	C5-102
ternal friction and elastic modulus of AuCd in the vicini- ty of martensitic transforma-		Xu Ziran See Wang Yening	C5-104
tion	C5-1049	Yamada (M.) See Tanaka (K.)	C5-139
Wang Yening and Zhu Jinsong A solute peak associated with the grain boundaries of iron		Yamane (T.) and Matsumoto (K.) Snök peaks in molybdenum	C5-80 5
containing a small amount of carbon	C5-457	Yamane (T.), Takahashi (Y.) and Hatano (K.) Internal fric- tion during steady state creep	
Weiner (D.) A low temperature peak in Young's modulus of copper single crystals	C5-325	of 25Cr-20Ni austenitic stain- less steel	C5-481
		Yang Quo-Ping See Li Wern-Bin.	C5-463
Weller (M.) and Moser (P.) Internal friction of chromium after 20 K electron-irradia-		Yang Guo-Ping See Li Wern-Bin.	C5-469
tion	C5-741	Yang Zhaojin See Wang Yening	C5-104
Weller (M.), Zhang (J.X.), Schulze (K.), Kê (T.S.) and Diehl (J.) The Snoek relaxation in tanta- lum and niobium with relati-		Yokoyama (T.) Internal friction of zinc single crystals under impact loading	C5-375
vely high concentrations of oxygen	C5-817	Yoshida (I.) and Ohsone (H.) Effect of ultrasonic vibration on the metallurgical properties	I
Weller (M.) See Wert (C.A.)	C5-581	of steel	
Wert (C.A.) and Weller (M.) In- ternal friction in coal	C5-581	Yoshida (I.), Sugai (T.), Tani (S.), Motegi (M.), Minamida (K.) and Hayakawa (H.) Auto-	
Wipf (H.) See Qi (Zh.)	C5-763	mation of internal friction	
Woirgard (J.), Mazot (Ph.) and Rivière (A.) Programmable		measurement apparatus of inver- ted torsion pendulum type	C5-112
system for the measurement of the shear modulus and internal		Yoshida (H.) See Igata (N.)	C5-61
friction, on small specimens, at very low frequencies	C5-1135	Yoshinari (0.) and Koiwa (M.) On the twist effect of V-H	
Woirgard (J.), Riviere (A.) and		alloys associated with the pre- cipitation of hydrides	- C5-893
De Fouquet (J.) Experimental and theoretical aspect of the		Zarembowitch (A.) See Degauque	
high temperature damping of pure metals	C5-407	(J.)	C5-607
		Zarembowitch (A.) See Vacher	

gadzaj (J.), Ilczuk (J.) and Moroń (J.W.) Internal fric-	Zhang Zhifang See Wang Yening.	C5-1049
tion and magnetic after-effects	Zhu Jinsong See Wang Yening	C5-457
in Fe-V-N alloys C5-835	Zhu Jinsong See Wang Yening	C5-1049
tion dynamic model of internal friction in the process of plastic deformation in metals C5-399	Zielinski (A.), Lunarska (E.) and Fiore (N.F.). Surface wave studies of H ingress and egress in stainless alloys	C5-181
Zhang (J.X.) See Weller (M.) C5-817		
Thang Rin See Li Wern-Bin C5-463	Zou Yifeng See Wang Yening	CJ1049

The following the second of th

.1

About the general Water for the composition of the second second

International Conference on Phonon Physics

Bloomington (U.S.A.), 1981

Abou-Ghantous (M.) See Khoder (A.F.)	C6-383	Arnold (W.), Billmann (A.), Doussineau (P.), Frenois (C.) and Levelut (A.) Ultrasonic absorption and dispersion in amorphous superconductors	
Adnot (A.) See Thiry (P.)	C6-843		C6-37
Agrawal (B.K.) Vibrational excit tions in a-Si:F and a-Si:F:H alloys	c6-93	Arnold (W.) See v. Löhneysen (H.)	C6-193
Aikawa (Y.) See Nakamura (T.).	C6-421	Arora (A.K.) See Sood (A.K.).	C6-572
Akao (F.) See Miyasato (T.)	C6-658	Asai (N.) See Inoue (K.)	C6-430
Alexandre (F.) See Jusserand (B.)	C6-43	Ashkenazi (J.) and Dacorogna (M.) Phonons in metals	 C6-355
Alig (R.C.), Bloom (S.) and Inoue (M.) Optical-mode deformation	n	Aubree (J.) See Schneck (J.)	C6-734
potential	C6-484	Aune (H.J.) See Fossum (J.O.).	C6-867
Aliotta (F.), Fontana (M.P.), Maisano (G.), Migliardo (P.), Vasi (C.) and Wanderlingh (F.) Vibrational dynamics in ionic liquids	C6-57	Ayache (C.) and Nuñez-Regueiro (M.) Basal thermal conductivity of TiSe2	C6-338
Almond (D.P.) and West (A.R.) A new interpretation of Ultra- sonic Attenuation in Superioni Conductors β and β"-alumina		Balasubramanian (K.R.) and Haridasan (T.M.).— A green's function study of the U centre in BaClF and SrClF	C6-905
Anastassakis (E.M.) and Cardona (M.) Simulation of long-wavelength optical phonons by generalized internal strains.	C6-537	Baldereschi (A.) See Resta (R.) Balkanski (M.) See Brunel	
Anikiev (A.A.), Umarov (B.S.), Gorelik (V.S.) and Vetelino (J.F.) Two-phonon bound state		(L.C.) Balkanski (M.) See Kanellis	C6-412
in ammonium chloride at finite temperatures	C6-152	(G.)	C6-341
Animalu (A.O.E.) See Onwuagba (B.N.)	C6-401	Ball (M.A.) The generalized pseudoatom formalism in lattice dynamics	e C6-519
Appel (J.) See Fe (D.)	06-490		
Arnold (G.B.) and Meron (M.) The local phonon density of states and its influence on superconducting properties.	c6-377	Ball (M.A.) and Leung (W.N.) The dielectric function matrix and the lattice dynamics of KCl	C6-911

Banerjee (P.K.), Bendow (B.),	Bellissent-Funel (M.C.) - See
Drexhage (M.G.), Goltman (J.T.), Mitra (S.S.) and Moynihan	Maisano (G.) C6-51
	Belushkin (A.I.) See Wasiutynski
(C.T.) Polarized Raman	
scattering studies of fluoro- zirconate and fluorohafnate	(T.) C6-59
glasses C6-75	Bendow (B.), Lipson (H.G.), Brown
	(R.N.), Marshall (R.C.), Billard
Bansal (M.L.), Deb (S.K.), Roy	(D.) and Mitra (S.S.) Tem-
(A.P.) and Sahni (V.C.)	perature dependence of multi-
Anomalous temperature dependence	phonon absorption in highly
of a LO mode in LiKSO ₄ C6-902	transparent cubic zirconia
Barchielli (A.) and Mulazzi (E.)	Bendow (B.) See Banerjee (P.K.) C6-75
Homogeneous broadening of zero-	Benedek (G.), Pollini (I.) and
phonon lines for a multilevel	Bauhofer (W.) Two-phonon pro-
system in a crystal : the role	gressions associated with
of the electron-phonon inter-	vibronic excitons in layered 3d-
actions	
	metal compounds
Basoon (S.A.) - See Kaplyanskii	Benedek (G.) See Bussmann-Holder
(A.A.)	(A.)
P (0 A)	(M))
Basoon (S.A.) See Kaplyanskii	Benedek (G.), Brusdeylins (G.),
(A.A.)	Doak (R.B.) and Toennies
n . 11 (7) n (7)	(J.P.) The spectroscopy of
Batallan (F.), Rosenman (I.),	surface phonons by inelastic
Simon (C.), Furdin (G.) and	atom scattering C6-79
Lauter (H.J.) - Phonons in	atom scattering
graphite intercalated with	Porhorich (F) and Vindor (U)
bromine	Berberich (F.) and Kinder (H.)
	The phonon spectrum emitted by
Bauernfeind (W.), Keller (J.) and	superconducting Sn tunnel
Schröder (U.) Theory of thermal	junctions
conductivity in molecular	P
crystals. Application to alcali	Beserman (R.) See Schmeltzer
cyanides C6-247	(D.)
	Beserman (R.) See Schmeltzer
Bauhofer (W.) See Benedek (G.) C6-301	(D.)
Baumgartner (R.), Engelhardt (M.)	(1)-)
and Renk (K.F.) Observation of	Beserman (R.) See Teicher (M.) C6-46
strongly frequency dependent	
lifetimes of acoustic phonons in	Bethoux (0.), Lasjaunias (J.C.)
CaF ₂ C6-119	and Ravex (A.) Effect of a
2	thermal annealing on the
Behera (S.N.) Effect of non-	lattice specific heat and
magnetic impurities on the	thermal conductivity of super-
Raman scattering by supercon-	conducting amorphous Zr.76Ni.24
ducting gap excitations in	alloys
layered compounds C6-292	
	Royalor (U II) - Cas w IShnoyan
Behera (S.N.) and Khare (A.)	Beyeler (H.U.) See v. Löhneysen
Dynamics and Thermodynamics of	(He)
a one dimensional non-linear	Riolyahkin (A.W.) Nathania (T.)
lattice in the continuum limit C6-314	Bielushkin (A.V.), Natkaniec (I.),
	Dolganov (V.K.) and Sheka
·	(E.F.). Vibrational density of
Behera (S.N.) and Samsur (Sk.)	states of amorphous and crys-
Hybridization of the two-phonon	talline liquid crystal subs-
bound state with the local mode	tances
in imperfect crystals 06_500	Pion (II) Con War (II)

Billard (D.) See Bendow (B.).	C6-140	Bortolani (V.), Nizzoli (F.),	
Billmann (A.) See Arnold (W.)	C6-37	Santoro (G.) and Marvin (A.) Theory of acoustic phonons at interfaces and Brillouin scattering	
Bilz (H.) See Brunel (L.C.)	C6-412		C6-804
Bilz (H.) See Bussmann-Holder (A.)	C6-409	Bosio (L.) See Copley (J.R.D.)	C6-386
Bilz (H.) See Büttner (H.)	C6-111	Bouillot (J.) See Cailleau (H.)	C6-704
Bilz (H.) See Kress (W.)	C6-3	Bouillot (J.) See Geick (R.).	C6-713
Bilz (H.) See Maradudin (A.A.)	C6-137	Boysen (H.) See Dorner (B.)	C6-752
Bist (H.D.) See Patel (M.B.).	C6-917	Brafman (0.) and Livescu (G.) A study of polaritons and	
Black (J.E.), Campbell (D.A.) and Wallis (R.F.) Atomic vibra- tions at (100) surfaces of		phonon peculiarities in CuCl _{1-x} Br _x	C6-96
FCC and BCC Metals	C6-837	Breazeale (M.A.) and Philip	
Blanchard (C.) See Deville (A.)	C6-465	(J.) Temperature dependence of phonon-phonon interaction in silicon	C6-134
Bloom (S.) See Alig (R.C.)	C6-484		60-134
Boatner (L.A.) See Salce (B.)	C6-424	Brito-Orta (R.) and Klemens (P.G.) Contribution of low-	
Boato (G.) See Cantini (P.)	C6-801	frequency longitudinal phonons to the lattice thermal conduc-	06 256
Boissier (M.), Vacher (R.) and Perrin (B.). Ultrasonic rela-		tivity of dielectric solids.	C6-256
xation in α-sulphur	C6-611	Brito-Orta (R.) and Klemens (P.G.) Contribution of low-	
Bokhenkov (E.L.), Kolesnikov (A.I.), Krivenko (T.A.), Sheka (E.F.) and Dementjev (V.A.)		frequency longitudinal phonons to the lattice thermal conduc- tivity of dielectric solids.	C6-256
Harmonic dynamics of anthracene and naphtalene crystals		Bron (W.E.) See Throwe (J.)	C6-849
Bokhenkov (E.L.), Kolesnikov		Brown (R.A.) The effect of	
(A.I.), Krivenko (T.A.), Sheka (E.F.), Dementjev (V.A.) and Natkaniec (I.) Neutron spec-		dislocations on thermal conduc- tivity	C6-271
troscopy of internal phonons or naphtalene and anthracene	E	Brown (R.N.) See Bendow (B.).	C6-140
crystals	C6-605	Brüesch (P.), Hibma (T.) and Bührer (W.) Phonons and ionic	c
Bokhenkov (E.L.) See Dorner (B.)	C6-602	conductivity in the two-dimen-	
	C6-404	sional super-ionic conductor AgCrS ₂	C6-178
Bömmel (H.E.) See Ströbel (B.)	C0-404	Brunel (L.C.), Landwehr (G.),	
Bonjour (E.), Calemczuk (R.), Lagnier (R.) and Salce (B.).— Low temperature thermal proper- ties of cyclohexanol: a glassy		Bussmann-Holder (A.), Bilz (H.) Balkanski (M.), Massot (M.) and Ziolkiewicz (M.K.) Influence of strong magnetic fields on) , i
crystal system	C6-63	ferroelectric phase transi-	C6-412
Bortolani (V.), Franchini (A.), Nizzoli (F.) and Santoro (G.).			
Surface lattice dynamics of nickel	C6-831	Brusdeylins (G.) See Benedek	C6-793

Buchenau (U.), Schober (H.R.) and Wagner (R.) Coherent neutron			C6-652
scattering from polycrystals	C6-395	Cardona (M.) See Pintschovius (L.)	C6-634
Bührer (W.). See Brüesch (P.).	C6-178	Cardona (M.) See Oka (Y.)	C6-459
Bussmann-Holder (A.), Benedek (G.) Bilz (H.) and Mokross (B.) Microscopic polarizability model of ferroelectric soft modes Bussmann-Holder (A.) See Brunel	C6-409	Carles (R.), Saint-Cricq (N.), Zwick (A.), Renucci (M.A.) and Renucci (J.B.) Low frequency Raman scattering in mixed Ga _{l-x} Al _x As and Ga _{l-x} In _x As alloys	C6-105
(L.C.)	C6-412		
Büttner (H.) and Bilz (H.) Some aspects of nonlinear lattice models	C6-111	Carlone (C.), Hota (N.K.), Stolz (H.J.), Elbert (M.), Kuzmany (H.) and Hochheimer (H.D.).— Gas phase Raman spectrum of TCNQ and pressure dependence of the modes in crystals of TCNQ	C6-323
	00 704	Carroll (P.J.) and Lannin	
Calemczuk (R.) See Bonjour (E.)	C6-63	(J.S.) Second order Raman scattering in crystalline sulphur selenium and	
Calleja (J.M.), Vogt (H.) and		tellurium	C6-643
Cardona (M.) Absolute cross sections for one-phonon Raman scattering from several insu- lators and semiconductors	C6-487	Casella (R.C.) Detection of Impurity tunneling in solids via coherent phonon coupling and direct neutron scattering	C6-923
Camassel (J.), Gil (B.), Merle (P.), Mathieu (H.) and Pascual		Castiel (D.) See Shen (T.J.).	C6-819
(J.) Resonant Raman scattering in TiO ₂	ng C6-682	Caudano (R.) See Thiry (P.)	C6-843
Camassel (J.) See Pascual (J.)	C6-719	Cazzanelli (E.), Fontana (A.), Mariotto (G.) and Rocca (F.) Vibrational dynamics of	
Campbell (D.A.).— See Black (J.E.)	C6-837	RbAg ₄ I ₅ by Raman spectros-	C6-190
Cancela (L.S.G.), Ramos (J.G.) and Gualberto (G.M.) Multi-			
phonon process in NiSO ₄ .6H ₂ O	C6-581	Cazzanelli (E.), Fontana (A.), Mariotto (G.), Rocca (F.) and	
Cantini (P.), Boato (G.) and Tatarek (R.).— Graphite surface phonons studied through He atoms resonant scattering	c6-801	Fontana (M.P.).— Observation of pretransitional effects on the shape of the $E_2(17~{\rm cm}^{-1})$ optical mode in β -AgI by	
Carabatos (C.) See Fontana		Raman. spectroscopy	C6-776
(M.D.)	C6-749	Cazzanelli (E.) See Mazacurati (V.)	C6-196
Carabatos (C.) See Kugel	06-904		
(G.E.)	C6-884	Celio (M.), Monnier (R.) and	
Cardona (M.) See Anastassakis (E.M.)	C6-537	Wachter (P.) Phonon and elastic anomalies in interme- diate valent Tm _x Se and	
Cardona (M.) See Calleia (J.M.)	C6-487	TmSe ₁ Te	C6-11

Cerdeira (F.), Melo (F.E.A.) and Katiyar (R.S.) Piezospectros copic study of the Raman spectrum of LiIO3	-	Colvard (C.), Merlin (R.), Klein (M.V.) and Gossard (A.C.) Light scattering from phonons: GaAs-AlAs superlattices	
Challis (L.J.), Ghazi (A.A.) and Wybourne (M.N.). Frequency crossing in bicrystals and inelastic scattering at surfaces	C6-229	Compaan (A.), Lo (H.W.), Lee (M.C and Aydinli (A.).— Pulsed Raman measurements of inhibited electron—phonon coupling at his plasma densities in silicon.	n
Challis (L.J.) See Ramdane (A.)	C6-244	Coombs (G.J.) See Lockwood (D.J.)	C6-761
Chao (M.H.) and Sladek (R.J.) Ultrasonic phonons in Hg _{0.8} Mn _{0.2} Te; dependences of elastic moduli on pressure and temperature	C6-667	Copley (J.R.D.), Bosio (L.), Cortes (R.), Lefebvre (J.) and Teuchert (W.D.) A neutron inelastic scattering study of phonons in metastable Beta- Galium	C6-386
Chaplot (S.L.), Mierzejewski (A.)		Cortes (R.) See Copley	
Lefebvre (J.), Pawley (G.S.) and Luty (T.) Internal and		(J.R.D.)	C6-386
external phonons in monoclinic		Couach (M.) See Khoder	
tetracyanoethylene	C6-584	(A.F.)	C6-383
Chaplot (S.L.) See Dorner (B.)	C6-602	Currat (R.) The role of phonons	
Chaplot (S.L.) See Rao (K.R.)	C6-637	in incommensurate phase transitions	C6-693
Cheeke (J.D.N.) and Madore (G.)			00 075
Acoustoelectric interaction in p-InSb at low temperature	C6-569	Czachor (A.) Gaps in phonon dispersion curves for substitutional alloys	C6-525
Cheeke (J.D.N.) See Poirier		Dacorogna (M.) See Ashkenazi	
(M.)	C6-876	(J.)	C6-355
Chernyshov (A.A.) See Dorner (B.),	C6-365	Dandoloff (R.) See Zeyer (R.)	C6-40
Chhi-Chong Wu Effect of the		Dattagupta (S.) See Sood	
propagation of ultrasounds on		(A.K.)	C6-572
magnetoacoustic phenomena in degenerate semiconductors	C6-655	Daudin (B.), Salce (B.) and Smith (S.H.) Specific heat	
Chih-Yuan Lu Ground State		and thermal conductivity of Tm VO ₄ and Dy VO ₄ with an	
energy of Frohlich electron- phonon system	C6-481	applied magnetic field	C6-277
phonon dyseemeet to the total total to the total total total to the total tota		Deb (S.K.) See Bansal (M.L.)	C6-902
Chin (M.A.) See Ulbrich		Dellepiane (G.) See Mulazzi	
(R.G.)	C6-226	(E.)	C6-493
Choi (B.K.) See Kim (J.J.)	C6-427	Delpech (P.) See Jusserand	C6-43
Ciplys (D.) and Prieur (J.Y.)			70 73
Ultrasonic properties of a superionic glass: B_2O_3 , 0.5 Li ₂ 0, 0.7 LiCl	C6-184	Dementjev (V.A.).— See Bokhenkov (E.L.)	C6-566
		Dementjev (V.A.) See Bokhenkov	
Coiret (M.) See Toledano (P.)	C6-476	(E.L.)	C6-605

Deville (A.), Gaillard (B.), Blanchard (C.) and Livage		Doussineau (P.) See Arnold (W.)	C6-37
(J.) Comparison of electron spin-lattice relaxation in amorphous and crystalline		Doussineau (P.) See Prieur (JY.)	C6-33
materials	C6-465	Drexhage (M.G.) See Banerjee	
Deville Cavellin (C.) See Petritis (D.)	C6-688	(P.K.)	C6-75
Devreese (J.T.) See Vanchamp		Duraiswamy (S.) and Haridasan (T.M.) Means square displa-	
(P.E.)	C6-625	cements of a substitutional defect and its nearest	
Dierker (S.B.) See Merlin (R.)	C6-392	neighbours in KI	C6-899
Dietsche (W.), Kirch (S.J.) and Wolfe (J.P.). Phonon spectros	-	Dynes (R.C.).— See Hensel (J.C.)	C6-212
copy of the electron-hole-	C6-447	Dynes (R.C.).— See Hensel (J.C.)	C6-308
Distants (U) - Cos Nowthwan		Ecolivet (C.) Linear chain models and sound velocities in	
Dietsche (W.) See Northrop (G.A.)	C6-209	molecular organic crystals	C6-578
Dletz (W.) See Fritsch (G.)	C6-143	Ecolivet (C.), Toudic (B.) and Sanquer (M.) Brillouin	
Dixon (G.S.) Two-Magnon-one- phonon scattering in the		scattering study of phase transitions in p-polyphenyls	C6-779
lattice thermal conductivity of magnetic insulators	C6-262	Einsenmenger (W.) Phonon	
		imaging	C6-201
Dixon (M.) See Elliott (R.J.)	C6-175	Elbert (M.) See Carlone (C.)	C6-323
Doak (R.B.) See Benedek (G.).	C6-793	Elliott (R.J.) and Dixon (M.)	
Dolganov (V.K.) See Bielushkin (A.V.)	C6-34	Vibrations and diffusion of atoms in superionic crystals and melts	C6-175
Dorner (B.), Bokhenkov (E.L.),			00-175
Sheka (E.F.), Chaplot (S.L.), Pawley (G.S.), Kalus (J.),		Engelhardt (M.). See Baumgartner (R.)	C6-119
Schmelzer (U.) and Natkaniec (I.) Phonon dispersion curve	S	Englman (R.) Dynamics in dis-	
in the molecular crystals naphtalene and anthracene		placive phase transitions	C6-755
measured by inelastic neutron scattering	C6-602	Englman (R.) Phonon pulses from	
	00 002	a relaxing system	C6-241
Dorner (B.), Boysen (H.), Frey (F.) and Grimm (H.).— On the		Errandonea (G.) Pressure depen-	
Si-O-Si bond angle in α - and β -quartz	C6-752	dence of the coupled optic	
Dorner (B.), Chernyshov (A.A.),		elastic transition of LaP ₅ 0 ₁₄	C6-710
Pushkarev (V.V.), Rumyantsev (A.Yu.) and Pynn (R.)			
Electron-phonon coupling in th nontransition metal cadmium.	e C6-365	Escribe-Filippini (C.) See Petitgrand (D.)	C6-782
Doussineau (P.) Attenuation and velocity change of acoustic		Etchepare (J.) See Merian (M.)	C6-608
waves in the amorphous metal		Every (A.G.) Phonon focusing and	
Pd Si Cu from 0.05 K to	06 70	the shape of the ray surface	06 215

Fain (B.) Instabilities in thermal bath: Phonon enhance- ment and saturation in a three-	-	Franchini (A.).— See Bortolani (V.)	C6-831
1 1	C6-540	Freire (J.D.) See Katiyar	
Falter (C.), Ludwig (W.) and Selmk	te	(R.S.)	C6-433
(M.) Renormalized response theory with applications to		Frenois (C.) See Arnold (W.).	C6-37
phonon anomalies	C6-516	Frenois (C.) See Prieur (JY.)	C6-33
Fasolino (A.), Santoro (G.) and Tosatti (E.) Effective mass		Frey (F.) See Dorner (B.)	C6-752
approximation for surface phonon states	C6-840	Frisken (B.), Guillon (F.), Harrison (J.P.) and Page (J.H.) Low frequency phonons	
Fasolino (A.), Santoro (G.) and Tosatti (E.) Surface Phonons		in sintered copper	C6-858
and the Incommensurate reconstruction of clean Mo(100)	C6-846	Fritsch (G.), Dietz (W.), Häring (R.) and Lüscher (E.)	
Fay (D.) and Appel (J.) Effect of phonons on the spin susceptibility	C6-490	Anharmonic contributions to the thermal diffuse X-Ray scatte- ring intensity from single	
		crystalline sodium.,,,	C6-143
Fishman (G.), Pinczuk (A.), Worloc (J.M.), Störmer (H.L.), Gossard		Furdin (G.) See Batallan (F.)	C6-344
(A.C.) and Wiegmann (W.) Coupled electron-LO phonon		Gaillard (B.) See Deville (A.)	C6-465
excitation in a two dimensional electron system	C6-305	Galeener (F.L.) Vibrational selection rules in disordered	
Fisk (Z.) See Merlin (R.)	C6-392	solids : vitreous GeO ₂	C6-24
Fontana (A.) See Cazzanelli	C6-19 0	Geick (R.), Rauch (H.), Lehner (N.), Bouillot (J.), Stirling (W.G.), Heger (G.) Lattice	
Fontana (A.) See Cazzanelli (E.)	C6-776	instabilities and phase transi- tions in fluoride perovskites	
Fontana (A.) See Mazzacurati	C6-196	Gerlinger (H.) and Schaack (G.) Magnetic field dependent Reststrahlen spectra in parama-	
Fontana (M.D.), Kugel (G.E.) and Carabatos (C.) Lattice		gnetic CeF ₃ ······	C6-499
dynamics of the cubic-tetra- gonal phase transition in KNbO ₃	C6-749	Gervais (F.) and Servoin (J.L.) Role of polar phonons in the chemical bound at structural phase transitions characterized	
Fontana (M.P.) See Maisano	C6-51	by repetitive Fourier spectros-	
Fontana (M.P.) See Aliotta	C6-57	Ghazi (A.A.) See Challis (L.J.)	C6-229
Fontana (M.P.) See Cazzanelli	C6-776	Gil (B.) See Camassel (J.)	C6-682
Fossheim (K.) See Fossum (J.O.)	C6-867	Gledhill (G.A.), Kudhail (S.S.),	
Fossum (J.O.), Aune (H.J.), Fossheim (K.) and Holt (R.M.) Echo properties of BGO and		Newman (R.C.), Woodhead (J.) and Zhang (G.Z.). The low frequency absorption of gallium phosphide induced by impurities	C6-685

De Goër (A.M.), Locatelli (M.) and Laermans (C.).— Thermal phonon transport study of the defects created in quartz single crystals by different irradia— tions (γ rays, electrons, neutrons)	Greenstein (M.) and Wolfe (J.P.) Modulation of ballis- tic phonon fluxes in Ge by a He Gas film at a solid/super- fluid interface
De Goër (A.M.), Locatelli (M.) and Lassmann (K.) A Study of the	Greig (D.) and Hardy (N.D.) The thermal conductivity of semicrystalline polymers at very low temperatures
ground state of acceptors in silicon from thermal transport experiments	Grier (B.H.) See Sato (M.) C6-770
De Goër (A.M.), Locatelli (M.) and	Grimm (H.) See Dorner (B.) C6-752
Nicolau (I.F.) Low tempera- ture thermal conductivity of Hg I ₂ single crystals C6-265	Gualberto (G.M.) See Cancela (L.S.G.)
De Goër (A.M.) See Salce (B.) C6-424	Guillon (F.) See Frisken (B.) C6-858
Goltman (J.T.) See Banerjee	Güntherodt (G.) See Kress (W.) C6-3
(P.K.)	Gupta (0.P.) Phonon dispersion and thermophysical properties
Gompf (F.) and Lannin (J.S.) The vibrational spectra of	of sodium
crystalline and amorphous phosphorous	Hamaguchi (C.) See Itoh (Y.). C6-810
Gompf (F.), Schmidt (H.J.) and Renker (B.) Inelastic neutron	Hamann (D.R.) See Harmon (B.N.)
scattering from amorphous Fe40Zr60 doped with different concentrations of Krypton	Hamdache (M.), King (P.J.), Murphy (D.T.) and Rampton
(0,1, 7 at .%)	(V.W.) Phonon spectroscopy of chromium-doped gallium arsenide using superconducting
Goodstein (D.) See Taborek (P.) C6-825	tunnel junctions
Goodstein (D.) See Taborek (P.) C6-855	Hanke (W.) See Muramatsu (A.) C6-834
Goodstein (D.) See Taborek (P.) C6-852 Gorelik (V.S.), Umarov (B.S.) and	Hardcastle (J.F.) and Zabel (H.) Thermal expansion of Alkali-
Umarov (M.) "Non-Central" peaks in light scattering spec-	graphite intercalation compounds
tra of lithium tantalate and quartz at phase transition. • C6-764	Hardy (N.D.) See Greig (D.) C6-69
Gorelik (V.S.) See Anikiev (A.A.)	Haridasan (T.M.) See Balasu- bramanian (K.R.) C6-905
Gossard (A.C.) See Colvard (C.) C6-631	Haridasan (T.M.) See Duraiswamy (S.)
Gossard (A.C.) See Fishman (G.) C6-305	Haridasan (T.M.) See Jacobs (P.W.M.)
Govindarajan (J.) See Jacobs (P.W.M.)	Haridasan (T.M.) See Ramachandran (K.)
Granato (A.V.) See Kneezel	Häring (R.) See Fritsch (C.). C6-1/3
1 Maria Janaanaanaanaanaanaanaanaanaanaan 176-750	HOLLIE UNA JAT DEE FEITSCH HELL. Challes

Harmon (B.N.), Weber (W.) and Hamann (D.R.) Vibrational frequencies via frozen phonons	C6-628	Horie (C.), Miyazaki (H.), Igarashi (S.) and Hatakeyama (S.) Role of phonons in orientational intra-plane ordering in graphite intercalation	
Harrison (J.P.) See Frisken (B.)	C6-858	compounds	C6-298
Hashimoto (S.) See Kuroda (N.)	C6-788	Horie (C.) See Miyazaki (H.). Horstman (R.E.) and Wolter (J.)	C6-335
Hatakeyama (S.) See Horie	C6-298	Thermalization of high- frequency phonons in silicon single crystals	C6-813
Hattori (T.), Imanishi (T.), Kurokawa (H.) and Mitsuishi		Hota (N.K.) See Carlone (C.).	C6-323
(A.) Brillouin scattering in silver halides	C6-920	Hu (P.) See Lax (M.)	C6-161
Hayes (W.) Phonons in super- ionic conductors	C6-167	Hwang (D.M.) and Solin (S.A.) Phonon Raman spectroscopy in graphite intercalation	
Heger (G.) See Geick (R.)	C6-713	compounds	C6-347
Henkel (J.H.) Lattice phonon		Igarashi (S.) See Horie (C.)	C6-298
lifetime calculation	C6-146	Imanishi (T.) See Hattori (T.)	C6-920
Hennion (B.).— See Petitgrand (D.)	C6-782	Inoue (K.) and Asai (N.) Hyper- Raman scattering study of ferro electric phase transitions in)
Hensel (J.C.), Dynes (R.C.) and Tsui (D.C.) Absorption of		Sr Ti O ₃ and Ba Ti O ₃	C6-430
ballistic phonons by the 2D electron gas in a Si mosfet.	C6-308	Inoue (M.) See Alig (R.C.)	C6-484
Hensel (J.C.), Dynes (R.C.), Unterwald (F.C.) and Simons		Inuishi (Y.) See Itoh (Y.)	C6-810
(A.L.) Singular effects in phonon focusing; formation of		Itoh (Y.), Hamaguchi (C.) and Inuishi (Y.) Resonant	
phonon focusing coustics	C6-212	Brillouin scattering in opaque region of CdS	C6-810
Heritier (M.) The phonon anomal	у	Itoh (S.) See Nakamura (T.)	C6-418
in b.c.c. He ⁴ : a phonon self- trapping ?	C6-890	Iwasa (J.) and Suzuki (H.)	
Heritier (M.) and Montambaux	00 070	Interaction of sound with vacancies in bcc ³ He crystals	C6-896
(G.) Interpretation of		Jacobs (P.W.W.), Nerenberg	
experiments in the solidon model for rotons	C6-534	(M.A.H.), Govindarajan (J.) and Haridasan (T.M.) Calcu-	
Hibma (T.) See Brüesch (P.)	C6-178	lation of defect entropies in ionic crystals in the	
Hidaka (Y.) See Yagi (T.)	C6-731	quasiharmonic approximation : application to fluorites	C6-914
		Jang (M.S.) See Nakamura (T.)	C6-418
Hikata (A.) See Poirier (M.).	C6-876	Janssen (T.) and de Lange (C.)	
Hochheimer (H.D.) See Carlone		Vibration spectra of crystals	04 242
(C.)	C6-323	in an incommensurate phase	C6-737
Holt (R.M.) See Fossum (J.O.)	C6-867	Jayaraman (A.) See Kress (W.)	C6-3

Joffrin (C.) See Schneck (J.)	C6-734	Keller (J.) See Bavernfeind	
Johnstone (I.W.), Jones (G.D.)		(W·)······	C6-247
and Lockwood (D.J.) Raman scattering from phonons in disorded CsMg _{1-x} Co _x		Kesavamoorthy (R.) See Umadevi (V.)	C6-81
Cl ₃ ····································	C6-90	Kesavasamy (K.). See Krishnamurthy (N.)	C6-908
Jones (G.D.) See Johnstone (I.W.)	C6-90	Khare (A.) See Behera (S.N.).	C6-314
Jørgensen (J.) See Merlin (R.)	C6-392	Khoder (A.F.), Couach (M.),	
Joukoff (B.) See Schneck (J.). Jusserand (B.), Sapriel (J.),	C6-734	Locatelli (M.), Abou-Ghantous (M.) and Senateur (J.P.) On the thermal and electrical	
Alexandre (F.) and Delpech (P.) Raman study of the anharmonicity and disorder-		conductivity of V ₃ Si and V ₅ Si ₃	C6-383
induced effects in Ga _{1-x} Al _x As	C6-43	Kim (J.J.), Won (J.W.) and Choi (B.K.) Soft-mode spectra and phase transition in KDP	
Kajimura (K.) Phonon echoes in bulk and powdered		crystal with ADP impurities.	C6-427
materials	C6-863	Kimura (T.) See Yoshikawa	C6-873
Kajimura (K.) See Tsuruoka	06_06/	(S.)	
	C6-864	Kinder (H.) See Berberich (F.)	C6-374
Kalus (J.) See Dorner (B.)	C6+602		
Kamitakahara (W.A.), Wada (N.) and .Solin (S.A.). Neutron spectros copy of phonons in Rb C ₈		King (P.J.) The effect of lattice strains on the acoustic relaxation losses in dielectric crystals	
Kanellis (G.), Morhange (J.F.) and Balkanski (M.) Electro-	đ	due to magnetic ions King (P.J.) See Hamdache (M.)	C6-456
static energy and lattice vibration in thin ionic	or 0/1		
slabs	C6-341	Kirch (S.J.) See Dietsche (W.)	C6-447
Kaplyanskii (A.A.), Basoon (S.A.) and Shekhtman (V.L.) Multiple resonant Raman spin-		Klemens (P.G.) Theory of the attenuation of elastic waves in	n
flip phonon scattering and 29 cm ⁻¹ phonon transport in excited ruby in a magnetic			C6-102
field	C6-462	Orta (R.)	C6-256
Kaplyanskii (A.A.), Basoon (S.A.) and Shekhtman (V.L.)		Vlote (W V)	
Phonons in excited ruby	C6-439	Klein (M.V.) Raman scattering from anomalous phonons in transition metals and	
Katayama (S.) See Suski (T.).	C6-758	compounds	C6-368
Katiyar (R.S.) and Freire (J.D.) Dynamical study of		Klein (M.V.) See Colvard (C.)	C6-631
phonons in ferrolectric lead titanate	C6-433	Klein (M.V.) See Merlin (R.).	C6-392
Katiyar (R.S.) See Cerdeira		Klein (M.V.) See Wada (N.)	C6-350
(F.)	C6-722	Klein (P.B.) See Strom (U.)	C6-30

(litsner (T.), Raychaudhuri (A.K.) and Pohl (R.O.) Connection between the low		<pre>Kugel (G.E.) See Fontana (M.D.) Kunc (K.) and Martin (R.M.)</pre>	C6-749
temperature thermal proper- ties of glasses and their glass transition temperature	C6+66	Ab initio force constants of germanium	C6-649
(neezel (G.A.) and Granato		Kunc (K.) See Pollian (A.)	C6-295
(A.V.) Phonon-Dislocation dipole interaction in Li F at low temperature	C6-250	Kuroda (N.), Hashimoto (S.) and Nishina (Y.) Spectroscopic ionicity and lattice instabi- lity in III-IV layer	
<pre>Cohli (M.) See Prohofsky (E.W.)</pre>	C6-560	compounds	C6-788
Kojima (S.) and Nakamura (T.) Raman scattering study of low frequency phonon polari-		Kuroda (N.) and Nishina (Y.) Two-exciton process resonant Raman scattering in InSe	C6-332
tons in several crystals	C6-767	Kurokawa (H.) See Hattori (T.)	C6-920
Kojima (S.) See Nakamura (T.)	C6-418	Kuzmany (H.) See Carlone (C.)	C6-323
Kolesníkov (A.I.) See Bokhenkov (E.L.)	C6-566	Laermans (C.) See De Goër (A.M.)	C6-78
(olesnikov (A.I.) See Bokhenkov (E.L.)	C6-605	Lagnier (R.) See Bonjour (E.)	C6-63
(ress (W.), Bilz (H.),Guntherodt (G.) and Jayaraman (A.) Phonons in mixed valence		Landwehr (G.) See Brunel (L.C.)	C6-412
compounds	C6-3	de Lange (C.) See Janssen (T.)	C6-737
Kress (W.) See Kugel (G.E.)	C6-884	Lannin (J.S.) See Carroll (P.J.)	C6-643
<pre>(rishnamurthy (N.) and Kesavasamy (K.) Phonon dispersions in calcium tungstate</pre>	C6-908	Lannin (J.S.) See Gompf (F.).	C6-27
<pre>Krivenko (T.A.) See Bokhenkov (E.L.)</pre>	C6-566	Lannin (J.S.) See Kshirsagar (S.T.)	C6-54
Krivenko (T.A.) See Bokhenkov (E.L.)	C6-605	Lasjaunias (J.C.) See Bethoux (0.)	C6-60
Kshirsagar (S.T.) and Lannin (J.J.) Raman scattering in		Lassmann (K.) See De Goër (A.M.)	C6-235
anneal stable amorphous silicon	C6-54	Lauger (K.) See Ströbel (B.).	C6-404
Kuchta (B.) and Luty (T.) Harmonic and anharmonic repre-		Lauter (H.J.) See Batallan (F.)	C6-344
sentation of librational motions in molecular crystals	C6-587	Lax (M.), Narayanamurti (V.), . Hu (P.) and Weber (W.) Lifetimes of high frequency	
Kudhail (S.S.) See Gledhill (G.A.)	C6-685	phonons	C6-161 C6-453
Kugel (G.E.), Carabatos (C.) and		Lefebvre (J.) See Compaan (A.)	00-433
Kress (W.) Lattice dynamics of cuprite (Cu ₂ 0)	C6-884	(S.L.)	C6-584

Legrand (S.) See Copley (J.R.D.)	C6-386	v.Löhneysen (H.), Schink (H.J.), Arnold (W.), Beyeler (H.U.), Pietronero (L.), Strässler (S.) Low-temperature spe-	
(D.)	C6-782	cific heat anomaly of the one-dimensional ionic con-	
Lehner (N.) See Geick (R.)	C6-713	ductor hollandite	C6-193
Leisure (R.G.) See Prieur (JY.)	C6-33	Lu (K.C.) See Prohofsky	C6-560
Lengfellner (H.) and Renk (K.F.) Detection of acoustic		Ludwig (W.) See Falter (C.).	C6-516
zone-boundary phonons by phonon difference absorption	C6-259	Luty (T.) Translational-	C6-143
Leung (W.N.) See Ball (M.A.).	C6-911	rotational coupling in s- triazine crystal	C6-590
Levelut (A.) See Arnold (W.).	C6-37	Luty (T.) See Chaplot (S.L.)	C6-584
Levelut (A.) See Prieur (JY.)	C6-33	Luty (T.) See Kuchta (B.)	C6-587
Lipson (H.G.) See Bendow (B.)	C6-140	Maciel (A.) and Ryan (J.F.) Lock-in type phase transition	06 716
Livage (J.) See Deville (A.).	C6-465	in KFeF ₄	C0-/10
Livescu (G.) See Brafman (O.)	C6-96	Maciel (A.) and Ryan (J.F.) Observation of coupled ampli-	
Lo (H.W.) See Compaan (A.)	C6-453	tude modes in the Raman spec- trum of incommensurate	0/ 705
Locatelli (M.), Suchail (R.) and .Zecchi (E.).— A study of the 'Pb precipitation in NaCl from thermal conductivity		γ-Na ₂ Co ₃	C6-723
experiments	C6-893	Magerl (A.) and Zabel (H.) Lattice dynamics of graphite	
Locatelli (M.) See de Goër (A.M.)	C6-78	intercalation compound-mo- delled by the phonon dis-	
Locatelli (M.) See de Goër (A.M.)	C6-235	persion of linear chains	C6-329
Locatelli (M.) See de Goër	00 233	Magerl (A.) See Zabel (H.)	C6-289
(A.M.)	C6-265	Mahan (G.D.) See Tua (P.F.).	C6-122
Locatelli (M.) See Khoder (A.F.)	· C6-383	Maier (J.) and Sigmund (E.).— The influence of the dynamical Jahn—Teller effect of	
Locatelli (M.) See Ramdane (A.)	C6-244	acceptors onto the phonon- transport mechanism in cubic semiconductors	C6-232
Lockwood (D.J.) and Coombs (G.J.) A Raman study of the structural phase transition		Maisano (G.), Migliardo (P.) Wanderlingh (F.), Fontana	
in disordered KMn _{0.85} Ni _{0.15} F ₃	C6-761	(M.P.), Bellissent-Funel (MC.) and Roth (M.)	
Lockwood (D.J.) - See Johnstone (I.W.)	C6-90	Local order and dynamics in liquid electrolytes : small angle neutron scattering	C6-51
Lowenhaupt (M.) See Yeh (C.T.)	C6-371	Maisano (G.) See Aliotta (F.)	C6-57

Yaradudin (A.A.), Martin (A.J.), Bilz (H.) and Wallis (R.F.) Non-linear dynamical excitation Maradudin (A.A.) See Shen	C6-137	Mazzacurati (V.), Ruocco (G.), Signorelli (G.), Cazzanelli (E.), Fontana (A.) and Mariotto (G.) A theoretical model for the continuous order-disorder transition at	
(T.J.)	C6-819	703 K in superionic α-AgI	C6-196
Maradudin (A.A.) See Suzuki (K.)	C6-640	Mei (M.N.) See Prohofsky (E.W.)	C6-560
Mariotto (G.) See Cazzanelli (E.)	C6-776	Melo (F.E.A.) See Cerdeira (F.)	C6-722
Mariotto (G.) See Cazzanelli (E.)	C6-190	Meltzer (R.S.) See Sox (D.J.)	C6-450
Mariotto (G.) See Mazzacurati (V.)	C6-196	Menn (K.) and Biem (W.) Coherent potential approximation and strongly anharmonic systems: crystals of	
Maroie (S.) See Thiry (P.) Marshall (R.C.) See Bendow	C6-843	para-hydrogen and neon resp. hydrogen atoms	C6-87
(B.)	C6-140	Menon (M.) See Arnold (G.B.).	C6-377
Martin (A.J.) See Maradudin (A.A.) Martin (D.J.)and Walker (R.P.)	C6-137	Meredith (D.J.), Pritchard (J.A.) and Wigmore (J.K.) Backward wave phonon echoes at 17 Ghz in single crystals of LiTaO ₃ , LiNbO ₃ and	
How much information can low energy ion scattering give about surface phonons ? Martin (R.M.) Recent advances	C6-807	Bi ₁₂ SiO ₂₀	C6-870
in the theory of phonons in semiconductors	C6-617	with the help of a single vibrational potential	C6-608
Martin (R.M.) See Kunc (K.)	C6-649	Merlin (R.) See Colvard (C.).	C6-631
Martinez (G.) See Petritis (D.)	C6-688	Merle (P.) See Camassel (J.). Merle (P.) See Pascual (J.)	C6-682 C6-719
Marvin (A.) See Bortolani (V.) Massa (N.E.) Raman scattering	C6-804	Merlin (R.), Dierker (S.B.), Klein (M.V.), Jørgensen (J.), Rasmussen (S.R.), Fisk (Z.) and Webb (G.W.) Raman	
spectra of Rb ₂ ZnCl ₄ under uniaxial stress	C6-593	scattering in V ₃ Si, V ₃ Ge, Nb ₃ Sb and Cr ₃ Si : correlation of E _g optical phonon line-	
Massot (M.) See Brunel (L.C.)	C6-412	width with magnetic suscep- tibility	C6-392
Mathieu (H.) See Camassel	C6-682	Michard (F.) See Perrin (B.).	C6-131
Mathieu (H.) See Pascual	0/ 710	Mierzejewsi (A.) See Chaplot (S.L.)	C6-584
(J.)	C6-719	Migliardo (P.) See Aliotta	C6-57
Maurer (D.) See Miller (V.).	C6-389	(F.)	00-37

Migliardo (P.) See Maisano (G.)	C6-51	Moussa (F.) See Cailleau (H.) Moynihan (C.T.) See Banerjee	C6-704
Mitra (S.S.) See Banerjee (P.K.)	C6-75	(P.K.)	C6-75
Mitra (S.S.) See Bendow (B.).	C6-140	Mulazzi (E.), Tubino (R.) and Dellepiane (G.).— On the origin of the red shift of	
Mitsuishi (A.) See Hattori	C6-920	Raman excitation profile from the absorption spectrum	C6-493
Miura (K.) See Yagi (T.)	C6-731	Mulazzi (E.) See Barchielli (A.)	C6-475
Miyajima (H.) See Sato (M.)	C6-770	Millor (V) Cohong (C)	
Miyasato (T.), Tokumura (M.) and Akao (F.). Effect of magnetic field on ballistic heatpulse scattering in n-type germanium	C6-658	Müller (V.), Schanz (G.), Unterhorst (E.J.) and Maurer (D.) Nuclear acoustic resonnance investigations of the longitudinal and transverse electron lattice interaction in transition	
Miyazaki (H.) and Horie (C.)		metals and alloys	C6-389
Effect of electron-phonon interaction on Raman spectra of graphite intercalation		Müller (V.) See Ströbel (B.).	C6-404
compounds	C6-335	Muramatsu (A.) and Hanke (W.) Microscopic theory of lattice	
Miyazaki (H.) See Horie (C.).	C6-298	dynamics and reconstruction of semiconductor surfaces	C6-834
Mokross (B.) See Bussmann- Holder (A.)	C6-409	Murase (K.) See Sugai (S.)	C6-740
Monnier (R.) See Celio (M.)	C6-11		
		Muraco (V) - Coo Cucoi (C)	06 220
Montambaux (G.) See Heritier	06 52/	Murase (K.) See Sugai (S.)	C6-320
Montambaux (G.) See Heritier (M.)	C6-534	Murase (K.) See Sugai (S.) Murase (K.) See Sugai (S.)	C6-320 C6-728
	C6-534		
(M.) Moore (D.R.), Tekippe (V.J.), Ramdas (A.K.) and Toledano	C6-534	Murase (K.) See Sugai (S.) Murphy (D.T.) See Hamdache (M.)	C6-728
(M.) Moore (D.R.), Tekippe (V.J.), Ramdas (A.K.) and Toledano (J.C.).— A Raman scattering study of the solid to	C6-534 C6-785	Murase (K.).— See Sugai (S.) Murphy (D.T.).— See Hamdache (M.)	C6-728
(M.) Moore (D.R.), Tekippe (V.J.), Ramdas (A.K.) and Toledano (J.C.).— A Raman scattering study of the solid to solid phase transition in		Murase (K.) See Sugai (S.) Murphy (D.T.) See Hamdache (M.) Mutsheller (W.) See Wagner (M.) Nakamura (T.), Kojima (S.), Jang (M.S.), Takashige (M.) and Itoh (S.) E(x)-A ₁ (z)	C6-728
(M.) Moore (D.R.), Tekippe (V.J.), Ramdas (A.K.) and Toledano (J.C.).— A Raman scattering study of the solid to solid phase transition in benzil Morhange (J.F.).— See Kanellis	C6-785	Murase (K.).— See Sugai (S.) Murphy (D.T.).— See Hamdache (M.) Mutsheller (W.).— See Wagner (M.) Nakamura (T.), Kojima (S.), Jang (M.S.), Takashige (M.)	C6-728
Moore (D.R.), Tekippe (V.J.), Ramdas (A.K.) and Toledano (J.C.).— A Raman scattering study of the solid to solid phase transition in benzil	C6-785	Murase (K.).— See Sugai (S.) Murphy (D.T.).— See Hamdache (M.)	C6-728 C6-664 C6-238
<pre>Moore (D.R.), Tekippe (V.J.), Ramdas (A.K.) and Toledano (J.C.) A Raman scattering study of the solid to solid phase transition in benzil Morhange (J.F.) See Kanellis (G.)</pre>	C6-785 C6-341 C6-876 C6-14	Murase (K.).— See Sugai (S.) Murphy (D.T.).— See Hamdache (M.) Mutsheller (W.).— See Wagner (M.) Nakamura (T.), Kojima (S.), Jang (M.S.), Takashige (M.) and Itoh (S.).— E(x)—A1(z) oblique phonons in tetragonal BaTiO3 Nakamura (T.) Takashige (M.) and Aikawa (Y.).— Raman	C6-728 C6-664 C6-238
Moore (D.R.), Tekippe (V.J.), Ramdas (A.K.) and Toledano (J.C.).— A Raman scattering study of the solid to solid phase transition in benzil	C6-785 C6-341 C6-876 C6-14	Murase (K.).— See Sugai (S.) Murphy (D.T.).— See Hamdache (M.)	C6-728 C6-664 C6-238
Moore (D.R.), Tekippe (V.J.), Ramdas (A.K.) and Toledano (J.C.).— A Raman scattering study of the solid to solid phase transition in benzil	C6-785 C6-341 C6-876 C6-14	Murase (K.).— See Sugai (S.) Murphy (D.T.).— See Hamdache (M.)	C6-728 C6-664 C6-238 C6-418 C6-421 C6-767
Moore (D.R.), Tekippe (V.J.), Ramdas (A.K.) and Toledano (J.C.).— A Raman scattering study of the solid to solid phase transition in benzil	C6-785 C6-341 C6-876 C6-14	Murase (K.).— See Sugai (S.) Murphy (D.T.).— See Hamdache (M.)	C6-728 C6-664 C6-238 C6-418

Varayanamurti (V.) See Ulbrich		Oka (Y.) and Cardona (M.)	
(R.G.)	C6-226	Raman scattering study of correlation between phonons	
Watkaniec (I.) See Bielushkin		and electron spins in CdS	
	C6-34	and ZnTe under resonance condition	C6-459
Natkaniec (I.). See Bokhenkov	06 60 5	01 (2) 0 0 0 0	
(E.L.)	C6-605	Olego (D.), Cardona (M.) and Vogl (P.) Volume and	
Natkaniec (I.) See Dorner (B.)	C6-602	temperature dependence of the transverse charge and the	
Natkaniec (I.) See Wasiutynski	C6-599	ionicity of tetrahedral semiconductors	C6-652
Nava (R.) and Rodriguez (M.)		Onwuagba (B.N.) and Animalu	
Frequency dependence of acoust saturation in smoky quartz		(A.O.E.) Screening of the short-range potential by the local field correction in	
Nemanich (R.J.) and Tsai (C.C.) Phonons of the metal/amorphous		the lattice dynamics of vanadium	C6-401
silicon interface studied by interference enhanced Raman		Padmanahhan (V W) - Can B	
scattering	C6-822	Padmanabhan (V.M.).— See Rao (K.R.)	C6-637
Nerenberg (M.A.H.).— See Jacobs (P.W.M.)	C6-914	Page (J.H.) See Frisken (B.)	C6-858
	00 714	Page (J.B.) See Schröder	
Newman (R.C.) See Gledhill	06 605	(U.)	C6-496
(G.A.)	C6-685	Parrinello (M.) and Rahman (A.) Polymorphic transi-	
Nicolau (I.F.) See de Goër		tions in alkali halides. A	
(A.M.)	C6-265	molecular dynamics study	C6-511
Nielsen (0.H.) Vibrations of		Pascoli (G.) See Toledano	
substitutional impurities in semiconductors	c6-673	(P.)	C6-746
Nishina (Y.). See Kuroda (N.)	C6-332	Pascual (J.), Camassel (J.), Merle (P.) and Mathieu	
Nishina (Y.) See Kuroda (N.)	C6-788	(H.) Gruneisen parameter and deformation potentials of	
Nishina (Y.) See Sasaki (Y.)	C6-557	Raman active modes in rutile structure compounds	C6-719
Nizzoli (F.) See Bortolani	-1 -001	Description Constraint (I)	06-692
(V.)	C6-804	Pascual (J.) See Camassel (J.)	C6-682
Nizzoli (F.) See Bortolani (V.)	C6-831	Patel (M.B.) and Bist (H.D.) Librational modes of water in crystal hydrates and phase-	
Northrop (G.A.), Dietsche (W.)		transitions	C6-917
Zdetsis (A.D.) and Wolfe (J.P.). Focussing of disper-		Pawley (G.S.) See Chaplot (S.L.)	C6-584
sive phonons in Ge	C6-209	(15020)	
	•	Pauley (G.S.) See Dorner (B.)	C6-602
Nücker (N.) See Yeh (C.T.)	C6-371	Pelous (J.) See Vacher (R.)	C6-125
Nunez-Regueiro (M.) See	C6-229	Perrin (B.) and Michard (F.)	
Ayache (C.)	. C6−338	Phonon-phonon interactions in molecular crystals studied by	
Nusser (HK.) - See Wagner (M.)	C6-238	ultrasonic methods	C6-131

Perrin (B.) See Boissier (M.).	C6-611	Prieur (J.Y.) See Ciplys (D.)	C6-184
Petitgrand (D.), Hennion (B.), Escribe-Filippini (C.) and Legrand (S.) Phonon disper-		Pritchard (J.A.) See Meredith (D.J.)	C6-870
sion and transverse mode sof- tening in RbFeCl ₃	C6-782	Prohofsky (E.W.), Van Zandt (L.L.), Kohli (M.), Lu	
Petritis (D.), Deville Cavellin (C.) and Martinez (G.) Resonant enhancement of the Raman linewidth of phonon		(K.C.), Mei (M.N.) and Putman (B.). Phonons in the DNA double helix	C6-560
modes induced by hydrostatic pressure	C6-688	Pushkarev (V.V.) See Dorner (B.)	C6-365
Philip (J.) See Breazeale		Putman (B.) See Prohofsky	
(M.A.)	C6-134	(E.W.)	C6-560
Pick (R.M.) See Poulet (H.)	C6-701	Pynn (R.) See Dorner (B.)	C6-365
Pietronero (L.) See v.Löhneysen (H.)	C6-193	Radhakrishnan (V.) See Sharma (P.C.)	C6-268
Pinczuk (A.) See Fishman (G.)	C6-305	Rahman (A.) See Parrinello	
Pintschovius (L.), Vergés (J.A.)		(M.)	C6-511
and Cardona (M.) Self- energy of phonons interacting with free carriers in		Ramachandran (K.) and Haridasan (T.M.) The localized modes	
silicon	C6-634	due to phospherous defects in cadmium telluride	C6-679
Pireaux (J.J.) See Thiry (P.)	C6-843	Ramdane (A.), Salce (B.),	
Pohl (R.O.) See Klitsner (T.)	C6-66	Challis (L.J.) and Locatelli (M.) Phonon scattering by	
Poirier (M.), Cheeke (J.D.N.), Morisseau (P.) and Hikata (A.) Attenuation of a		Cr ions in GaAs and the effect of uniaxial stress	C6-244
Static polarization echo by He film	C6-876	Ramdas (A.K.) and Rodriguez (S.) Time reversal symmetry and the Raman scattering	
Pollian (A.), Kunc (K.) and Rossat-Mignod (J.) Inelas-		by crystals	C6-887
tic neutron scattering and lattice dynamics of the		Ramdas (A.K.).— See Moore (D.R.)	C6-785
layer semiconductor GaS	C6-295		
Pollini (I.) See Benedek (G.)	C6-301	Ramos (J.G.) See Cancella (L.S.G.)	C6-581
Poulet (H.) and Pick (R.M.)		Rampton (V.W.) See Hamdache	
Are phasons in biphenyl detectable by light		(M.)	C6-664
scattering ?	C6-701	Rao (K.R.), Padmanabhan (V.M.), Vijayaraghavan (P.R.) and	
Prasad (P.N.) Dynamics of phonon interactions inorganic		Chaplot (S.L.) Neutron, X-ray and lattice dynamical	
solids	C6-563	studies of Sb ₂ S ₃	C6-637
Prieur (JY.), Doussineau (P.), Frenois (C.), Levelut (A.)		Rasmussen (S.R.) See Merlin (R.)	C6-392
and Leisure (R.G.) Low- temperature acoustical pro-		Rauh (H.) See Geick (R.)	C6-713
perties of Na doped		(m) bee deter (m) ·····	00 713
β-A1 ₂ 0 ₃	C6-33	Ravex (A.) See Bethoux (O.)	C6-60

Raychaudhuri (A.K.) See Klitsner (T.)	C6-66	Rossat-Mignod (J.) See Pollian (A.)	C6-295
Rayne (J.A.) See Mountfield (K.R.)	C6-468	Roth (M.) See Maisano (G.)	C6-51
Rebane (L.A.) and Rebane (K.K.)		Roy (A.P.) See Bansal (M.L.).	C6-902
Phonons in resonant secondary emission of impurity molecules		Roy (K.P.) See Sharma (P.C.).	C6-268
in crystals	C6-505	Rumyantsev (A.Yu.) See Dorner (B.)	C6-365
Rebane (K.K.) See Rebane (L.A.)	C6-505	Ruocco (G.) See Mazzacurati	
Rebane (K.) See Saari (P.)	C6-502	(V.)	C6-196
Reichardt (W.) See Yeh (C.T.)	C6-371	Ryan (J.F.) See Maciel (A.)	C6-716
Reiger (R.) See Schröder (U.)	C6-496	Ryan (J.F.) See Maciel (A.)	C6-725
Renk (K.F.) Voir Baumgartner	C6-119	Rynee (T.M.) See Spector (H.N.)	C6-670
Renk (K.F.) See Lengfellner	30 117	Saari (P.) and Rebane (K.) Phonon relaxation in crystals	
(H.)	C6-259	studied by hot luminescence.	C6-502
Renker (B.) See Gompf (F.)	C6-49	Sahni (V.C.) On the possible use of optical rotation mea-	
Renker (B.) See Yeh (C.T.)	C6-371	surements for detecting weakly IR active modes	C6-881
Renucci (J.B.) See Carles (R.)	C6-105	Sahni (V.C.) See Bansal (M.L.)	C6-902
		Saint-Cricq (N.) See Carles	
Renucci (M.A.) See Carles		(R.)	C6-105
(R.)	C6-105	Salce (B.), De Goër (A.M.) and	
Resta (R.) and Baldereschi		Boatner (L.A.) Study of "pure" KTaO ₃ and KTN	
(A.) Local-field effects and zone-center phonons in		single crystals by thermal	
polar and covalent cubic	0/ //1	conductivity measurement	C6-424
semiconductors	C6-661	down to 50 mK	00-424
Rives (J.E.) See Sox (D.J.)	C6-450	Salce (B.) See Bonjour (E.)	C6-63
Rocca (F.) See Cazzanelli (E.)	C6-190	Salce (B.) See Daudin (B.)	C6-277
Rocca (F.) See Cazzanelli (E.)	C6-776	Salce (B.) See Ramdane (A.)	C6-244
Rodriguez (M.) See Nava (R.).	C6-99	Samsur (Sk.) See Behera	C6-528
Rodriguez (S.) See Ramdas			
(A.K.)	C6-887	Sanquer (M.) See Ecolivet (C.)	C6-779
Rogers (S.J.), Shaw (C.J.) and Wiederick (H.D.) Limita-		Santoro (G.):- See Bortolani (V.)	C6-804
tions on the usefulness of		Santoro (G.) See Bortolani	
metallic thin film semi- conductors for phonon detec-		(V.)	C6-831
tion	C6-317	Santoro (G.) See Fasolino (A.)	C6-840
Rosenman (I.) See Batallan (F.)	C6-344	Santoro (G.) See Fasolino (A.)	C6-846
(1.0)			

Sapriel (J.) See Jusserand (B.) Sasaki (Y.) and Nishina (Y.)	C6-43	Schröder (U.), Weiss (M.), Reiger (R.) and Page (J.B.) Generalization of Fröhlich's operator for the electron-	
Vibrational spectra of solid and liquid sulfur	C6-557	phonon coupling in polar crystals	C6-496
Sasaki (W.), Sasaki (Y.), Ushioda (S.) and Taylor (W.) Low frequency light scattering		Schröder (U.) See Bauernfeind (W.)	C6-247
spectra of AgI	C6-181	Schröder (U.) See Strauch	C6-707
Sasaki (Y.) See Sasaki (W.)	C6-181	Schwartz (C.) and Schaich (W.L.) Phonons at metal	06 939
Sato (M.), Grier (B.H.), Shapiro (S.M.) and Miyajima (H.) Sof	t	surfaces	C6-828
phonons and magnetic ordering in the γ-phase transition meta	1 C6-770	Selmke (M.) See Falter (C.) Senateur (J.P.) See Khoder	C0-J10
alloys Fe _{1-x} Pd _x		(A.F.)	C6-383
Schaack (G.) See Gerlinger (H.)	C6-499	Servoin (J.L.) See Gervais	C6-415
Schaich (W.L.). See Schwartz	C6-828	Shapiro (S.M.) See Sato (M.).	C6-770
Schanz (G.) See Müller (V.)	C6-389		
Schink (H.J.) See v.Löhneysen (H.)	C6-193	Sharma (P.C.), Roy (K.P.) and Radhakrishnan (V.) The	
Schmeltzer (D.) and Beserman (R.) Anharmonic interaction in zinc selenide	C6-158	effect of non-metal-metal transition due to doping on lattice thermal conducti- vity : applications to Ge-	
Schmeltzer (D.) and Beserman (R.) Renormalization group method for vibrational beha-		Shaw (C.J.) See Rogers (S.J.)	C6-268
vior in mixed crystals	C6-84	Shaw (P.B.) See Young (E.W.).	C6-522
Schmeltzer (D.) See Suzuki (K.)	C6-640		
Schmeltzer (D.) See Teicher (M.)	C6-46	Sheard (F.W.) Spin-lattice interaction in cerium ethyl-sulphate	C6-596
Schmelzer (U.) See Dorner (B.)	C6-602	Sheka (E.F.) See Bielushkin (A.V.)	C6-34
Schmidt (H.J.) See Gompf (F.)	C6-49	Sheka (E.F.) See Bokhenkov (E.L.)	C6-566
Schneck (J.), Toledano (J.C.), Aubree (J.), Joukoff (B.) and		Sheka (E.F.) See Bokhenkov	
Joffrin (C.) Soft phonon valley near the transition to an incommensurate phase		(E.L.)	C6-605
in barium sodium niobate	C6-734	Shekhtman (V.L.) See	06 /20
Scholar (U.D.)		Kaplyanskii (A.A.)	C6-439
Schober (H.R.).— See Buchenau	C6-395	Shekhtman (V.L.) See Kaplyanskii (A.A.)	C6-462

<pre>ihen (T.J.), Castiel (D.) and Maradudin (A.A.) Effect of surface roughness on the Kapitza resistance</pre>	C6-819	Spector (H.N.) and Rynee (T.M.).— Phonon assisted transitions in the free carrier absorption of semi- conductors in quantizing	
Shiren (N.S.). Surface roughness Kapitza conductance: dependence on material properties and phonon frequency	C6-816	Srivastava (G.P.) Phonon conductivity due to non-	C6-670
Sigmund (E.) See Maier (J.)	C6-232	diagonal energy-flux operator	C6-253
Signorelli (G.) See Mazzacurati (V.)	C6-196	Srivastava (G.P.) Role of thermal expansion in the phonon conductivity of solids	06.170
Singh (T.J.) and Verma (G.S.) Four-Phonon processes in solids	C6-155	Stirling (W.G.) See Geick	C6-149
			00,20
Simon (C.) See Batallan (F.).	C6-344	Stolz (H.J.) See Carlone (C.).	C6-323
Simons (A.L.) See Hensel (J.C.)	C6-212	Störmer (H.L.) See Fishman (G.)	C6-305
Sinvani (M.) See Taborek (P.)	C6-852	Strässler (S.) See v.Löhneysen (H.)	C6-193
Sinvani (M.) See Taborek (P.)	C6-855	Strauch (D.) and Schröder (U.)	
Sinvani (M.) See Taborek (P.)	C6-825	On the mode softening in SnTe and PbTe	C6-707
Sladek (R.J.) See Chao (M.H.)	C6-667	Ströbel (B.), Läuger (K.), Bömmel (H.E.) and Müller (V.).	
Sladek (R.J.) See Wu (A.Y.)	C6-646	The ultrasound-induced electric field gradient	
Smith (S.H.) See Daudin (B.).	C6-277	(EFG) in metals	C6-404
Solin (S.A.) Phonons in gra- phite intercalation compounds	C6-283	Strom (U.), Klein (P.B.), Weiser (K.) and Wolf (S.A.) - Time-	
Solin (S.A.) See Hwang (D.M.)	C6-347	resolved phonon spectroscopy of amorphous As ₂ S ₃	C6-30
Solin (S.A.) See Kamitakahara (W.A.)	C6-311	Suchail (R.) See Locatelli (M.)	C6-893
Sood (A.K.), Arora (A.K.), Dattagupta (S.) and Venkataraman (G.) Orienta-		Sugai (S.), Murase (K.), Uchida (S.) and Tanaka (S.). Investigation of the charge density waves in 1T-VSe ₂ by	
tional dynamics of sulphate ions in potash alum	C6-572	Raman scattering	C6-740
Sood (A.K.) See Umadevi (V.).	C6-81	Sugai (S.), Murase (K.), Uchida (S.) and Tanaka (S.). Investigation of the charge	
Sox (D.J.), Rives (J.E.) and Meltzer (R.S.) Stimulated		density waves in 2H-TaSe ₂ by one and two phonons Raman	06 300
emission of tunable high frequency phonons in LaF ₃ :Er ³⁺ .	C6-450	scattering	C6-728
Spector (H.N.) Parametric		Sugai (S.), Ueda (T.) and Murase (K.) Raman scattering in	
amplification of microwave	C6-676	MoS ₂ MoSe ₂ and α-MoTe ₂ at	C6-320

Suski (T.) and Katayama (S.) Phonon induced anomalous resistivity in structural phase transition of PbSnTe	C6-758	Thiry (P.), Maroie (S.), Pireaux (J.J.), Caudano (R.) and Adnot (A.) Adsorbate induced vibrations measured by EELS on a CuZn	
Suzuki (K.), Schmeltzer (D.) and Maradudin (A.A.) Vibrational		α 75/25 (110) surface during its early stage of oxidation	C6-843
properties of vacancies in homopolar semiconductors	C6-640	Thorpe (M.F.) Phonons in amorphous materials	C6-19
Suzuki (H.) See Iwasa (I.)	C6-896	Throwe (J.) and Bron (W.E.) Mode conversion coefficients	
Taborek (P.), Sinvani (M.), Weimer (M.) and Goodstein (D.) Film thickness		of acoustic waves at a crystal-vacuum interface	C6-849
dependence of heat trans- mission into helium	C6-825	Toennies (J.P.).— See Benedek	C6-793
Taborek (P.), Sinvani (M.), Weimer (M.) and Goodstein (D.) Polarization depen-		Tokumura (M.) See Miyasato	C6-658
dence of phonon induced desorption of He atoms at low temepratures	C6-855	Toledano (P.), Pascoli (G.) and Coiret (M.) Simultaneous condensation of several modes at structural phase tran-	
Taborek (P.), Sinvani (M.), Weimer (M.) and Goodstein		sitions	C6-746
(D.) Time constant for phonon induced desorption of Helium	C6-852	Toledano (J.C.). See Moore (D.R.)	C6-785
Takaoka (Y.) See Motizuki	C6-773	Toledano (J.C.) See Schneck	C6-734
Takashige (M.) See Nakamura	C6-418	Tosatti (E.) See Fasolino (A.)	C6-840
Takashige (M.) See Nakamura	C6-421	Tosatti (E.) See Fasolino (A.)	C6-846
Talwar (D.N.) See Zigone		Toudic (B.) See Ecolivet (C.)	C6-779
(M.)	C6-743	Tsai (C.C.) See Nemanich (R.J.)	C6-822
Tanaka (S.) See Sugai (S.)	C6-728	Tsui (D.C.) See Hensel (J.C.).	C6-308
Tanaka (S.) See Sugai (S.)	C6-740	Tsuruoka (F.) and Kajimura (K.) Static polarization echoes in	
	C6-801	metal powders	C6-864
Taylor (W.) See Sasaki (W.).	C6-181	Tua (P.F.) and Mahan (G.D.)	
Teicher (M.), Schmeltzer (D.) and Beserman (R.) Phonon-		Anharmonicity effects in CaF ₂ at low temperature	C6-122
line-shape and disorder correlation in mixed		Tubino (R.) See Mulazzi (E.)	C6-493
GaP _{l-x} As _x	C6-46	.Tucker (J.W.) Phonon scattering at frequencies near the excitor	
Tekippe (V.J.) - See Moore (D.R.)	C6-785	frequency of two-level defects	C6-478
Teuchert (W.D.) See Copley		Uchida (S.) See Sugai (S.)	C6-740
(J.R.D.)	C6-386	Uchida (S.) See Sugai (S.)	C6-728

Ueda (T.) See Sugai (S.)	C6-320	Venkataraman (G.) See Umadevi	06 01
Ulbrich (R.G.), Narayanamurti		(, , , , , , , , , , , , , , , , , , ,	C6-81
(V.) and Chin (M.A.)		Vergés (J.A.) See Pintschovius	
Ballistic transport and decay of near zone-edge non-		(L.)	C6-634
thermal phonons in semi-	C6-226	Verma (G.S.) See Singh (T.J.)	C6-155
Condition	00 -220	Vetelino (J.F.) See Anikiev	
Umadevi (V.), Kesavamoorthy (R.), Sood (A.K.) and		(A.A.)	C6-152
Venkataraman (G.) Surface		Vijayaraghavan (P.R.) See Rao	
polaritons in irradiated		(K.R.)	C6-637
α-quartz	C6-81	11-1 /9 \ 0 01	
Haran (R.C.) - Coo Anihio		Vog1 (P.) See Olego (D.)	C6-652
Umarov (B.S.).— See Anikiev (A.A.)	C6-152	Vogt (H.) See Calleja (J.M.)	C6-487
()	00 132	0 () = ================================	00 407
Umarov (M.) See Gorelik (V.S.)	C6-764		
(7.0) 0 0 141		Wachter (P.) See Celio (M.)	C6-11
Umarov (B.S.) See Gorelik	C6-764	Wachter (P.) See Mörke (I.)	06 17
(00-704	wachter (1.) See Mothe (1.)	C6-14
Unterhorst (E.J.) See Müller		Wada (N.), Klein (M.V.) and Zabel	
(V.)	C6-389	(H.) Raman scattering studies	
11 (7 0) 0 1		of folded shearing phonons in	06 0 0 0
Unterwald (F.C.). See Hensel	C6-212	KC _{12n} (n= 2 ~ 6)	C6-350
(3.0.)	CU-212	Undo (N) . Coo World believe	
Ushioda (S.) See Sasaki (W.)	C6-181	Wada (N.) See Kamitakahara	C6-311
		(00 311
Vacher (R.) and Pelous (J.)		Wagh (A.S.) - A Microscopic	
Phonon damping in crystalline and amorphous solids at		transport theory of electron-	
hypersonic frequences	C6-125	phonon systems	C6-471
Vacher (R.) See Boissier (M.).	C6-611	Harris (W.) Mutachalles (H.) and	
Wallama (A M) - Wigh tomporature		Wagner (M.), Mutscheller (W.) and	
		Nusser (HK.) Phonon and	
Vallera (A.M.) High temperature	C6-398	Nusser (HK.) Phonon and pseudo-magnon transport in coo-	
phonons in iron	C6-398	Nusser (HK.) Phonon and pseudo-magnon transport in cooperative Jahn-Teller systems.	C6-238
phonons in iron	C6-398	pseudo-magnon transport in cooperative Jahn-Teller systems.	C6-238
Van Camp (P.E.), Van Doren (V.E.) and Devreese (J.T.) First	C6-398	pseudo-magnon transport in coo-	C6-238
Van Camp (P.E.), Van Doren (V.E.) and Devreese (J.T.). First principles calculation of the		pseudo-magnon transport in cooperative Jahn-Teller systems. Wagner (R.) See Buchenau (U.).	C6-238 C6-395
Van Camp (P.E.), Van Doren (V.E.) and Devreese (J.T.) First	C6-398	pseudo-magnon transport in cooperative Jahn-Teller systems.	C6-238 C6-395
phonons in iron	C6-625	pseudo-magnon transport in cooperative Jahn-Teller systems. Wagner (R.) See Buchenau (U.).	C6-238 C6-395 C6-807
van Camp (P.E.), Van Doren (V.E.) and Devreese (J.T.). First principles calculation of the phonon spectra of solids		pseudo-magnon transport in cooperative Jahn-Teller systems. Wagner (R.) See Buchenau (U.). Walker (R.P.) See Martin (D.J.) Wallis (R.F.) See Black (J.E.).	C6-238 C6-395 C6-807
phonons in iron	C6-625	pseudo-magnon transport in cooperative Jahn-Teller systems. Wagner (R.) See Buchenau (U.). Walker (R.P.) See Martin (D.J.) Wallis (R.F.) See Black (J.E.). Wallis (R.F.) See Maradudin	C6-238 C6-395 C6-807 C6-837
phonons in iron	C6-625	pseudo-magnon transport in cooperative Jahn-Teller systems. Wagner (R.) See Buchenau (U.). Walker (R.P.) See Martin (D.J.) Wallis (R.F.) See Black (J.E.). Wallis (R.F.) See Maradudin (A.A.)	C6-238 C6-395 C6-807 C6-837
phonons in iron	C6-625 C6-743	pseudo-magnon transport in cooperative Jahn-Teller systems. Wagner (R.) See Buchenau (U.). Walker (R.P.) See Martin (D.J.) Wallis (R.F.) See Black (J.E.). Wallis (R.F.) See Maradudin	C6-238 C6-395 C6-807 C6-837
phonons in iron	C6-625 C6-743	pseudo-magnon transport in cooperative Jahn-Teller systems. Wagner (R.) See Buchenau (U.). Walker (R.P.) See Martin (D.J.) Wallis (R.F.) See Black (J.E.). Wallis (R.F.) See Maradudin (A.A.)	C6-238 C6-395 C6-807 C6-837
phonons in iron	C6-625 C6-743 C6-625	pseudo-magnon transport in cooperative Jahn-Teller systems. Wagner (R.) See Buchenau (U.). Walker (R.P.) See Martin (D.J.) Wallis (R.F.) See Black (J.E.). Wallis (R.F.) See Maradudin (A.A.)	C6-238 C6-395 C6-807 C6-837 C6-137 C6-128
phonons in iron	C6-625 C6-743 C6-625	pseudo-magnon transport in cooperative Jahn-Teller systems. Wagner (R.) See Buchenau (U.). Walker (R.P.) See Martin (D.J.) Wallis (R.F.) See Black (J.E.). Wallis (R.F.) See Maradudin (A.A.)	C6-238 C6-395 C6-807 C6-837
phonons in iron	C6-625 C6-743 C6-625	pseudo-magnon transport in cooperative Jahn-Teller systems. Wagner (R.) See Buchenau (U.). Walker (R.P.) See Martin (D.J.) Wallis (R.F.) See Black (J.E.). Wallis (R.F.) See Maradudin (A.A.)	C6-238 C6-395 C6-807 C6-837 C6-137 C6-128
phonons in iron	C6-625 C6-743 C6-625 C6-560 C6-57	pseudo-magnon transport in cooperative Jahn-Teller systems. Wagner (R.) See Buchenau (U.). Walker (R.P.) See Martin (D.J.) Wallis (R.F.) See Black (J.E.). Wallis (R.F.) See Maradudin (A.A.)	C6-238 C6-395 C6-807 C6-837 C6-137 C6-128
phonons in iron	C6-625 C6-743 C6-625	pseudo-magnon transport in cooperative Jahn-Teller systems. Wagner (R.) See Buchenau (U.). Walker (R.P.) See Martin (D.J.) Wallis (R.F.) See Black (J.E.). Wallis (R.F.) See Maradudin (A.A.)	C6-238 C6-395 C6-807 C6-837 C6-137 C6-128
phonons in iron	C6-625 C6-743 C6-625 C6-560 C6-57	pseudo-magnon transport in cooperative Jahn-Teller systems. Wagner (R.) See Buchenau (U.). Walker (R.P.) See Martin (D.J.) Wallis (R.F.) See Black (J.E.). Wallis (R.F.) See Maradudin (A.A.)	C6-238 C6-395 C6-807 C6-837 C6-137 C6-128

Wasiutynski (T.) Self-consistent phonon calculations in solid nitrogen C6-575	Wu (A.Y.) and Sladek (R.J.) Ultrasonic phonon velocities in Cd _{1-x} Mn _x Te between 1.5 and
Wasiutynski (T.), Natkaniec (I.) and Belushkin (A.I.) Lattice	96 K: anomalies near the magnetic transition C6-646
dynamics calculations for	Wybourne (M.N.) See Challis
solid biphenyl in the high temperature phase	(L.J.)
Webb (W.) See Merlin (R.) C6-392	Yagi (T.), Hidaka (Y.) and Miura (K.) Brillouin scattering study of the incommensurate
Weber (W.) See Harmon (B.N.) C6-628	(Antiferroelectric). Phase transition in sodium nitrite C6-731
Weber (W.) See Lax (M.) C6-161	
Weimer (M.) See Taborek (P.) C6-825	Yeh (C.T.), Reichardt (W.), Renker (B.), Nücker (N.) and
Weimer (M.) See Taborek (P.) C6-855	Loewenhaupt (M.) Lattice dynamics of YAl ₂ and LaAl ₂ - a
Weimer (M.) See Taborek (P.) C6-852	further contribution to the La-Y puzzle C6-371
Weiser (K.) See Strom (U.) C6-30	Yoshida (Y.) See Motizuki (K.) C6-773
	Yoshikawa (S.) and Kimura (T.)
Weiss (K.) Attenuation of	Dipolar field fluctuation
second sound in solids C6-531	effects on phonon echoes in piezoelectric powder C6-873
Weiss (M.) See Schröder (U.) C6-496	
West (A.R.) See Almond (D.P.) C6-187	Young (E.W.) and Shaw (P.B.) A strong coupled acoustic
Wiederick (H.D.) See Rogers (S.J.)	deformation polaron in one dimension
(5:50), (5:50), (5:50)	Zabel (H.) and Magerl (A.)
Wiegmann (W.) See Fishman (G.) C6-305	Phonons in alkali graphite intercalation compounds C6-289
Wigmore (J.K.) See Meredith	Zabel (H.) See Hardcastle
(D.J.)	(S.E.)
Wolf (S.A.) See Strom (V.) C6-30	Zabel (H.) See Magerl (A.) C6-329
Wolfe (J.P.) See Dietsche (W.) C6-447	Zabel (H.) See Wada (N.) C6-350
Wolfe (J.P.) See Greenstein (M.) C6-274	Zdetsis (A.D.) See Northrop (G.A.)
Wolfe (J.P.) See Northrop	Zecchi (E.) - See Locatelli
(G.A.)	(M.)
Wolter (J.) See Horstman	Zeyher (R.) and Dandoloff (R.)
(R-E-)	Entropy of disordered solids at low temperatures C6-40
Won (J.W.) See Kim (J.J.) C6-427	
Woodhead (J.) See Gledhill	Zeyen (C.M.E.) Improved energy resolution with neutron spin
(G.A.)	echo triple-axis spectrometers C6-543
Worlock (J.M.) See Fishman	Zeyen (C.M.E.) See Cailleau
(G.)	(H.)

hang (G.Z.). See Gledhill (G.A.)	Raman scattering spectra in ZnS crystals	C6-743
igone (M.), Vandevyver (M.) and Talwar (D.N.) Pressure	Ziolkiewicz (M.K.) See Brunel (L.C.)	C6-412
dependence of impurity induced	Zwick (A.) See Carles (R.)	C6-105

Third International Conference on Hot Carriers in Semiconductors

Montpellier (France), 1981


Allen Jr. (S.J.), Tsui (D.C.), Derosa (F.), Thornber (K.K.) and Wilson (B.A.).— Direct measurement of velocity over- shoot by hot-electron, sub-		Bimberg (D.), Münzel (H.) and Steckenborn (A.).— Inhibited relaxation of nonresonantly excited electrons in a cold lattice	C7-137
millimeter wave conductivity in Si inversion layers	C7-369	Boggess (T.F.) See Smirl (A.L.)	C7-463
Amand (T.) See Cornet (A.)	C7-471	Boittiaux (B.).— See Constant (E.)	C7-73
Aoki (K.), Kobayashi (T.) and Yamamoto (K.) Periodic oscil- lations and turbulence of hot-		Brousseau (M.) See Cornet (A.)	
carrier plasma at 4.2 K in	C7-51	Brunetti (R.), Jacoboni (C.) and Reggiani (L.) On intervalley	
Asauskas (R.), Balynas (V.),		diffusion of hot-electrons	C7-117
Dobrovolskis (Z.), Krotkus (A.) and Hoerstel (W.) Electron transfer effect in intrinsic		Brunetti (R.) See Jacoboni (C.)	C7-123
tellurium single crystals	C7-329	Brunetti (R.) See Reggiani (L.)	C7-111
Asche (M.), Kostial (H.) and Sarbey (O.G.). The multi- valued distribution of electrons between equivalent	a7 222	Calecki (D.) and Pottier (N.) A convenient gauge treatment of the high electric field elec- tron distribution function	C7-271
valleys	C7-323	Castagné (R.) See Hesto (P.)	C7-177
Balynas (V.) See Asauskas (R.)	C/-329		
Banik (N.C.) See Conwell (E.M.)	C7-315	Castagné (R.) See Hesto (P.) Collet (J.) See Cornet (A.)	C7-235
Bareikis (V.), Galdikas (A.), Miliušytė (R.), Pozhela (J.) and Viktoravičius (V.) Noise and diffusivity of hot elec- trons in n-type InSb	C7-215	Constant (E.) and Boittiaux (B.). Non steady state carrier transport in semiconductor, application to the modelling of submicron devices	-
Barker (J.R.) Quantum physics of retarded transport	C7-245	Constant (E.) See Lippens (D.)	C7-207
Barker (J.R.) and Lowe (D.) Quantum theory of hot electron-		Constant (E.) See Rolland (P.A.)	C7-171
phonon transport in inhomoge- neous semiconductors	c7-293	Conwell (E.M.) and Banik (N.C.) Nonlinear conductivity in	
Bauer (G.) See Schmitte (F.J.)	C7-407	quasi-one-dimensional organic crystals	C7-315
Baumann (K.) See Kocevar (P.).	C7-431	Cornet (A.), Pugnet (M.), Collet (J.), Amand (T.) and Brousseau	
Bensoussan (M.) and Moison (J.M.) Hot electron emission from silicon under pulsed la-	67.170	(M.) Spatial expansion of hot electron-hole plasma at high density in CdSe	C7-471
ser excitation	C7-149		

Dargys (A.) See Dienys (V.) Das (P.), Ferry (D.K.) and Grubin (H.) Hot electron noise properties of semiconductors in	C7-33	Graudszus (W.) and Goebel (E.O.). Picosecond luminescence studies of hot carrier relaxation in pure and highly doped GaAs	C7-437
the non-zero collision duratio	n	VALUE CONTRACTOR CONTR	
regime	C7-227	Grubin (H.L.), Iafrate (G.J.) and Ferry (D.K.) Hot carrier spa-	
Derosa (F.) See Allen Jr. (S.J.)	C7-369	ce and time dependent tran- sients in short channel gallium arsenide devices	
Dienys (V.) and Dargys (A.) Mi- crowave experiments including		Grubin (H.) See Das (P.)	C7-227
avalanche	C7-33	Country (N. Y.) Con Tofonto	
Dobrovolskis (Z.) See Asauskas	a7 222	Grubin (H.L.) See Iafrate	C7-307
(R.)	C7-329	Hashimoto (H.) See Inoue (M.).	C7-19
Dorda (G.) See Leburton (J.P.)	C7-193	Herbert (D.C.) and Till (S.J.)	
Eastman (L.F.) Experimental studies of ballistic trans-		Field dependent scattering and hot electron kinetics	C7-277
port in semiconductors	C7-263	4	
Esposito (F.P.) See Jauho		Hess (K.) Lateral transport in superlattices	C7-3
(A.P.)	C7-301	Hess (K.) See Tang (J.Y.)	C7-63
Ferry (D.K.) Transport in sub-			
micron devices	C7 -2 53	Hesto (P.), Pône (J.F.) and Castagné (R.). Unidimensional	
Ferry (D.K.) See Das (P.)	C7-227	particle model of conduction in N type GaAs at low tempera-	
Ferry (D.K.) See Grubin (H.L.)	C7-201	ture : contacts influence	C7-177
Ferry (D.K.) See Infrate		Hesto (P.), Vaissière (J.C.),	
(G.J.)	C7-307	Gasquet (D.), Castagné (R.) and Nougier (J.P.) A sugges-	
Ferry (D.K.) See Lugli (P.)	C7-103	tion of noise experiment for showing ballistic transport	C7-235
Ferry (D.K.) See Zimmermann			
(J.)	C7-95	Hida (H.) See Inoue (M.)	C7-19
Friscourt (M.R.).— See Rolland	C7-171	Hill (G.) and Robson (P.N.) Hot electron transport in the Ga _{1-x} Al _x As system	C7-335
Fujii (K.) See Otsuka (E.)	C7-393	A 6	
Galdikas (A.) See Bareikis		Hiyamizu (S.) See Inoue (M.)	C7-19
(V.)	°C7-215	Hoerstel (W.) See Asauskas	C7-329
Gasquet (D.) See Hesto (P.)	C7-235		0, 32)
Gasquet (D.) See Nougier	C7-283	Honlein (W.) and Landwehr (G.) Energy loss of warm electrons	
		at the interface of (100) silicon MOSFETs	C7-25
Gasquet (D.). See Reggiani (L.)	C7-357	lafrate (G.J.), Grubin (H.L.) and	
Goebel (E.O.) See Graudszus		Ferry (D.K.) Utilization of	
(W.)	C7-437	quantum distribution functions for ultra-submicron device	
Gornik (E.) See Lindemann (G.)	C7-399	transport	C7-307

Iafrate (G.I.). See Grubin (H.L.)	C7-201	Lecoy (G.) See Touboul (A.)	C7-183
Iafrate (G.J.) See Tang (J.Y.)	C7-63	Lindemann (G.) and Gornik (E.) Non Maxwellian electron dis- tribution function in GaAs	C7-399
Inoue (M.), Hiyamizu (S.), Hida (H.), Hashimoto (H.) and I Inuishi (Y.). Hot electron effects in a 2D electron gas at the GaAs/AlGaAs interface.	C7-19	Lippens (D.) and Constant (E.).— A semiparticular simulation to study high frequency beha- viour of hot carriers in IMPATT devices	C7-207
Inuishi (Y.) See Inoue (M.)	C7-19	Lopez (J.C.) See Touboul (A.).	
Ishikawa (T.) See Sawaki (N.).	C7-221	Lowe (D.) See Barker (J.R.)	C7-183
Jacoboni (C.), Reggiani (L.) and Brunetti (R.). Frequency and wavevector dependent diffusion coefficient of electrons from Monte Carlo calculations	C7-123	Lugli (P.), Zimmermann (J.) and Ferry (D.K.) Non-equilibrium hot-carrier diffusion phenomenon in semiconductors. II. An experimental Monte Carlo	0, 2,3
Jacoboni (C.) See Brunetti (R.)	C7-117	approach	C7-103
Jacoboni (C.) See Reggiani		Lugli (P.) See Zimmermann (J.)	C/-95
(L.) Jauho (A.P.), Wilkins (J.W.) and Esposito (F.P.) A non- pertubative Green's function	C7-111	Maksym (P.A.) The effect of electron-electron scattering on hot photoexcited electrons in GaAs	C7-143
technique for calculations of non-linear transport pro- perties	c7-301	Miliušytė (R.) See Bareikis (V.)	C7-215
Kobayashi (T.) See Aoki (K.)	C7-51	Miller (A.) See Smirl (A.L.)	C7-463
Kocevar (P.), Baumann (K.), Vogl (P.) and Potz (W.). Theory of one- and two-phonon defor- mation potentials in semicon-		Moison (J.M.) See Bensoussan (M.) Morohashi (M.) See Sawaki (N.)	C7-149 C7-221
ductors	C7-431	Münzel (H.) See Bimberg (D.)	C7-137
Komiyama (S.) and Spies (R.) Hot carrier population inver-		Nishinaga (T.) See Sawaki (N.)	C7-221
sion in p-Ge	C7-387	Nougier (J.P.), Vaissière (J.C.) and Gasquet (D.) Duration of	
Kostial (H.) See Asche (M.)	C7-323	collisions in semiconductors.	C7-283
Kozlov (V.A.) Hot electron po- pulation inversion and bulk		Nougier (J.P.) See Hesto (P.).	C7-235
NDC in semiconductors	C7-413	Nougier (J.P.) See Reggiani	C7-357
Krotkus (A.) See Asauskas (R.)	C7-329	Ohyama (T.) See Otsuka (E.)	C7-393
Kurosawa (T.) Peculiar hot electron effects in crossed electric and magnetic fields.	C7-377	Otsuka (E.), Ohyama (T.) and Fujii (K.) Far-infrared cyclotron resonance of photoexcited hot	L
Landwehr (G.) See Honlein (W.)	C7-25	carriers in indium antimonide	C7-393
Leburton (J.P.) and Dorda (G.) The v-E relation and the field		Perryman (G.P.) See Smirl (A.L.)	C7-463
distribution in submicron MOSFET's	C7-193	Pone (J.F.) See Hesto (P.)	C7-177

Pottier (N.) See Caleki (D.) C7-2	
	carrier relaxation with pico-
Pötz (W.) See Kocevar (P.) C7-4	31 second laser pulses C7-445
Pozhela (J.) See Barakeis (V.) C7-2	15 Shichijo (H.) See Tang (J.Y.). C7-63
Pugnet (M.) See Cornet (A.) C7-4	71 Smirl (A.L.), Miller (A.), Perryman (G.P.) and Boggess
Rees (H.D.) Hot electron devices for millimetre and submillime-	(T.F.) Picosecond interband saturation and intraband re-
tre applications C7-1	1 laxation of photoexcited car- riers in germanium C7-463
Reggiani (L.), Brunetti (R.) and Jacoboni (C.) The method of moments for the analysis of	Spies (R.) See Komiyama (S.) C7-387
transient hot-carrier pheno-	Steckenborn (A.).— See Bimberg (D.)
mena C7-1	
Reggiani (L.), Vaissière (J.C.), Nougier (J.P.) and Gasquet	Tauc (J.) See Vardeny (Z.) C7-477
(D.) Transient regimes of hot carriers in p-type silicon C7-3!	structure dependent impact io-
Reggiani (L.) See Brunetti (R.)	nization in silicon and gallium arsenide
Reggiani (L.) See Jacoboni (C.)	Thornber (K.K.). See Allen Jr (S.J.)
Robson (P.N.) See Hill (G.) C7-3	Till (S.J.) See Herbert (D.C.) C7-277
Rolland (P.A.), Friscourt (M.R.), Salmer (G.) and Constant (E.).— Theoretical study of 100 GHz GaAs transferred-electron de- vices	Touboul (A.), Lopez (J.C.) and Lecoy (G.) Hot carriers in reduced geometry surface-chan- nel charge-coupled devices C7-183
Rosencher (E.) Ballistic trans-	Tsui (D.C.) See Allen Jr. (S.J.) C7-369
port in semiconductors : a Displaced Maxwellian formula-	Ulbrich (R.G.) Electron-phonon
tion	phonon transport and decay C7-423
Sarbey (0.G.) See Asche (M.) C7-3	Vaigaiàre (I.C.) - Con Hooks
Sawaki (N.), Ishikawa (T.),	
Morohashi (M.) and Nishinaga (T.). Hot holes and noise in	Vaissière (J.C.) See Nougier (J.P.)
space charge limited current flow in p-type silicon C7-2	21 Vaissière (J.C.) See Reggiani (L.) C7-357
Schmitte (J.F.), Bauer (G.) and Zawadzki (W.) — Magnetophonon oscillations in the lifetime of hot photoexcited carriers. C7-4	Van de Roer (Th.G.) See Van Someren Greve (S.C.) C7-343
Schöll (E.) An impact ioniza- tion model for current control- led negative resistance and	Van Someren Greve (S.C.) and Van de Roer (Th.G.) Time- dependent carrier velocities in III-V compounds calculated
g-r induced nonequilibrium phase transitions C7-5	by the Legendre-polynomial iterative method C7-343

tical picosecond studies of hot carriers in amorphous	Wilson (B.A.).— See Allen Jr. (S.J.)	C7-369
semiconductors C7-477	Yamamoto (K.) See Aoki (K.)	C7-51
iktoravičius (V.) See Bareikis (V.) C7-215	Zawadzki (W.) See Schmitte (F.J.)	C7-407
ogl (P.) See Kocevar (P.) C7-431	Zimmermann (J.), Lugli (P.) and Ferry (D.K.) Non-equilibrium	
autelet (M.) Diffusion of elec- trons and holes in doped semi- conductors at high lattice and electronic temperatures C7-131	hot-carrier diffusion phenomenon in semiconductors. I. A theoretical non-Markovian approach	C7-95
filkins (J.W.) See Jauho	Zimmermann (J.) See Lugli (P.)	C7-103



Troisième Symposium sur les Etalons de Fréquence et la Métrologie

Aussois (France), 1981

Alekseev (V.A.). The effect of collisions in precision spectroscopy	n	Basov (N.G.), Gubin (M.A.), Nikitin (V.V.), Nikulchin (A.V.), Tyurikov (D.A.), Petrovskiy (V.N.) and Protscenko (E.D.).— A highly sensitive technique of de- tecting supernarrow spectral lines based on the frequency resonances of a double—mode laser	C8-89
Alley (C.O.) Voir Allan (D.W.)	C8-395	Bassi (D.) Voir Bordé (Ch.J.)	C8-15
André (J.), Teboul (A.), Védel (F.) and Védel (M.) A simplified model for computing temperature of a dilute population of laser cooled stored		Bava (E.), Godone (A.) and De Marc (A.) Phase-coherent frequency synthesis suitable for direct frequency measurements up to 30 THz	
ions	C8-315	Becker (W.), Blatt (R.) and	00 40.7
Arditi (M.) Microwave Ramsey pattern in a laser pumped cesium beam. Application to		Werth (G.) Precise determination of ¹³⁵ Ba+ and ¹³⁷ Ba+ hyperfine structure	C8-339
velocity distribution and 2 nd order Doppler - shift		Bergquist (J.C.) Voir Lewis	
in frequency standard	C8-261	(L.L.)	C8-271
Ashby (N.) Voir Allan (D.W.)	C8-395	Bergquist (J.C.) Voir Wineland (D.J.)	C8-307
Audoin (C.), Viennet (J.) and Lesage (P.) Hydrogen maser: active or passive ?	C8-159	Besson (R.J.) New advances in precision quartz oscillators and their impact on quantum	
Avrillier (S.) Voir Bordé (Ch.J.)	C8-15	standards	C8-199
Baer (T.) Voir Hall (J.L.)	C8-59	Billing (H.) Voir Rüdiger (A.)	C8-451
Bagayev (S.N.), Chebotayev		Blatt (R.) Voir Becker (W.)	C8-339
(V.P.), Dychkov (A.S.) and Maltsev (S.V.).— Supernarrow resonances in methane on E-line of the P(7) transition of the v ₃ band and their application in optical frequency standards	C8-21	Bordé (Ch.J.), Avrillier (S.), Van Lerberghe (A.), Salomon (Ch.), Bassi (D.) and Scoles (G.).— Observation of optical Ramsey fringes in the 10 µm spectral region using a super- sonic beam of SF ₆	C8-15
Baird (K.M.) Optical frequency		Bordé (Ch.J.) Voir Salomon	
measurement at NRC and progress towards frequency measurement	of	(Ch.)	C8-127
visible light	C8-485	Bordé (Ch.J.) Voir Clairon (A.)	00 127
Barger (R.L.) Voir Salomon (Ch.)	C8-3	Boulanger (J.S.) Voir Mungall (A.G.)	C8-233


Bréant (Ch.) Voir Clairon (A.)	C8-127	Drullinger (R.E.) Voir Wineland (D.J.)	C8-307
Bréant (Ch.) Voir Salomon (Ch.)	C8-3	Dychkov (A.S.) Voir Bagayev	
Brillet (A.) and Cérez (P.) Laser frequency stabilization		(S.N.)	C8-21
by saturated absorption	C8-73	Evenson (K.M.), Jennings (D.A.) and Petersen (F.R.).— The	
Brisson (A.) Voir Vanier (J.)	C8-139	frequency measurement of visible light	C8-473
Busca (G.) Voir Thomann (P.).	C8-189	Ezekiel (S.) Voir Peuse (B.W.)	C8-53
Camy (G.) Voir Clairon (A.)	C8-127	Ezekiel (S.) Voir Tench (R.E.)	C8-45
Cérez (P.) Voir Brillet (A.).	C8-73	Feldman (M.) Voir Lewis (L.L.)	C8-271
Chebotayev (V.P.) Optical time scale	C8-505	Gagnepain (J.J.), Théobald (G.) and Uebersfeld (J.) Analysis	
Chebotayev (V.P.) Voir Bagayev (S.N.)	C8-21	of 1/f ² and 1/f frequency noises in quartz resonators.	C8-201
Clairon (A.), Van Lerberghe (A.), Bréant (Ch.), Salomon (Ch.), Camy (G.) and Bordé		Garvey (R.M.) Approaches to the realization of improved clock performance in cesium based	
(Ch.J.) A new absolute frequency reference grid in the		devices	C8-185
28 THz range	C8-127	Giacomo (P.) Prospects for the new definition of the metre.	C8-513
Costain (C.C.) Voir Mungall (A.G.)	C8-233	Glaze (D.), Gray (J.), Levine (J.), Stein (S.) and Walls	
Couillaud (B.) Generation of CW coherent radiation in the near U.V	C8-115	(F.) New NBS time scale hardware	C8-427
Crampton (S.B.), Krupczak (J.J.)	00-113	Glaze (D.J.) Voir Lewis (L.L.)	C8-241
and Souza (S.P.) Progress of the state-selected beam		Godone (A.) Voir Bava (E.)	C8-463
low temeprature hydrogen maser	C8-181	Goy (P.) Voir Raimond (J.M.).	C8-37
Daams (H.) Voir Mungall		Gubin (M.A.) Voir Basov (N.G.)	C8-89
(A.G.)	C8-233	Guinot (B.) Status of the International Atomic Time	
Decher (R.) Voir Allan (D.W.)	C8-395	TAI	C8-219
Dehmelt (H.) Coherent spectros- copy on single atomic system		Gray (J.) Voir Glaze (D.)	C8-427
at rest in free space II	C8-299	Hall (J.L.), Hollberg (L.), Ma Long-sheng, Baer (T.)	
De Marchi (A.) Improved performances in Cs standards with		and Robinson (H.G.) Pro- gress toward phase-stable	
new-design resonators	C8-289	optical frequency standards.	C8-59
De Marchi (A.) Voir Bava (E.) Desaintfuscien (M.) Voir	C8-463	Hardy (W.N.) and Morrow (M.) Prospects for low tempera-	
Jardino (M.)	C8-327	ture H masers using liquid helium coated walls	C8-171
Dorenwendt (K.) Development and		Haroche (S.) Voir Raimond	
performance of primary standard	ds - C8-229	(J.M.)	C8-37

<pre>layashi (R.) Voir Nakagi (K.) lemmati (H.) Voir Winela</pre>	C8-253	Lewis (L.L.), Feldman (M.) and Bergquist (J.C.) Impact of lasers on primary frequency	
(D.J.)		standards and precision spec- troscopy	C8-271
emmer (P.R.) Voir Tench		Lewis (L.L.), Walls (F.L.) and Glaze (D.J.) Design considerations and performance of	
bllberg (L.) Voir Hall (J.L.)	C8-59	NBS-6, the NBS primary frequen standard	cy C8-241
owe (D.A.) Voir Walls ((F.L.) C8-151	Lewis (L.L.) Voir Itano (W.M.)	C8-283
tano (W.M.), Lewis (L.L.) Wineland (D.J.). Shift ² S _k hyperfine splitting	of	Ma Long-sheng Voir Hall (J.L.)	C8-59
due to blackbody radiat and its influence on fr quency standards	re-	Maischberger (K.) Voir Rüdiger (A.)	C8-451
tano (W.M.) Voir Winela		Maltsev (S.V.) Voir Bagayev (S.N.)	C8-21
ardino (M.), Desaintfusci		Minogin (V.G.) Voir Letokhov (V.S.)	C8-347
(M.) and Plumelle (F.). Prospects for a mercury	,-	Moreau (R.) The impact of very	
ion frequency standard.	C8-327	stable oscillators on navi- gation systems	C8-415
ennings (D.A.) Voir Eve		Morrow (M.) Voir Hardy (W.N.)	C8-171
night (D.J.E.) Optical quency synthesis at NPI	C8-495	Mungall (A.G.); Daams (H.), Boulanger (J.S.) and Costain (C.C.) Performance of the	
loga (Y.), Nakadan (Y.) ar Yoda (J.) The cesium quency standard NRLM-I	beam fre-	NRC primary time standards CsV, CsVI A, CsVI B, CsVI C.	C8-233
irupczak (J.J.) Voir Cra		Nakadan (Y.) Voir Koga (Y.)	C8-247
(S.B.)		Nakagiri (K.), Shibuki (M.), Urabe (S.), Hayashi (R.) and Saburi (Y.) Cesium beam	
(uhnle (P.F.) and Sydnor (Present and future free and timing capabilities	quency	frequency standard at the Radio Research Laboratories.	C8-253
deep space network		Nikitin (V.V.) Voir Basov	
Kunski (R.) Voir Vanier	(J.). C8-139	(N.G.)	C8-89
eiby Jr. (C.C.) Voir Te		Nikulchin (A.V.) Voir Basov (N.G.)	C8-89
esage (P.) Voir Audoin	(C.). C8-159	Ohtsu (M.) Voir Tako (T.)	C8-83
etokhov (V.S.) and Minogi		Paulin (P.) Voir Vanier (J.).	C8-139
(V.G.) Laser cooling atoms and its applicati in frequency standards.	lon	Petersen (F.R.) Voir Evenson (K.M.)	C8-473
		Petrovskiy (V.N.) Voir Basov (N.G.)	C8-89
Taraban (T) Wada Class	(D) C8-427		

Peuse (B.W.), Prenties (M.G.)		Shibuki (M.) voir Nakagiri	
and Ezekiel (S.) Distor-		(K.)	C8-253
tion in atomic beam absorption			
lineshapes	C8-53	Souza (S.P.) Voir Crampton	
1		(S.B.)	C8-181
Peuse (B.W.) Voir Tench (R.E.)	C8-45		
rease (bowe). Volt lenen (Male)	00 15	Stein (S.) Voir Glaze (D.)	C8-427
		000211 (00)0 1022 1222 (120)	
Picard (R.H.) Voir Tench (R.E.)	C8-45	Strumia (F.) Frequency control	
		and frequency modulation of	
Plumelle (F.) Voir Jardino (M.)	C8-327	far infrared laser by Stark	
			C8-105
Prenties (M.G.) Voir Peuse		effect	00-103
(B.W.)	C8-53		
		Sydnor (R.L.) Voir Kuhnle	00 073
Protscenko (E.D.) Voir Basov		(P.F.)	C8-373
(N.G.)	C8-89		
(Tako (T.), Ohtsu (M.) and	
Raimond (J.M.), Goy (P.),		Tsuchida (H.) Frequency	
Vitrant (G.) and Haroche (S.).		stabilization of AlGaAs	
Millimeter-wave spectroscopy		lasers based on the H ₂ O and	
of cesium Rydberg states and		Rb-D ₂ lines	C8-83
, ,		2	
possible applications to fre-	00 27	Teboul (A.) Voir André (J.)	C8-315
quency metrology	C8-37		
		Tench (R.E.), Peuse (B.W.),	
Reinhardt (V.S.) A statistical		Hemmer (P.R.), Thomas (J.E.),	
model for flicker noise	C8-211	Ezekiel (S.), Leiby Jr. (C.C.),	
		Picard (R.H.) and Willis	•
Ritter (R.C.) Macroscopic rotors	3		
and gravitational effects	C8-429	(C.R.) - Two laser Raman	
		difference technique applied	
Robinson (H.G.) Voir Hall		to high precision spectros-	00 / 5
(J.L.)	C8-59	copy	C8-45
Rüdiger (A.), Schilling (R.),		Théobald (G.) Voir Gagnepain	2000
Schnupp (L.), Winkler (W.),		(J.J.)	C8-201
Billing (H.), Maischberger			
(K.) Demands on laser		Thomann (P.) and Busca (G.)	
purity in an interferometric		Modulation effects in passive	
gravitational wave detector	C8-451	Rb frequency standards	C8-189
		Thomas (J.E.) Voir Tench (R.F.)	C8-45
Saburi (Y.) Voir Nakagiri (K.)	C8-253		
		Tourrenc (Ph.) Gravitational	
Salomon (Ch.), Bréant (Ch.), Borde	5	experiments with stable electro) -
(Ch.J.) and Barger (R.L.)		magnetic devices	C8-441
Ramsey fringes using transi-			
tions in the visible and 10 µm		Tsuchida (H.) Voir Tako (T.).	C8-83
spectral regions : experiments	a1		
methods	C8-3	Tyurikov (D.A.) - Voir Basov	١ .
		(N.G.)	C8-89
Salomon (Ch.) Voir Bordé			
(Ch.J.)	C8-15	Uebersfeld (J.) Voir Gagnepain	
		(J.J.)	C8-201
Salomon (Ch.) Voir Clairon (A.)	C8-127		
(114)		Urabe (S.) Voir Nakagiri (K.)	C8-253
Schilling (R.) Voir Rüdiger		(1)	
(A.)	C8-451	Vanier (J.), Kunski (R.), Brisson	1
(, , , , , , , , , , , , , , , , , ,	30 431	(A.) and Paulin (P.) Pro-	
Schnupp (I.) - Wody Dildion			
Schnupp (L.) Voir Rüdiger	CO / 51	gress and prospects in rubi-	CO_120
(A.)	C8-451	dium frequency standards	C8-139
Cooles (C) - Water Break		Wan Jarharaha (A) Water Barrie	
Scoles (G.) Voir Bordé		Van Lerberghe (A.) Voir Bordé	- 1.
			CO 1 F

Van Lerberghe (A.).- Voir Clairon Werth (G.).- Voir Becker (W.).. C8-339

Vedel (F.) Voir André (J.)	C8-127	Willenberg (G.D.) Voir Weiss (C.O.)	C8-97
Vedel (M.) Voir André (J.)	C8-315	Willis (C.R.) Voir Tench (R.E.)	C8-45
Vessot (R.F.C.) Gravitation and relativity experiments using atomic clocks	C8-359	Wineland (D.J.), Bergquist (J.C.), Drullinger (R.E.), Hemmati (H.), Itano (W.M.) and Walls (F.L.) Laser cooled, stored ion experi- ments at NBS and possible	
Vessot (R.F.C.) Voir Allan	C8-395		
Viennet (J.) Voir Audoin (C.)	C8-159	applications to microwave and optical frequency standards.	C8-307
Vitrant (G.) Voir Raimond (J.M.)	C8-37	Wineland (D.J.) Voir Itano (W.M.)	C8-283
Walls (F.L.) and Howe (D.A.) Timekeeping potentials using passive hydrogen masers	C8-151	Winkler (G.M.R.) Voir Allan (D.W.)	C8-395
Walls (F.) Voir Glaze (D.)	C8-427	Winkler (W.) Voir Rüdiger (A.)	C8-451
Walls (F.L.) Voir Lewis (L.L.)	C8-241	Yang Xiaoren Works on Chinese primary frequency standards.	C8-257
Walls (F.L.) Voir Wineland (D.J.)	C8-307	Yannoni (N.F.) Clock require- ments for navigation, commu-	
Weiss (C.O.) and Willenberg (G.D.) Generation of cw FIR radiation using 2- and 3- pho-		nication and identification systems	C8-383
ton interactions	C8-97	Yoda (J.) Voir Koga (Y.)	C8-247



Commission paritaire Nº 26.741

Editions de Physique 1982

Directrice de la Publication : Jeanne BERGER

Imprimé en France. — Imprimerie JOUVE, 18, rue Saint-Denis, 75001 PARIS N° 10036, Dépôt légal : Mai 1982 to be of more polesters.

A STATE OF THE PARTY OF THE PAR

THE RESIDENCE OF STREET

COLUMN TO SERVICE STATE OF THE SERVICE STATE OF THE

